

## 1. NAME OF THE MEDICINAL PRODUCT

**Sativex®**

Oromucosal Spray.

## 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each 100 microlitre spray contains:

2.7 mg delta-9-tetrahydrocannabinol (THC) and 2.5 mg cannabidiol (CBD).

Each ml contains:

38-44 mg and 35-42 mg of two extracts (as soft extracts) from *Cannabis sativa* L., folium cum flore (Cannabis leaf and flower) corresponding to 27 mg delta-9-tetrahydrocannabinol and 25 mg cannabidiol.

Extraction solvent: Liquid carbon dioxide.

Excipient(s) with known effect: each 100 microlitre spray also contains up to 0.04 g ethanol.

For the full list of excipients, see section 6.1.

## 3. PHARMACEUTICAL FORM

Oromucosal spray.

A yellow/brown solution in a spray container.

## 4. CLINICAL PARTICULARS

### 4.1 Therapeutic indications

Sativex is indicated as add-on treatment for symptom relief in patients with moderate to severe spasticity due to multiple sclerosis (MS) who have not responded adequately to other medication and who demonstrate at least 20% improvement in spasticity related symptoms during a four week trial of therapy.

Sativex is indicated as adjunctive treatment for the symptomatic relief of neuropathic pain in multiple sclerosis (MS) in adults.

The physician who elects to use Sativex for extended periods should periodically re-evaluate the long-term usefulness of Sativex for the individual patient.

Sativex may be useful as adjunctive analgesic treatment in adult patients with advanced cancer who experience moderate to severe pain during the highest tolerated dose of strong opioid therapy for persistent background pain.

### 4.2 Posology and method of administration

Sativex is for oromucosal use only.

Sativex is intended to be used in addition to the patient's current anti-spasticity medication.

Treatment must be initiated and supervised by a physician with specialist expertise in treating this patient population.

**Adults:**

The spray container should be shaken before use and the spray should be directed at different sites on the oromucosal surface changing the application site each time the product is used.

Patients should be advised that it might take up to two weeks to find the optimal dose and that undesirable effects can occur during this time, most commonly dizziness. These undesirable effects are usually mild and resolve in a few days. However, physicians should consider maintaining the current dose, reducing the dose or interrupting, at least temporarily, the treatment depending on seriousness and intensity.

To minimise variability of bioavailability in the individual patient, administration of Sativex should be standardised as far as possible in relation to food intake (see section 4.5). In addition, starting or stopping some concomitant medicinal products may require a new dose titration (see section 4.5).

**Titration period:**

A titration period is required to reach optimal dose. The number and timing of sprays will vary between patients.

The number of sprays should be increased each day following the pattern given in the table below. The afternoon/evening dose should be taken at any time between 4 pm and bedtime. When the morning dose is introduced, it should be taken at any time between waking and midday. The patient may continue to gradually increase the dose by one spray per day, up to a maximum of 12 sprays per day, until they achieve optimum symptom relief. There should be at least a 15 minute gap between sprays.

Day	Number of sprays in the morning	Number of sprays in the evening	(Total number of sprays per day)
1	0	1	1
2	0	1	1
3	0	2	2
4	0	2	2
5	1	2	3
6	1	3	4
7	1	4	5
8	2	4	6
9	2	5	7
10	3	5	8
11	3	6	9
12	4	6	10
13	4	7	11
14	5	7	12

**Maintenance period:**

Following the titration period, patients are advised to maintain the optimum dose achieved. The median dose in clinical trials for patients with multiple sclerosis is eight sprays per day. Once the optimum dose has been achieved, patients may spread the doses throughout the day according to individual response and tolerability. Re-titration upwards or downwards may be appropriate if there are any changes in the severity of the patient's condition, changes in their concomitant medication or if troublesome adverse reactions develop. Doses of greater than 12 sprays per day are not recommended.

#### Review by the physician - patients with spasticity in MS

A thorough evaluation of the severity of spasticity related symptoms and of the response to standard anti-spasticity medication should be performed prior to initiation of treatment. Sativex is only indicated in patients with moderate to severe spasticity that have responded inadequately to other anti-spasticity medication. The patient's response to Sativex should be reviewed after four weeks of treatment. If a clinically significant improvement in spasticity related symptoms is not seen during this initial trial of therapy, then treatment should be stopped. In the clinical trials this was defined as at least a 20% improvement in spasticity related symptoms on a 0-10 patient reported numeric rating scale (see section 5.1). The value of long term treatment should be re-evaluated periodically.

#### **Children**

Sativex is not recommended for use in children or adolescents below 18 years of age due to lack of safety and efficacy data.

#### **Elderly**

No specific studies have been carried out in elderly patients, although patients up to 90 years of age have been included in clinical trials. However, as elderly patients may be more prone to develop some CNS adverse reactions, care should be taken in terms of personal safety such as preparation of hot food and drinks.

#### **Patients with significant hepatic or renal impairment**

There are no studies in patients with impaired hepatic or renal function. However, in these sub-populations the effects of Sativex may be exaggerated or prolonged. Frequent clinical evaluation by a clinician is recommended in these patient populations (see section 4.4).

### **4.3 Contraindications**

Sativex is contraindicated in patients:

- With hypersensitivity to cannabinoids or to any of the excipients listed in section 6.1.
- With any known or suspected history or family history of schizophrenia, or other psychotic illness; history of severe personality disorder or other significant psychiatric disorder other than depression associated with their underlying condition.
- Who are breast feeding (in view of the considerable levels of cannabinoids likely in maternal breast milk and the potential adverse developmental effects in infants).

### **4.4 Special warnings and precautions for use**

Mild or moderate dizziness is commonly reported. This most frequently occurs in the first few weeks of treatment.

Sativex is not recommended for use in children or adolescents below 18 years of age due to lack of safety and efficacy data.

Alterations in pulse rate and blood pressure have been observed following initial dose introduction so caution during initial dose titration is essential. Fainting episodes have been observed with use of Sativex. Use of Sativex is not recommended in patients with serious cardiovascular disease. However, following dosing in healthy volunteers with Sativex up to 18 sprays twice daily, there were no clinically relevant changes in QTc, PR or QRS interval duration, heart rate, or blood pressure.

Until further information is available, caution should be taken when treating patients with a history of epilepsy, or recurrent seizures.

Psychiatric symptoms such as anxiety, illusions, changes in mood, and paranoid ideas have been reported during treatment with Sativex. These are likely to be the result of transient CNS effects and are generally mild to moderate in severity and well tolerated. They can be expected to remit on reduction or interruption of Sativex medication.

Disorientation (or confusion), hallucinations and delusional beliefs or transient psychotic reactions have also been reported and in a few cases a causal association between Sativex administration and suicidal ideation could not be ruled out. In any of these circumstances, Sativex should be stopped immediately and the patient monitored until the symptom has completely resolved.

No specific studies have been carried out in patients with significant hepatic or renal impairment. THC and CBD are metabolised in the liver, and approximately one third of the parent drugs and their metabolites are excreted in the urine (the remainder via the faeces). Several THC metabolites may be psychoactive. Thus, the systemic exposure and the effects of Sativex are dependent on both renal and hepatic function and in patients with significant impaired hepatic or renal function; the effects of Sativex may be exaggerated or prolonged. Frequent clinical evaluation by a clinician is recommended in these patient populations.

Sativex contains approximately 50% v/v of ethanol. Each actuation contains up to 0.04g of ethanol. A small glass of wine (125 mL) of nominal ethanol content 12% v/v would contain approximately 12g ethanol. Most patients respond at doses up to and including 12 sprays a day which would contain less than 0.5 g of ethanol.

There is a risk of an increase in incidence of falls in patients whose spasticity has been reduced and whose muscle strength is insufficient to maintain posture or gait. In addition to an increased risk of falls, the CNS adverse reactions of Sativex, particularly in elderly patients, could potentially have an impact on various aspects of personal safety, such as with food and hot drink preparation.

Although there is a theoretical risk that there may be an additive effect with muscle-relaxing agents such as baclofen and benzodiazepines, thereby increasing the risk of falls, this has not been seen in clinical trials with Sativex. However, patients should be warned of this possibility.

Although no effect has been seen on fertility, independent research in animals found that cannabinoids affected spermatogenesis. Female patients of child-bearing potential and male patients with a partner of childbearing potential should ensure that reliable contraceptive precautions are maintained for the duration of therapy and for three months after discontinuation of therapy (see section 4.6).

Patients who have a history of substance abuse, may be more prone to abuse Sativex as well (see section 5.1).

The abrupt withdrawal of long-term Sativex treatment has not resulted in a consistent pattern or time-profile of withdrawal-type symptoms and the likely consequence will be limited to transient disturbances of sleep, emotion or appetite in some patients. No increase in daily dosage has been observed in long-term use, and patient self-reported levels of 'intoxication' are low. For these reasons, dependence on Sativex is unlikely.

Adverse reactions have been reported which could be associated with the route of administration of the medicine. Application site type reactions consisted of mainly mild to moderate stinging at the time of application. Common application site reactions include application site pain, oral pain and discomfort, dysgeusia, mouth ulceration and glossodynia. Two cases of possible leukoplakia were observed but neither was confirmed histologically; a third case was unrelated. In view of this, patients who observe discomfort or ulceration at the site of application of the medicine are advised to vary the site of application within the mouth and should not continue spraying onto sore or inflamed mucous membrane. Regular inspection of the oral mucosa is also advised in long-term administration. If lesions or persistent soreness are observed, medication should be interrupted until complete resolution occurs.

Patients should be advised that if they travel to another country it may not be legal for them to take this medicine into some countries. They should be encouraged to check the legal status before travelling with Sativex.

Pregnancy and lactation: refer to section 4.6.

#### **4.5 Interaction with other medicinal products and other forms of interaction**

The two main components of Sativex, delta-9-tetrahydrocannabinol (THC) and cannabidiol (CBD) are metabolised by the cytochrome P<sub>450</sub> enzyme system.

The *in vitro* inhibitory effects of Sativex on the major CYP450 enzymes CYP3A4 and CYP2C19 occurs at concentrations substantially higher than the maximum observed in clinical trials. No interactions with regards to at risk-CYP3A4 substrates are expected.

In an *in vitro* study with 1:1% (v/v) THC botanical drug substance (BDS) and CBD BDS, no relevant induction of cytochrome P<sub>450</sub> enzymes was seen for CYP1A2, CYP2C9, CYP2C19 and CYP3A4 enzymes in human hepatocytes, at doses of up to 1 µM (314 ng/mL).

Concomitant treatment with the CYP3A4 inhibitor ketoconazole produced an increase in C<sub>max</sub> and AUC of THC (1.2- and 1.8-fold, respectively), its primary metabolite (3- and 3.6-fold, respectively) and of CBD (2- and 2-fold, respectively). Therefore, if concomitant drug treatment with CYP3A4 inhibitors (e.g. itraconazole, ritonavir, clarithromycin) is started or stopped during treatment with Sativex, a new dose titration may be required (see section 4.2).

Following treatment with the CYP3A4 inducer rifampicin reductions in C<sub>max</sub> and AUC of THC (40% and 20% reduction, respectively), its primary metabolite (85% and 87% reduction, respectively) and CBD (50% and 60% reduction, respectively) were observed. Therefore, concomitant treatment with strong enzyme inducers (e.g. rifampicin, carbamazepine, phenytoin, phenobarbital, St John's Wort) should be avoided whenever possible. If deemed necessary, careful titration is recommended, notably within the two weeks following the stop of the inducer.

Care should be taken with hypnotics, sedatives and drugs with potential sedating effects as there may be an additive effect on sedation and muscle relaxing effects.

Although there has been no greater rate of adverse events in patients already taking anti-spasticity agents with Sativex, care should be taken when co-administering Sativex with such agents since a reduction in muscle tone and power may occur, leading to a greater risk of falls.

Sativex may interact with alcohol, affecting co-ordination, concentration and ability to respond quickly. In general, alcoholic beverages should be avoided whilst using Sativex, especially at the beginning of treatment or when changing dose. Patients should be advised that if they do drink alcohol while using Sativex the additive CNS effects may impair their ability to drive or use machines, and increase the risk of falls.

#### **4.6 Fertility, pregnancy and lactation**

There is insufficient experience in humans regarding the effects of Sativex on reproduction. Therefore men and women of child bearing potential should take reliable contraceptive precautions for the duration of therapy and for three months after discontinuation of therapy.

##### **Pregnancy**

Sativex should not be used during pregnancy unless the potential risks to the fetus and/or embryo are considered to be outweighed by the benefit of treatment.

## Lactation

Available pharmacodynamics / toxicological data in animals have shown excretion of Sativex / metabolites in milk (for details see section 5.3).

A risk to the breastfed child cannot be excluded. Sativex is contraindicated during breast-feeding (see section 4.3).

## Fertility

In fertility studies in rodents, there was no effect of treatment with Sativex in males or females. There was no effect on fertility of the offspring from mothers treated with Sativex.

### 4.7 Effects on ability to drive and use machines

Sativex may produce undesirable effects such as dizziness and somnolence which may impair judgement and performance of skilled tasks. Patients should not drive, operate machinery or engage in any hazardous activity if they are experiencing any significant CNS effects such as dizziness or somnolence. Patients should be aware that Sativex has been known to cause a few cases of loss of consciousness.

### 4.8 Undesirable effects

The Sativex clinical program has so far involved over 1500 patients with MS in placebo controlled trials and long-term open label studies in which some patients used up to 48 sprays per day. Patients in clinical trials for relief of pain in cancer are described separately.

The most commonly reported adverse reactions in the first four weeks of exposure were dizziness, which occurs mainly during the initial titration period, and fatigue. These reactions are usually mild to moderate and resolve within a few days even if treatment is continued (see section 4.2). When the recommended dose titration schedule was used, the incidence of dizziness and fatigue in the first four weeks was much reduced.

The frequency of adverse events with a plausible relationship to Sativex, from placebo controlled trials in patients with MS, according to System Organ Classes (SOC) are given below (some of these adverse events may be part of the underlying condition).

MedDRA SOC	Very Common ≥ 1/10	Common ≥ 1/100 to < 1/10	Uncommon ≥ 1/1000 to < 1/100
Infections and infestations			pharyngitis
Metabolism and nutrition disorders		anorexia (including appetite decreased), appetite increased	
Psychiatric disorders		depression, disorientation, dissociation, euphoric mood,	hallucination (unspecified, auditory, visual), illusion, paranoia, suicidal ideation, delusional perception*
Nervous system disorders	dizziness	amnesia, balance disorder, disturbance in attention, dysarthria, dysgeusia, lethargy, memory impairment somnolence	syncope
Eye disorders		vision blurred	
Ear and labyrinth disorders		vertigo	

Cardiac disorders			palpitations, tachycardia
Vascular disorders			hypertension
Respiratory, thoracic and mediastinal disorders			throat irritation
Gastrointestinal disorders		constipation, diarrhoea, dry mouth, glossodynia, mouth ulceration, nausea, oral discomfort, oral pain, vomiting	abdominal pain (upper), oral mucosal discolouration*, oral mucosal disorder, oral mucosal exfoliation*, stomatitis, tooth discolouration
General disorders and administration site conditions	fatigue	application site pain, asthenia, feeling abnormal, feeling drunk, malaise	application site irritation
Injury, poisoning and procedural complaints		fall	

\* reported in long-term open-label studies:

A single case of ventricular bigeminy has been reported though this was in the context of acute nut allergy.

See also sections 4.4, 4.5 and 4.7.

#### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions.

Any suspected adverse events should be reported to the Ministry of Health according to the National Regulation by using an online form:

(<http://forms.gov.il/globaldata/getsequence/getsequence.aspx?formType=AdversEffectMedic@moh.health.gov.il>) or by email (adr@MOH.HEALTH.GOV.IL).

#### **Adverse events in patients with pain in cancer**

Treatment-emergent adverse events that occurred in 3% or more of patients given Sativex or placebo in a trial for patients with pain in cancer are given below in Table 3.

**Table 3: Treatment Emergent Adverse Events for Sativex in a placebo-controlled study in patients with pain in cancer**

	<b>Sativex n = 60 (%)</b>	<b>Placebo n = 59 (%)</b>
<b>Blood and Lymphatic System Disorders</b>		
Anaemia Nos	0	5
<b>Cardiac Disorders</b>		
Cardio-Respiratory Arrest	0	3
<b>Ear and Labyrinth Disorders</b>		
Vertigo	5	2
<b>Gastrointestinal Disorders</b>		
Nausea	12	10
Vomiting	8	7
Constipation	5	10
Oral Pain	2	5

	<b>Sativex n = 60 (%)</b>	<b>Placebo n = 59 (%)</b>
Diarrhoea	7	3
Glossodynia	3	0
Abdominal pain upper	2	3
Dry mouth	0	3
Stomatitis	2	3
<b>General Disorders and Administration Site Conditions</b>		
Pain exacerbated	0	3
Pyrexia	0	3
Weakness	5	0
Disease Progression	3	0
<b>Hepatobiliary Disorders</b>		
Hepatic cytolysis	0	3
<b>Infections and Infestations</b>		
Oral Candidiasis	3	2
Urinary Tract Infection	0	7
Lower Respiratory Tract Infection	0	5
<b>Investigations</b>		
GGT Increased	3	5
Blood Urea Increased	2	5
Liver Function Tests Abnormal	5	3
Blood Creatinine Increased	2	3
Blood Calcium Increased	0	5
<b>Musculoskeletal and Connective Tissue Disorders</b>		
Pain in Limb	2	3
Buttock Pain	0	3
<b>Neoplasms Benign, Malignant and Unspecified (incl.Cysts and Polyps)</b>		
Neoplasm Progression	10	5
Malignant Neoplasm Progression	2	5
<b>Nervous System Disorders</b>		
Somnolence	15	14
Dizziness	12	5
Disturbance in Attention	3	0
Dysgeusia	3	0
Headache	3	0
<b>Psychiatric Disorders</b>		
Confusion	7	3
Hallucination	3	2
Insomnia	3	2
Panic Attack	3	0
Euphoric Mood	3	0



	<b>Sativex n = 60 (%)</b>	<b>Placebo n = 59 (%)</b>
<b>Renal and Urinary Disorders</b>		
Urinary Retention	5	0
Haematuria	3	0
<b>Respiratory, Thoracic and Mediastinal Disorders</b>		
Dyspnoea	2	3
<b>Vascular Disorders</b>		
Hypotension	5	0

### **Urinary Retention and Infections**

The combined incidence of urinary retention and urinary infections appear to be increased in the cancer patients taking Sativex over those on placebo. Caution is advised in the urinary care of the cancer patients who are using Sativex.

### **Abnormal Haematologic and Clinical Chemistry Findings**

No consistent effect of Sativex on haematologic and clinical chemistry parameters has been observed.

### **Post-Market Adverse Drug Reactions**

Adverse event profile, based on post-market spontaneous reports, is consistent with those observed in clinical trials.

#### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product.

Any suspected adverse events should be reported to the Ministry of Health according to the National Regulation by using an online form

(<http://forms.gov.il/globaldata/getsequence/getsequence.aspx?formType=AdversEffectMedic@moh.health.gov.il>) or by email ([adr@MOH.HEALTH.GOV.IL](mailto:adr@MOH.HEALTH.GOV.IL)).

## **4.9 Overdose**

There is no experience of deliberate overdose with Sativex in patients. However, in a Thorough QT study of Sativex in 257 subjects, with 18 sprays taken over a 20-minute period twice daily, signs and symptoms of overdose/poisoning were observed. These consisted of acute intoxication type reactions including dizziness, hallucinations, delusions, paranoia, tachycardia or bradycardia with hypotension. In three of 41 subjects dosed at 18 sprays twice a day, this presented as a transient toxic psychosis which resolved upon cessation of treatment. Twenty-two subjects who received this substantial multiple of the recommended dose successfully completed the 5-day study period.

In the case of overdose, treatment should be symptomatic and supportive.

## **5. PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic Group: Other Analgesics and Antipyretics  
ATC Code: N02BG10

The European Medicines Agency has deferred the obligation to submit the results of studies with Sativex in one or more subsets of the paediatric population in spasticity. See section 4.2 for information on paediatric use.

### **Mechanism of action**

As part of the human endocannabinoid system (ECS), cannabinoid receptors, CB<sub>1</sub> and CB<sub>2</sub> receptors are found predominantly at nerve terminals where they have a role in retrograde regulation of synaptic function. THC acts as a partial agonist at both CB<sub>1</sub> and CB<sub>2</sub> receptors, mimicking the effects of the endocannabinoids, which may modulate the effects of neurotransmitters (e.g. reduce effects of excitatory neurotransmitters such as glutamate).

In animal models of MS and spasticity CB receptor agonists have been shown to ameliorate limb stiffness and improve motor function. These effects are prevented by CB antagonists, and CB<sub>1</sub> knockout mice show more severe spasticity. In the CREAE (chronic relapsing experimental autoimmune encephalomyelitis) mouse model, Sativex produced a dose-related reduction in the hind limb stiffness.

### **Clinical experience**

#### Treatment for the relief of spasticity in MS

Sativex has been studied at doses of up to 48 sprays/day in controlled clinical trials of up to 19 weeks duration in more than 1500 patients with MS. In the pivotal trials to assess the efficacy and safety of Sativex for symptom improvement in patients with moderate to severe spasticity due to multiple sclerosis (MS) the primary efficacy measure was a 0 to 10 point Numeric Rating Scale (NRS) on which patients indicated the average level of their spasticity related symptoms over the last 24 hours where 0 is no spasticity and 10 is the worst possible spasticity.

In a first Phase 3 placebo controlled trial over a 6-week treatment period the difference from placebo reached statistical significance but the difference between treatments of 0.5 to 0.6 points on the 0-10 point NRS was of questionable clinical relevance. In a responder analysis 40% Sativex and 22% placebo responded to treatment using the criterion of greater than a 30% reduction in NRS score. A second 14 week Phase 3 study failed to show a significant treatment effect. The difference from placebo on the NRS score was 0.2 points.

It was postulated that a clinically useful treatment effect in some patients might be partly masked by data from non-responders in the analyses of mean changes. In analyses comparing NRS scores with patient global impression of change (PGI), a 19% NRS response was estimated to represent a clinically relevant improvement on the PGI and a response of 28% “much improved” on the PGI. In post hoc exploratory combined analyses of the above two studies, a 4-week trial period using a 20% NRS response threshold was predictive of eventual response defined as a 30% reduction.

A third Phase 3 trial incorporated a formalised 4-week therapeutic trial period prior to randomisation. The aim of the trial was to assess the benefit of continued treatment for patients who achieve an initial response to treatment. 572 patients with MS and refractory spasticity all received single blind Sativex for four weeks. After four weeks on active treatment 273 achieved a reduction of at least 20% on the spasticity symptom NRS, of which 241 met the entry criteria for randomisation, with a mean change from the start of treatment of -3.0 points on the 10 point NRS. These patients were then randomised to either continue to receive active or switch to placebo for the 12 week double-blind phase, for a total of 16 weeks treatment overall.

During the double-blind phase the mean NRS scores for patients receiving Sativex generally remained stable (mean change from randomisation in NRS score -0.19), while the mean NRS scores for patients switched to placebo increased (mean change in NRS score was +0.64 and median change was +0.29). The difference\* between treatment groups was 0.84 (95% CI -1.29, -0.40).

\* Difference adjusted for centre, baseline NRS and ambulatory status

Of those patients who had a 20% reduction from screening in NRS score at week 4 and continued in the trial to receive randomised treatment, 74% (Sativex) and 51% (placebo) achieved a 30% reduction at week 16.

The results over the 12-week randomised phase are shown below for the secondary endpoints. The majority of secondary endpoints showed a similar pattern to the NRS score, with patients who continued to receive Sativex maintaining the improvement seen from the initial 4-week treatment period, while patients switching to placebo declined:

Modified Ashworth Score for spasticity:	Sativex -0.1; Placebo +1.8; Adjusted Difference -1.75 (95% CI -3.80, 0.30)
Spasm frequency (per day):	Sativex -0.05; Placebo +2.41 Adjusted Difference -2.53 (95% CI -4.27, -0.79)
Sleep disruption by spasticity: (0 to 10 NRS)	Sativex -0.25; Placebo +0.59; Adjusted Difference -0.88 (95% CI -1.25, -0.51)
Timed 10 metre walk (seconds):	Sativex -2.3; Placebo +2.0; Adjusted Difference -3.34 (95% CI -6.96, 0.26)
Motricity index (arm and leg):	No differences between treatment groups were seen.
Barthel Activities of Daily Living:	Odds ratio for improvement: 2.04

Subject global impression of change (OR=1.71), carer global impression of change (OR=2.40) and physician global impression of change (OR=1.96) all showed statistically significant superiority of Sativex over placebo.

The benefit of continued treatment in the long-term was studied in a placebo controlled, parallel group, randomised withdrawal trial in subjects taking long-term Sativex. Thirty six patients with a mean duration of Sativex use prior to the trial of 3.6 years were randomised to either continue with Sativex treatment or switch to placebo for 28 days. The primary endpoint was time to treatment failure, defined as the time from the first day of randomised treatment to a 20% increase in NRS or premature withdrawal from randomised treatment. Treatment failure was experienced by 44% of Sativex patients, and 94% of placebo patients, hazard ratio 0.335 (95% CI 0.16, 0.69).

#### Treatment for the symptomatic relief of neuropathic pain in multiple sclerosis

The potential efficacy of Sativex as an adjunct treatment for the symptomatic relief of neuropathic pain in multiple sclerosis was demonstrated by the results of a randomized, double-blind, placebo-controlled, parallel group, 4-week clinical study in multiple sclerosis patients with neuropathic pain (Study GWMS0107). There were 66 patients (14 male, 52 female) ranging in age from 27 to 71 (mean  $49 \pm 8.3$  standard deviation). The primary efficacy measure was the change from baseline of the mean BS-11, 11-box Numerical Rating Scale (NRS). To enter the study, the patient was required to have a pain severity score  $\geq 4$  on the 11-box NRS on at least four occasions during the 7-10 day baseline period. Regular medication for neuropathic pain had to have been stable for at least two weeks prior to entry and was maintained during the study. Sativex was self-titrated to symptom resolution or maximum tolerated dose.

Secondary efficacy measures included the Neuropathic Pain Scale (NPS) and sleep disturbance (also on an 11 point NRS). Completing patients from this study had the opportunity to enter an open-label extension study.

The baseline pain severity was 6.5 in the Sativex group and 6.4 in the placebo group. Analysis of the change from baseline of the mean 11-box NRS pain score showed a statistically significant treatment difference of -1.25 in favour of Sativex ( $p=0.005$ ; 95% CI: -2.11, -0.39 units).

Efficacy was also observed in the following secondary outcome measures. A pain reduction on the 11-box NRS of at least 50% was seen in 48% of the patients treated with Sativex, compared with 12% of the placebo group. Analysis of the change from baseline of the mean NPS showed a statistically significant treatment difference of -6.82 in favour of Sativex ( $p=0.039$ ; 95% CI: -13.28, -0.37). The NRS score for sleep improved by 2.73 from a baseline in the Sativex group, and by 1.41 in the placebo group. The treatment difference of -1.39 was significantly in favour of Sativex ( $p=0.003$ ; 95% CI: -2.27, -0.50).

The study medication was well tolerated. There were no serious adverse events during the study, and only one patient on Sativex discontinued due to an adverse event. Sixty-three of 66 (95%) eligible patients completing study GWMS0107 entered the long-term extension study.

#### Adjunctive treatment for the relief of pain in adult patients with advanced cancer who experience inadequate analgesia during the highest tolerated dose of strong opioid therapy for persistent background pain

The efficacy of Sativex was investigated in a two-week placebo-controlled, three-arm study in patients who reported moderate to severe pain despite already taking a strong opioid for persistent pain. The three groups were Sativex ( $n=53$ ), placebo ( $n=52$ ), and THC cannabis extract ( $n=56$ ). Some of the patients took high dose tramadol.

The primary comparison was between the Sativex group and the placebo group. There were two co-primary outcome endpoints: reduction from baseline at the last observation on the 11-point Numerical Rating Scale (NRS) and the use of escape medication at the last observation. The analysis was based on the intent-to-treat (ITT) population.

Baseline NRS scores were 5.68 in the Sativex group and 6.05 in the placebo group. The study outcomes showed a reduction from baseline on the NRS of 1.37 points on Sativex compared with 0.69 points on placebo. The difference between the two primary comparison groups was significant in favour of Sativex ( $p=0.024$ ). An improvement of greater than 30% in pain score was reported by 43% of patients on Sativex compared with 21% on placebo (Odds Ratio = 2.81; 95% CI: 1.22, 6.50). There was no difference in the use of escape medication between the two groups.

#### Clinical experience - abuse potential

In a study designed to identify its abuse potential, Sativex at a dose of 4 sprays taken at one time did not differ significantly from placebo. Higher doses of Sativex of 8 to 16 sprays taken at one time did show abuse potential comparable to equivalent doses of dronabinol, a synthetic THC. In a QTc study a dose of Sativex 4 sprays over 20 minutes twice daily was well-tolerated, but a substantially supratherapeutic dose of 18 sprays over 20 minutes twice daily resulted in significant psychoactivity and cognitive impairment.

## **5.2 Pharmacokinetic properties**

### **Absorption**

Following administration of Sativex (four sprays), both THC and CBD are absorbed fairly rapidly and appear in the plasma within 15 minutes after single oromucosal administration. With Sativex, a mean  $C_{max}$  of about 4 ng/mL was reached some 45-120 minutes after a single dose administration of a 10.8 mg THC dose, and was generally well tolerated with little evidence of significant psychoactivity.

When Sativex is co-administered with food the mean  $C_{max}$  and AUC for THC were 1.6- and 2.8-fold higher compared with fasting conditions. Corresponding parameters for CBD increased 3.3- and 5.1-fold.

There is a high degree of variability in pharmacokinetic parameters between patients. Following a single dose administration of Sativex (four sprays) under fasted conditions, the mean plasma level of THC showed a 57.3% CV for  $C_{max}$  (range 0.97-9.34ng/mL) and a 58.5% CV for AUC (range 4.2-

30.84 h\*ng/mL). Similarly the %CV for CBD was 64.1% (range 0.24-2.57ng/mL) and 72.5% (range 2.18-14.85 ng/mL) for the same parameters respectively. After nine consecutive days of dosing the % CV values for the same parameters were 54.2% ( $C_{\max}$  range = 0.92-6.37) and 37.4% ( $AUC_{0-t}$  = 5.34-15.01 h\*ng/mL) for THC and 75.7% ( $C_{\max}$  range 0.34-3.39 ng/mL) and 46.6% ( $AUC_{0-t}$  = 2.40-13.19 h\*ng/mL) for CBD respectively.

There is a high degree of variability in pharmacokinetic parameters within patients following single and repeat dosing. Of 12 subjects who received four sprays of Sativex as a single dose, eight had reductions in  $C_{\max}$  after nine days of multiple dosing, whilst three had increases (1 drop-out). For CBD, seven had reductions in  $C_{\max}$  after multiple dosing, whilst four had increases.

When Sativex is administered oromucosally, plasma levels of THC and other cannabinoids are lower compared with the levels achieved following inhalation of cannabinoids at a similar dose. A dose of 8 mg of vaporised THC extract, administered by inhalation resulted in mean plasma  $C_{\max}$  of more than 100 ng/mL within minutes of administration, with significant psychoactivity.

**Table to show PK parameters for Sativex, for vaporised THC extract and smoked cannabis**

	$C_{\max}$ THC ng/mL	$T_{\max}$ THC minutes	$AUC_{(0-t)}$ THC ng/mL/min
Sativex (providing 21.6 mg THC)	5.40	60	1362
Inhaled vaporised THC extract (providing 8 mg THC)	118.6	17.0	5987.9
Smoked cannabis* (providing 33.8 mg THC)	162.2	9.0	No data

\*Huestis et al, Journal of Analytical Toxicology 1992; 16:276-82.

### Distribution

As cannabinoids are highly lipophilic, they are quickly absorbed and distributed into body fat. The resultant concentrations in the blood following oromucosal administration of Sativex are lower than those obtained by inhaling the same dose of THC because absorption is slower and redistribution into fatty tissues is rapid. Additionally some of the THC undergoes hepatic first pass metabolism to 11-OH-THC, the primary metabolite of THC, and CBD similarly to 7-OH-CBD. Protein binding of THC is high (~97%). THC and CBD may be stored for as long as four weeks in the fatty tissues from which they are slowly released at sub-therapeutic levels back into the blood stream, then metabolised and excreted via the urine and faeces.

### Metabolism

THC and CBD are metabolised in the liver. Additionally some of the THC undergoes hepatic first pass metabolism to 11-OH-THC, the primary metabolite of THC, and CBD similarly to 7-OH-CBD. Human hepatic  $P_{450}$  2C9 isozyme catalyses the formation of 11-OH-THC, the primary metabolite, which is further metabolised by the liver to other compounds including 11-nor-carboxy- $\Delta^9$ -THC (THC-COOH), the most abundant metabolite in human plasma and urine. The  $P_{450}$ -3A subfamily catalyses the formation of other hydroxylated minor metabolites. CBD is extensively metabolised and more than 33 metabolites have been identified in urine. The major metabolic route is hydroxylation and oxidation at C-7 followed by further hydroxylation in the pentyl and propenyl groups. The major oxidized metabolite identified is CBD-7-oic acid containing a hydroxyethyl side chain.

See section 4.5 for information on drug interaction and metabolism by the cytochrome  $P_{450}$  enzyme system.

## **Elimination**

From clinical studies with Sativex, a non-compartmental PK analysis shows that the first order terminal elimination half life from plasma is 1.94, 3.72 and 5.25 hours for THC and 5.28, 6.39 and 9.36 for CBD following the administration of 2, 4 and 8 sprays respectively.

From the literature, elimination of oral cannabinoids from plasma is bi-phasic with an initial half-life of approximately four hours, and the terminal elimination half-lives are of the order of 24 to 36 hours or longer. Cannabinoids are distributed throughout the body; they are highly lipid soluble and accumulate in fatty tissue. The release of cannabinoids from fatty tissue is responsible for the prolonged terminal elimination half-life.

## **5.3 Preclinical safety data**

Effects in non-clinical studies were observed only at exposures considered sufficiently in excess of the maximum human exposure indicating little relevance to clinical use.

Reprotoxicity studies carried out with the THC and CBD extracts present in Sativex showed no adverse effects on either male or female fertility in terms of numbers of animals mating; number of fertile males and females, or on copulation or fertility indices. There were reduced absolute weights of epididymides, with a "no-effect" dosage level of 25 mg/kg/day (150 mg/m<sup>2</sup>) for male fertility. The "no-effect" dosage levels for effects on early embryonic and fetal survival, in rat studies, were approximately 1 mg/kg/day (6 mg/m<sup>2</sup>), which is close to or less than the likely maximum human dosage level of Sativex. There was no evidence to suggest any teratogenic activity in either rats or rabbits at dosage levels considerably in excess of likely human maximum dosage levels. However, in a rat pre- and post-natal study, pup survival and nursing behaviour were impaired at doses of 2 and 4 mg/kg/day (12 and 24 mg/m<sup>2</sup> respectively). Data from the literature have shown negative effects of THC and/or CBD on sperm number and motility.

In studies in animals, as expected, due to the lipophilic nature of cannabinoids, considerable levels of cannabinoids were found in the maternal breast milk. Following repeat dosing, cannabinoids are concentrated in breast milk (40 to 60 times the plasma level). Doses in excess of normal clinical doses may affect growth rates of breast-fed infants.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Ethanol anhydrous.

Propylene glycol.

Peppermint oil.

### **6.2 Incompatibilities**

In the absence of compatibility studies, this medicinal product must not be mixed with other medicinal products.

### **6.3 Shelf life**

2 years

In use stability after first opening: 42 days from date of opening.

### **6.4 Special precautions for storage**

Store in a refrigerator (2 to 8°C).

Once the spray container is opened and in use, refrigerated storage is not necessary but do not store above 25°C.  
Store upright.

#### **6.5 Nature and contents of container**

A Type I amber glass spray container (10 ml container has amber plastic-coated glass) fitted with a metering pump possessing a polypropylene dip tube and elastomer neck covered with a polyethylene cap. The metering pump delivers 100 microlitres per spray.

Pack Size: 10 mL.

10 mL pack size allows delivery after priming of up to 90 actuations (sprays) of 100 microlitres.

1, 2, 3, 4, 5, 6, 10 or 12 glass sprays containers per carton.

Not all pack sizes may be marketed.

#### **6.6 Special precautions for disposal**

Any unused product or waste material should be disposed of in accordance with local requirements.

### **7. MANUFACTURER**

GW Pharma Limited,  
Wiltshire,  
UK.

### **8. MARKETING AUTHORISATION HOLDER**

Neopharm Scientific Ltd.  
8 Hashiloach Street  
POB 7063  
Petach Tikva 49170  
Israel



### **9. REGISTRATION NO.**

147-48-33290-00

The format of this leaflet has been determined by the Ministry of Health and its content has been examined and approved- March 2015