

## 1. NAME OF THE MEDICINAL PRODUCT

Actrapid® Penfill® 100 iu/ml solution for injection in cartridge.

## 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

1 ml solution contains 100 iu insulin human\* (equivalent to 3.5 mg). 1 cartridge contains 3 ml equivalent to 300 iu.

\*Human insulin is produced in *Saccharomyces cerevisiae* by recombinant DNA technology.

Excipient with known effect:

100 iu of Actrapid® contains approximately 5 mcmol sodium, i.e. Actrapid® contains less than 1 mmol sodium (23 mg) per dose and is therefore considered essentially 'sodium-free'.

For the full list of excipients, see section 6.1.

## 3. PHARMACEUTICAL FORM

Solution for injection in cartridge. Penfill®.

The solution is clear, colourless and aqueous.

## 4. CLINICAL PARTICULARS

### 4.1 Therapeutic indications

Treatment of diabetes mellitus.

### 4.2 Posology and method of administration

#### Posology

The potency of human insulin is expressed in international units (iu).

Actrapid® dosing is individual and determined in accordance with the needs of the patient. It can be used alone or in combination with intermediate-acting or long-acting insulin before a meal or a snack.

The individual insulin requirement is usually between 0.3 and 1.0 iu/kg/day. The daily insulin requirement may be higher in patients with insulin resistance (e.g. during puberty or due to obesity) and lower in patients with residual, endogenous insulin production.

In patients with diabetes mellitus optimised glycaemic control delays the onset of late diabetic complications. Close blood glucose monitoring is therefore recommended. Adjustment of dose may be necessary if patients undertake increased physical activity, change their usual diet or during concomitant illness.

## Special populations

### Elderly ( $\geq 65$ years old)

Actrapid® can be used in elderly patients.

As with all insulin medicinal products, in elderly patients, glucose monitoring should be intensified and the insulin dose adjusted on an individual basis.

### Renal and hepatic impairment

Renal or hepatic impairment may reduce the patient's insulin requirements.

As with all insulin medicinal products, in patients with renal or hepatic impairment, glucose monitoring should be intensified and the human insulin dose adjusted on an individual basis.

### Paediatric population

Actrapid® can be used in children and adolescents.

### Transfer from other insulin medicinal products

When transferring from other insulin medicinal products, adjustment of the Actrapid® dose and the dose of the basal insulin may be necessary.

Close glucose monitoring is recommended during the transfer and in the initial weeks thereafter (see section 4.4).

## **Method of administration**

Actrapid® is a fast-acting human insulin and may be used in combination with intermediate or long-acting insulin medicinal products.

Actrapid® is administered subcutaneously in the abdominal wall, the thigh, the gluteal region or the deltoid region. Injection into a lifted skin fold minimises the risk of unintended intramuscular injection. The needle should be kept under the skin for at least 6 seconds to make sure the entire dose is injected. Injection sites should always be rotated within the same region in order to reduce the risk of lipodystrophy. As with all insulin medicinal products, subcutaneous injection in the abdominal wall ensures a faster absorption than other injection sites. As with all insulin medicinal products, the duration of action will vary according to the dose, injection site, blood flow, temperature and level of physical activity.

An injection should be followed within 30 minutes by a meal or snack containing carbohydrates.

Due to the risk of precipitation in pump catheters, Actrapid® should not be used in insulin pumps for continuous subcutaneous insulin infusion.

### Administration with an insulin delivery system

Actrapid® Penfill® is designed to be used with Novo Nordisk insulin delivery systems and NovoFine or NovoTwist needles.

Actrapid® Penfill® is accompanied by a package leaflet with detailed instructions for use to be followed.

### Intravenous use

If necessary, Actrapid® can be administered intravenously which should be carried out by healthcare professionals.

For intravenous use, infusion systems with Actrapid® at concentrations from 0.05 iu/ml to 1.0 iu/ml human insulin in the infusion fluids 0.9% sodium chloride, 5% dextrose and 10% dextrose inclusive 40 mmol/l potassium chloride using polypropylene infusion bags, are stable at room temperature for 24 hours. Although stable over time, a certain amount of insulin will initially be adsorbed to the material of the infusion bag. Monitoring of blood glucose is necessary during the insulin infusion.

### 4.3 Contraindications

Hypersensitivity to the active substance or to any of the excipients (see section 6.1).

### 4.4 Special warnings and precautions for use

Before travelling between different time zones, the patient should seek the doctor's advice since this may mean that the patient has to take the insulin and meals at different times.

#### Hyperglycaemia

Inadequate dosing or discontinuation of treatment, especially in type 1 diabetes, may lead to hyperglycaemia and diabetic ketoacidosis. Usually, the first symptoms of hyperglycaemia develop gradually over a period of hours or days. They include thirst, increased frequency of urination, nausea, vomiting, drowsiness, flushed dry skin, dry mouth, loss of appetite as well as acetone odour of breath. In type 1 diabetes, untreated hyperglycaemic events eventually lead to diabetic ketoacidosis, which is potentially lethal.

#### Hypoglycaemia

Omission of a meal or unplanned, strenuous physical exercise may lead to hypoglycaemia.

Hypoglycaemia may occur if the insulin dose is too high in relation to the insulin requirement. In case of hypoglycaemia or if hypoglycaemia is suspected Actrapid® must not be injected. After stabilisation of patient's blood glucose adjustment of the dose should be considered (see sections 4.8 and 4.9).

Patients, whose blood glucose control is greatly improved, e.g. by intensified insulin therapy, may experience a change in their usual warning symptoms of hypoglycaemia and should be advised accordingly. Usual warning symptoms may disappear in patients with longstanding diabetes.

Concomitant illness, especially infections and feverish conditions, usually increases the patient's insulin requirement. Concomitant diseases in the kidney, liver or affecting the adrenal, pituitary or thyroid gland can require changes in the insulin dose.

When patients are transferred between different types of insulin medicinal products, the early warning symptoms of hypoglycaemia may change or become less pronounced than those experienced with their previous insulin.

#### Transfer from other insulin medicinal products

Transferring a patient to another type or brand of insulin should be done under strict medical supervision. Changes in strength, brand (manufacturer), type, origin (animal, human insulin or human insulin analogue) and/or method of manufacture (recombinant DNA versus animal source insulin) may result in a need for a change in dose. Patients transferred to Actrapid® from another type of insulin may require an increased number of daily injections or a change in dose from that used with their usual insulin medicinal products. If an adjustment is needed, it may occur with the first dose or during the first few weeks or months.

#### Injection site reactions

As with any insulin therapy, injection site reactions may occur and include pain, redness, hives, inflammation, bruising, swelling and itching. Continuous rotation of the injection site within a given area may help to reduce the risk of developing these reactions. Reactions usually resolve in a few days to a few weeks. On rare occasions, injection site reactions may require discontinuation of Actrapid®.

## Combination of Actrapid® with pioglitazone

Cases of cardiac failure have been reported when pioglitazone was used in combination with insulin, especially in patients with risk factors for development of cardiac heart failure. This should be kept in mind if treatment with the combination of pioglitazone and Actrapid® is considered. If the combination is used, patients should be observed for signs and symptoms of heart failure, weight gain and oedema. Pioglitazone should be discontinued if any deterioration in cardiac symptoms occurs.

### **4.5 Interaction with other medicinal products and other forms of interaction**

A number of medicinal products are known to interact with glucose metabolism.

#### **The following substances may reduce the patient's insulin requirement:**

Oral antidiabetic medicinal products, monoamine oxidase inhibitors (MAOI), beta-blockers, angiotensin converting enzyme (ACE) inhibitors, salicylates, anabolic steroids and sulphonamides.

#### **The following substances may increase the patient's insulin requirement:**

Oral contraceptives, thiazides, glucocorticoids, thyroid hormones, sympathomimetics, growth hormone and danazol.

Beta-blockers may mask the symptoms of hypoglycaemia.

Octreotide/lanreotide may either increase or decrease the insulin requirement.

Alcohol may intensify or reduce the hypoglycaemic effect of insulin.

### **4.6 Fertility, pregnancy and lactation**

#### Fertility

Animal reproduction studies with human insulin have not revealed any adverse effects on fertility.

#### Pregnancy

There are no restrictions on treatment of diabetes with insulin during pregnancy, as insulin does not pass the placental barrier.

Both hypoglycaemia and hyperglycaemia, which can occur in inadequately controlled diabetes therapy, increase the risk of malformations and death *in utero*. Intensified blood glucose control and monitoring of pregnant women with diabetes are recommended throughout pregnancy and when contemplating pregnancy. Insulin requirements usually fall in the first trimester and increase subsequently during the second and third trimesters. After delivery, insulin requirements normally return rapidly to pre-pregnancy values.

#### Breast-feeding

There is no restriction on treatment with Actrapid® during breast-feeding. Insulin treatment of the nursing mother presents no risk to the baby. However, the Actrapid® dose may need to be adjusted.

### **4.7 Effects on ability to drive and use machines**

The patient's ability to concentrate and react may be impaired as a result of hypoglycaemia. This may constitute a risk in situations where these abilities are of special importance (e.g. driving a car or operating machinery).

Patients should be advised to take precautions to avoid hypoglycaemia while driving. This is particularly important in those who have reduced or absent awareness of the warning signs of hypoglycaemia or have frequent episodes of hypoglycaemia. The advisability of driving should be considered in these circumstances.

## 4.8 Undesirable effects

### a. Summary of the safety profile

The most frequently reported adverse reaction during treatment is hypoglycaemia. The frequencies of hypoglycaemia vary with patient population, dose regimens and level of glycaemic control, please see section c below.

At the beginning of the insulin treatment, refraction anomalies, oedema and injection site reactions (pain, redness, hives, inflammation, bruising, swelling and itching at the injection site) may occur. These reactions are usually of transitory nature. Fast improvement in blood glucose control may be associated with acute painful neuropathy, which is usually reversible. Intensification of insulin therapy with abrupt improvement in glycaemic control may be associated with temporary worsening of diabetic retinopathy, while long-term improved glycaemic control decreases the risk of progression of diabetic retinopathy.

### b. Tabulated list of adverse reactions

Adverse reactions listed below are based on clinical trial data and classified according to MedDRA frequency and System Organ Class. Frequency categories are defined according to the following convention: Very common ( $\geq 1/10$ ); common ( $\geq 1/100$  to  $< 1/10$ ); uncommon ( $\geq 1/1,000$  to  $< 1/100$ ); rare ( $\geq 1/10,000$  to  $< 1/1,000$ ); very rare ( $< 1/10,000$ ); not known (cannot be estimated from the available data).

Immune system disorders	Uncommon – Urticaria, rash
	Very rare – Anaphylactic reactions*
Metabolism and nutrition disorders	Very common – Hypoglycaemia*
Nervous system disorders	Uncommon – Peripheral neuropathy (painful neuropathy)
Eye disorders	Uncommon – Refraction disorders
	Very rare – Diabetic retinopathy
Skin and subcutaneous tissue disorders	Uncommon – Lipodystrophy*
General disorders and administration site conditions	Uncommon – Injection site reactions, Oedema

\* see section c

### c. Description of selected adverse reactions

#### *Anaphylactic reactions:*

The occurrence of generalised hypersensitivity reactions (including generalised skin rash, itching, sweating, gastrointestinal upset, angioneurotic oedema, difficulty in breathing, palpitation and reduction in blood pressure) is very rare but can potentially be life threatening.

#### *Hypoglycaemia:*

The most frequently reported adverse reaction is hypoglycaemia. It may occur if the insulin dose is too high in relation to the insulin requirement. Severe hypoglycaemia may lead to unconsciousness and/or convulsions and may result in temporary or permanent impairment of brain function or even death.

The symptoms of hypoglycaemia usually occur suddenly. They may include cold sweats, cool pale skin, fatigue, nervousness or tremor, anxiousness, unusual tiredness or weakness, confusion, difficulty in concentration, drowsiness, excessive hunger, vision changes, headache, nausea and palpitation.

In clinical trials, the frequency of hypoglycaemia varied with patient population, dose regimens and level of glycaemic control.

#### *Lipodystrophy:*

Lipodystrophy (including lipohypertrophy, lipoatrophy) may occur at the injection site. Continuous rotation of the injection site within the particular injection area may help to reduce the risk of developing these reactions.

#### d. Paediatric population

Based on post-marketing sources and clinical trials, the frequency, type and severity of adverse reactions observed in the paediatric population do not indicate any differences to the broader experience in the general population.

#### e. Other special populations

Based on post-marketing sources and clinical trials, the frequency, type and severity of adverse reactions observed in elderly patients and in patients with renal or hepatic impairment do not indicate any differences to the broader experience in the general population.

### **4.9 Overdose**

A specific overdose of insulin cannot be defined, however, hypoglycaemia may develop over sequential stages if too high a dose relative to the patient's requirement is administered:

- Mild hypoglycaemic episodes can be treated by oral administration of glucose or sugary products. It is therefore recommended that the diabetic patient always carries sugar-containing products.
- Severe hypoglycaemic episodes, where the patient has become unconscious, can be treated with glucagon (0.5 to 1 mg) given intramuscularly or subcutaneously by a trained person, or with glucose given intravenously by a healthcare professional. Glucose must be given intravenously, if the patient does not respond to glucagon within 10 to 15 minutes. Upon regaining consciousness, administration of oral carbohydrates is recommended for the patient in order to prevent a relapse.

## **5. PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: Drugs used in diabetes. Insulins and analogues for injection, fast-acting, insulin (human). ATC code: A10AB01.

#### Mechanism of action and pharmacodynamic effects

The blood glucose lowering effect of insulin is due to the facilitated uptake of glucose following binding of insulin to receptors on muscle and fat cells and to the simultaneous inhibition of glucose output from the liver.

A clinical trial in a single intensive care unit treating hyperglycaemia (blood glucose above 10 mmol/l) in 204 diabetic and 1344 non-diabetic patients undergoing major surgery showed that normoglycaemia (blood glucose 4.4 – 6.1 mmol/l) induced by intravenous Actrapid® reduced mortality by 42% (8% versus 4.6%).

Actrapid® is a fast-acting insulin.

Onset of action is within ½ hour, reaches a maximum effect within 1.5–3.5 hours and the entire duration of action is approximately 7–8 hours.

## 5.2 Pharmacokinetic properties

Insulin in the blood stream has a half-life of a few minutes. Consequently, the time-action profile of an insulin preparation is determined solely by its absorption characteristics.

This process is influenced by several factors (e.g. insulin dose, injection route and site, thickness of subcutaneous fat, type of diabetes). The pharmacokinetics of insulin medicinal products are therefore affected by significant intra- and inter-individual variation.

### Absorption

The maximum plasma concentration is reached within 1.5–2.5 hours after subcutaneous administration.

### Distribution

No profound binding to plasma proteins, except circulating insulin antibodies (if present) has been observed.

### Metabolism

Human insulin is reported to be degraded by insulin protease or insulin-degrading enzymes and possibly protein disulfide isomerase. A number of cleavage (hydrolysis) sites on the human insulin molecule have been proposed; none of the metabolites formed following the cleavage are active.

### Elimination

The terminal half-life is determined by the rate of absorption from the subcutaneous tissue. The terminal half-life ( $t_{1/2}$ ) is therefore a measure of the absorption rather than of the elimination *per se* of insulin from plasma (insulin in the blood stream has a  $t_{1/2}$  of a few minutes). Trials have indicated a  $t_{1/2}$  of about 2-5 hours.

### Paediatric population

The pharmacokinetic profile of Actrapid® has been studied in a small number (n=18) of diabetic children (aged 6–12 years) and adolescents (aged 13–17 years). The data are limited but suggest that the pharmacokinetic profile in children and adolescents may be similar to that in adults. However, there were differences between age groups in  $C_{max}$ , stressing the importance of individual dose titration.

## 5.3 Preclinical safety data

Non-clinical data reveal no special hazard for humans based on conventional studies of safety pharmacology, repeated dose toxicity, genotoxicity, carcinogenic potential, toxicity to reproduction and development.

## 6. PHARMACEUTICAL PARTICULARS

### 6.1 List of excipients

Zinc chloride  
Glycerol  
Metacresol  
Sodium hydroxide (for pH adjustment)  
Hydrochloric acid (for pH adjustment)  
Water for injections

## **6.2 Incompatibilities**

Insulin medicinal products should only be added to compounds with which it is known to be compatible. Medicinal products added to the insulin solution may cause degradation of the insulin, e.g. if the medicinal products contain thiols or sulphites.

## **6.3 Shelf life**

Before opening: 30 months.

During use or when carried as a spare: The product must be stored for a maximum of 6 weeks. Store below 30°C.

## **6.4 Special precautions for storage**

Before opening: Store in a refrigerator (2°C – 8°C). Do not freeze.

During use or when carried as a spare: Store below 30°C. Do not refrigerate. Do not freeze.

Keep the cartridge in the outer carton in order to protect from light.

For storage conditions of the medicinal product, see section 6.3.

## **6.5 Nature and contents of container**

3 ml solution in cartridge (type 1 glass) with a plunger (bromobutyl) and a stopper (bromobutyl/polyisoprene).

Pack sizes of 5 cartridges.

Not all pack sizes may be marketed.

## **6.6 Special precautions for disposal and other handling**

Needles and Actrapid® Penfill® must not be shared. The cartridge must not be refilled.

Do not use this medicinal product if you notice that the solution is not clear, colourless and aqueous.

Actrapid® which has been frozen must not be used.

The patient should be advised to discard the needle after each injection.

In case of emergency in current Actrapid® users (hospitalisation or insulin pen malfunction), Actrapid® can be withdrawn with an U100 insulin syringe from a cartridge.

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

## **7. REGISTRATION NUMBER:**

Actrapid® Penfill®: 049-82-26016-00



**8. MANUFACTURER:**

Novo Nordisk A/S  
Novo Allé  
DK-2880 Bagsværd  
Denmark

**9. MARKETING AUTHORISATION HOLDER:**

Novo Nordisk Ltd.  
20 HaTa'as St. Indust. Area,  
Kfar Saba, 4442520

The content of this leaflet was checked and approved by the Ministry of Health in May 2014.