

Read this package insert carefully in its entirety before using this medicine

The format of this leaflet was determined by the Ministry
of Health and its content was checked and approved by it in 07/2011

SALIKAREN

Ointment

Composition: Salicylic Acid 3%

Inactive Ingredients: Paraffin liquid, Hard Paraffin, Propylene Glycol, Vaseline, Olive Oil, Lanolin, Hard Fat, Sodium EDTA.

Indication: This medicine is indicated as a topical aid in the removal of excessive keratin in hyperkeratotic skin disorders, including dandruff, keratosis palmaris and plantaris, psoriasis - keratin of scalp, hands, legs and elbows.

Therapeutic group: Keratolytic from salicylic group.

When Should The Preparation Not Be Used?

Do not use this medicine if you are sensitive to any of its ingredients.

Do not take this medicine without consulting a doctor before starting treatment:

If you are pregnant or breast-feeding.

If you are suffering or have suffered in the past, from impaired function of the circulatory system or diabetes mellitus.

In case of inflammation, irritation, or infection of the skin.

Warnings: Do not use this medicine frequently, or for a long period, without consulting a doctor.

Do not use on open wounds or cuts. Do not use on large surface of the skin without consulting a doctor.

If you are sensitive to any type of food or medicine, inform your doctor before commencing treatment with this medicine.

Avoid contact with eyes, nose, mouth or other mucous membranes. If contact occurs flush with a lot of water.

Do not use occlusive dressing unless directed by the physician.

Do not use on moles, birthmarks and hairy warts.

Drug Interactions: If you are taking another drug concomitantly or if you have just finished treatment with another medicine inform the attending doctor, in order to prevent hazards or lack of efficacy arising from drug interactions.

Do not simultaneously use (at same site, time) with other preparations containing peeling agents such as: Tretinoin, Retinoic acid, Resorcin, Benzoyl Peroxide, Sulfur and like, without consulting a doctor.

Simultaneous use of ointments containing salicylic acid can cause

irritation and redness in the treated area. Apply only one of the ointments.

Side effects: In addition to the desired effect of the medicine, adverse reactions may occur during the course of taking this medicine, for example: rash, irritation, burning, peeling, ulcers, local inflammation - discontinue treatment and contact your doctor.

In the event that you experience side effects not mentioned in this leaflet, or if there is a change in your general health, consult your doctor immediately.

Dosage: This medicine is not usually intended for administration to children and infants under 2 years of age. At the beginning of the treatment, use this medicine on a small skin surface or apply only once a day to ensure that skin irritation does not occur.

Attention: Do not swallow! This medicine is intended for external use only.

Directions for use: Before the beginning of treatment, wash your hands with soap. Soften the affected skin with warm water and dry thoroughly. Then with the help of your fingers apply enough ointment to cover the area. At the beginning of the treatment, use this medicine on a small skin surface or apply only once a day to ensure that skin irritation does not occur.

After few days of treatment, you may apply gently 2-3 times a day.

If the treatment is not for the hands, wash them thoroughly after every use. If within 6 weeks of treatment no improvement or deterioration exists, consult a doctor.

AVOID POISONING!

This medicine, and all other medicines, must be stored in a safe place out of the reach of children and/or infants, to avoid poisoning.

If you have taken an overdose, or if a child has accidentally swallowed the medicine, proceed immediately to a hospital emergency room and bring the package of the medicine with you. Do not induce vomiting unless explicitly instructed to do so by a doctor! Do not take medicines in the dark! Check the label and the dose each time you take your medicine. Wear glasses if you need them.

Storage: Store in a dark place below 25°C

Even if kept in their original container and stored as recommended, medicines may be kept for a limited period only. Please note the expiry date of the medicine! In case of doubt, consult the pharmacist who dispensed the medicine to you.

Do not store different medications in the same package.

Licence Number: 18382004700

Manufacturer: Rekah Pharm. Prod. Ltd., 30 Hamelacha St. Holon, Israel.