

**PATIENT PACKAGE INSERT IN ACCORDANCE WITH THE PHARMACISTS' REGULATIONS (PREPARATIONS) - 1986**

Read the package insert carefully in its entirety before using this medicine

This leaflet format was determined by the Ministry of Health and the content thereof was checked and approved by it in April 2011

# Strepsils® Sugar Free

## Lemon Lozenges

### COMPOSITION

Each lozenge contains:  
2,4-Dichlorobenzyl alcohol 1.2 mg  
Amylmetacresol 0.6 mg

### Inactive Ingredients:

Flav P Lemon Flavor 74940-74, Sodium Saccharin, Tartaric Acid, Maltitol liquid, Isomalt, Quinoline Yellow (E104 CI 47005), Water Purified.

### THERAPEUTIC GROUP

Antiseptic.

### INDICATION

Sugar free antiseptic for the relief of sore throat in children and adults.

### WHEN SHOULD THE PREPARATION NOT BE USED?

Do not use this medicine if you are sensitive to any of its ingredients.

### WARNINGS

Do not use this medicine frequently, or for a long period, without consulting a doctor.  
If you are sensitive to any type of food or medicine, inform the doctor before commencing treatment with this medicine.

### DRUG INTERACTIONS

If you are taking another drug concomitantly or if you have just finished treatment with another medicine, inform the doctor, in order to prevent hazards or lack of efficacy arising from drug interactions.

### SIDE EFFECTS

In addition to the desired effect of the medicine, adverse reactions may occur during the course of taking this medicine.

### Effects that require special attention

Irritation, local infection or rash (didn't develop before the use of this medicine) (rare): stop treatment and refer to your doctor.

In the event that you experience side effects not mentioned in this leaflet, or if there is a change in your general health, consult the doctor.

### ADVERSE REACTIONS AND DRUG INTERACTIONS IN CHILDREN

Parents must inform the attending doctor about any side effect, as well as any additional medicine being taken by a child!  
See above for details of special side effects.

### RECOMMENDED DOSAGE UNLESS OTHERWISE PRESCRIBED BY THE DOCTOR

Adults and children: one lozenge to be sucked every 2-3 hours.

Do not exceed the recommended dosage.

This medicine is not usually intended for administration to children and infants under 6 years of age.

If there is no improvement in your condition within a few days, consult a doctor.

Sore throat that is severe or persistent or accompanied by fever, headache or nausea and vomiting requires immediate consultation with a doctor.

### DIRECTIONS FOR USE

The lozenge should be dissolved in the mouth. Do not chew or break the lozenge.

Suck the lozenge slowly.

### AVOID POISONING!

This medicine, and all other medicines, must be stored in a safe place out of the reach of children and/or infants, in order to avoid poisoning.

If you have taken an overdose, or if a child has accidentally swallowed the medicine, proceed immediately to a hospital emergency room and bring the package of the medicine with you.

Do not induce vomiting unless explicitly instructed to do so by a doctor!

Do not take medicines in the dark! Check the label and the dose each time you take your medicine. Wear glasses if you need them.

### STORAGE

Store this medicine below 25°C.  
Even if kept in their original container and stored as recommended, medicines may be kept for a limited period only. Please note the expiry date of the medicine! In case of doubt, consult the pharmacist who dispensed the medicine to you. Do not store different medications in the same package.

**DRUG REGISTRATION NUMBER:** 052-52-24917-00

### MANUFACTURER:

Reckitt Benckiser Healthcare International Ltd., Nottingham, England.

### LICENSE HOLDER:

Reckitt Benckiser (Near East) Ltd.,  
6 Hanagar St., Hod Hasharon 45240.