PATIENT PACKAGE INSERT The medicine is not Before beginning Use in children and If a child or anyone twice. Shake the bottle associated with the use other medicine, should cocamidopropylamine treatment with Agispor, IN ACCORDANCE WITH intended for infants below infants else has accidentally well, apply a measured of Bifonazole: be kept in a closed oxide, Perfume 199-THE PHARMACISTS' tell the doctor if amount of shampoo 95. purified water the age of one month. Use in infants and swallowed the medicine place out of the sight Reactions on the skin REGULATIONS vou have a known proceed immediately to to the hair or affected and reach of children toddlers must be under and subcutaneous What does the (PREPARATIONS) - 1986 . WHAT IS THE sensitivity to antifungal the doctor or a hospital area of the skin and and/or infants in order to medical supervision. tissue: medicine look like and MEDICINE INTENDED leave the shampoo on medicines of the emergency room and avoid poisoning. Do not what are the contents The medicine is dispensed If there is no Contact dermatitis. FOR? imidazole group such as: bring the package of the the scalp for 5 minutes induce vomiting without allergic of the package: improvement in your skin with a doctor's explicit instructions econazole, clotrimazole, medicine with you. before washing off. Use For the treatment of mycotic condition within 6 inflammation, reddening, A plastic bottle prescription only. a sufficient quantity from the doctor. miconazole. diseases of the scalp, such weeks, refer to the itching, rash, urticaria, containing 100 If you forgot to take of shampoo to ensure vou are sensitive to Do not use the medicine milliliters of Agispor **AGISPOR** as pityriasis versicolor and doctor again. blistering, peeling of the the medicine do not good foaming of the seborrheic dermatitis. any type of food or skin, eczema, dry skin, after the expiry date Shampoo. Avoid contact of the take a double dose. shampoo. (exp. date) appearing medicine. skin irritation, softening preparation with the Name of manufacture SHAMPOO Take the next dose at Therapeutic group Do not take medicines on the package. The of the skin, and burning and registration eves. the scheduled time and An antifungal belonging to III you are taking, or in the dark! Check the expiry date refers to the sensation. holder: Perrigo Israel consult the doctor. the family of imidazoles. have recently taken label and the dose last day of that month. Each bottle contains: 3. HOW SHOULD Pain and/or edema in Pharmaceuticals Ltd. Adhere to the treatment other medicines, each time you take the Can be used for 6 the treated area. P.O.B. 16, Yeruham. Bifonazole 1% YOU USE THE regimen as recommended including nonmedicine. Wear glasses 2. BEFORE USING THE months after first MEDICINE? This leaflet was Inactive ingredients and These side effects by the doctor. prescription medicines MEDICINE if you need them. openina. allergens in the preparation disappear after cessation checked and approved Always use according Even if there is an and nutritional If you have further Store in a cool place by the Ministry of see Section 6. to the doctor's of treatment, however it improvement in your ☑Do not use the supplements, tell the questions regarding below 25°C. these effects persist and/ Health in August instructions. health, do not stop medicine: doctor or pharmacist. use of the medicine, Read this leaflet carefully or are bothersome, refer Store in the original Check with the doctor treatment with the · If you are sensitive Especially inform the in its entirety before using consult the doctor or to the doctor. packaging. Registration number or pharmacist if you are medicine without doctor or pharmacist if the medicine. This leaflet (allergic) to the active pharmacist. of the medicine in the uncertain consulting the doctor or If any of the side effects you are taking warfarin ingredient or to any of contains concise information 6. FURTHER National Drug Registry The dosage and manner pharmacist. worsen, or if you suffer the other ingredients 4. SIDE EFFECTS about the medicine. If you INFORMATION of the Ministry of from a side effect that APregnancy and of treatment will be Method of use in the have further questions. found in the medicine. As with any medicine. Health: breastfeeding determined by the absence of other is not mentioned in the In addition to the active refer to the doctor or use of Agispor Shampoo ingredient, the medicine 12623.26745 leaflet, consult with the Do not use the medicine doctor only. instructions from the pharmacist Special warnings may cause side effects in doctor. also contains: if vou are pregnant Do not exceed the regarding use of the some users. Do not be This medicine has been Sodium lauryl sulphate, or breastfeeding, recommended dose. Use for 4 weeks. The medicine prescribed to treat you. Do alarmed by reading the 5. HOW SHOULD sodium lauryl ether except with the explicit recommended dosage is For external use only. Do not use the medicine list of side effects. You not pass it on to others. I THE MEDICINE BE sulphate, ethyl alcohol, recommendation of the 3 times a week. At every frequently, or for a long Do not swallow. may not suffer from any may harm them, even i STORED? lactic acid, cocamide doctor. shampooing, the scalp period, without consulting it seems to you that their Shake before use. of them Avoid poisoning! This diethanolamine. should be shampooed the doctor. medical condition is similar Known side effects medicine, and any