

**PATIENT PACKAGE INSERT IN ACCORDANCE
WITH THE PHARMACISTS' REGULATIONS
(PREPARATIONS) - 1986**

Read this package insert carefully in its entirety
before using this medicine

The format of this leaflet was determined by
the Ministry of Health and its content was
checked and approved by it in May 2012

CORSODYL™ DENTAL GEL

Composition:

Active ingredient and concentration:

Chlorhexidine Digluconate 1% W/W

Inactive ingredients:

Purified Water, Isopropyl Alcohol 4%,
Hydroxypropylcellulose, Sodium Acetate,
Peppermint Oil, Levomenthol, Macrogolglycerol
Hydroxystearate

Therapeutic group: Antiseptic.

Therapeutic activity:

Disinfectant and antiseptic, for use in the oral
cavity.

The medicine is intended:

- For inhibition of formation of dental plaque.
- As an aid in the treatment and prevention of gingivitis.
- As an aid in the maintenance of oral hygiene.
- For use following dental surgery or treatments, to promote healing of the gums.
- For use in mouth ulcers, inflammations due to dentures and oral thrush.
- As an aid in the prevention of dental caries in high risk patients, as an adjunct to fluoride.

When should the preparation not be used?

Do not use if you know of sensitivity to chlorhexidine digluconate or to any of the medicine's ingredients.

Do not use this medicine without consulting a physician before starting treatment:

If you are pregnant or breastfeeding.

In children under 12 years of age.

Warnings:

Toothpastes may impair the action of Corsodyl Dental Gel. Therefore, after brushing - rinse the mouth thoroughly with water and wait 5 minutes after brushing the teeth with toothpaste and before using Corsodyl Dental Gel, or use Corsodyl Dental Gel at times of the day different from the times of toothbrushing.

When using Corsodyl Dental Gel to help in the treatment and prevention of gingivitis, the gums may bleed when brushing.

Do not bleach fabrics that have been in contact with the gel.

Drug interactions:

If you are taking another medicine, or if you have just finished treatment with another medicine, inform the attending physician, in order to prevent hazards or lack of efficacy arising from drug interactions.

Side effects:

In addition to the desired effect of the medicine, side effects may occur during the course of treatment with this medicine, such as: change in sense of taste or burning sensation.

These effects usually disappear within a short time following the period of adaptation to the preparation. If the effects persist, refer to the physician.

Side effects that require special attention:

Temporary staining of the tongue. This effect disappears after stop using the preparation.

Temporary staining of the teeth that can usually be removed by brushing. Staining can be avoided by not drinking tea, coffee or red wine - especially one hour after use, and by brushing daily with toothpaste. If staining remains it can be removed with a scale and polish from the dentist or hygienist. You can clean your dentures with denture cleaner.

In case of pain, swelling or irritation in the mouth: stop treatment and refer to the physician.

Severe allergic reaction that can be manifested by rash, swelling of the mouth or face or difficulty breathing (rare): stop treatment and refer to the physician immediately!

In any event that you experience side effects not mentioned in this leaflet, or if there is a change in your general health, consult your physician immediately.

Dosage:

Recommended dosage unless otherwise instructed by the physician:

For treatment of gingivitis:

Brush the teeth for at least one minute, once or twice daily with 2 cm of gel. Continue the treatment for about one month.

For treatment of mouth ulcers and oral thrush:

Apply the gel directly on the sore areas using the finger tip or a cotton bud once or twice daily for at least one minute. Continue the treatment for an additional two days after healing.

For treatment of mouth inflammations due to dentures:

Before inserting the dentures, apply the gel gently on the dentures in the surfaces that come in contact with the gum. Clean and soak the dentures in Corsodyl Mint Mouthwash for 15 minutes twice daily or as directed by the dentist.

As an aid in the prevention of dental caries or after dental surgery: Use according to the dentist's instructions.

Do not exceed the recommended dosage.

If there is no improvement in your condition, refer to your dentist.

Attention:

Do not swallow! This medicine is intended for oral use only.

Avoid contact with the eyes or ears. In case of contact with the eyes or ears, rinse them well with water.

Avoid poisoning!

This medicine, and all other medicines, must be stored in a safe place out of the reach of children and/or infants, to avoid poisoning.

If you have taken an overdose, or if a child has accidentally swallowed the medicine, proceed immediately to a hospital emergency room and bring the package of the medicine with you.

Do not induce vomiting unless explicitly instructed to do so by a physician!

Do not take medicines in the dark! Check the label and the dose each time you take your medicine. Wear glasses if you need them.

Storage:

Do not store above 25°C.

Even if kept in their original container and stored as recommended, medicines may be kept for a limited period only.

Please note the expiry date of the medicine! In case of doubt, consult the pharmacist who dispensed the medicine to you.

Do not store different medications in the same package.

License number: 114-62-29571

Manufacturer:

Purna Pharmaceuticals, Puurs, Belgium.

License Holder and Importer:

GlaxoSmithKline (Israel) Ltd., 25 Basel St., Petach Tikva.