

**PATIENT PACKAGE INSERT IN ACCORDANCE WITH THE  
PHARMACISTS' REGULATIONS (PREPARATIONS) - 1986**

Read this leaflet carefully in its entirety  
before using this medicine

The format of this leaflet was determined by  
the Ministry of Health and its content  
was checked and approved by it in November 2011

## BABYZIM

### Drops

- **Composition:** Each teaspoonful (5 ml) contains:
- Pepsin 1:10,000 30 mg
- Sodium Glycerophosphate 40 mg
- Nicotinic Acid 10 mg

**Inactive ingredients:**

Calcium Chloride, Hydrochloric Acid, Sodium Methylparaben (Nipagin), Glycine, Glycerin, Ethanol, Lactic Acid, Tutti Frutti, Purified Water.

Each 5 ml contains approx. 0.3 g alcohol

**Therapeutic group:** Digestive enzymes.

**Indication:** For the treatment of constipation and flatulence.

**When should the preparation not be used?**

Do not use this medicine if there is a known sensitivity to any of its ingredients.

**Do not take this medicine without consulting a physician before starting treatment:**

If you are suffering or have suffered in the past from impaired function of: the heart and/or blood vessels, the kidney/urinary system.

**Warnings:**

Do not use this medicine frequently or for a prolonged period without consulting your physician.

If you are sensitive to any kind of food or drugs, inform your physician before taking this medicine.

**Drug interactions:**

If you are taking another drug concomitantly, or if you have just finished treatment with another medicine, including non-prescription medicines and food supplements, inform the attending physician, in order to prevent hazards or lack of efficacy arising from drug interactions.

**Side effects:**

In addition to the desired effect of the medicine, during treatment unwanted side effects may also occur.

In the event that you experience side effects not mentioned in this leaflet, or if there is a change in your general health, consult your physician immediately.

**Adverse reactions and drug interactions in children and infants:**

Parents must inform the attending physician about any side effects, as well as any additional medicine being taken by the child!

**Dosage:**

Recommended dosage unless otherwise prescribed by your physician:

Infants up to one year: 5 drops before food and 5 drops after food at every feeding (up to 50 drops daily).

Infants 1-2 years: 20-40 drops after every meal.

Children 2-5 years: 40-60 drops after every meal.

Children 5-14 years: 0.5-1 teaspoonful of 5 ml after every meal.

Do not exceed the recommended dosage.

In the treatment of constipation, if there is no improvement within a week refer to your physician.

**Directions for use:**

Take this medicine with a meal.

Drip the medicine first into a teaspoon and not directly into the baby's mouth.

**Avoid poisoning!**

This medicine, and all other medicines, must be stored in a safe place out of the reach of children and/or infants, to avoid poisoning. In case of overdose, or if a child has accidentally swallowed the medicine, proceed immediately to a hospital emergency room and bring the package of the medicine with you.

Do not induce vomiting without explicit instructions from the physician!

Do not take any medicine in the dark! Carefully check the label and dose each time you take the medicine. Wear your glasses if you need them.

**Storage:** Keep tightly closed at room temperature - 25°C.

Even if kept in their original container and stored as recommended, medicines may be kept for a limited period only. Please note the expiry date of the medicine! In case of doubt, consult the pharmacist who dispensed the medicine to you.

Do not store different medicines in the same container.

The information appearing in this leaflet is based on the reference: MARTINDALE

**Drug registration no.:** 25-17-21245

**Manufacturer:** Sam-On Ltd., 25 Ha-avoda St., Bat-Yam.