

CARDOXIN FORTE

Tablets

Composition:

Active ingredient

Each tablet contains: Dipyridamole 75 mg

Inactive ingredients

Starch corn, silicon dioxide colloidal, starch soluble, magnesium stearate, calcium phosphate dibasic, lactose, hydroxypropylmethylcellulose, titanium dioxide, macrogol, sunset yellow 85 (E110).

Each tablet contains approximately 2 mg of lactose.

Therapeutic group: Anticoagulants.

Therapeutic activity: Cardoxin is an anticoagulant (anti-thrombotic) medicine which affects platelet activity to inhibit aggregation and adhesion.

When should this medicine not be used?

Do not use this medicine if you are hypersensitive to any of its ingredients.

Do not take this medicine without consulting a doctor before starting treatment, in the following cases:

If you are pregnant or breastfeeding.

If you suffer or have suffered in the past from impaired function of the: heart and/or vascular system (angina pectoris, blood circulation disturbances, heart valve problems, if you recently suffered a heart attack), kidney/urinary tract, blood system (e.g. bleeding, coagulation disorders) or if you suffer or have suffered in the past from hypotension or myasthenia gravis (muscle weakness).

Warnings:

If you are sensitive to any type of food or medicine, inform your doctor before starting treatment with this medication.

The tablets contain lactose. If you are sensitive to lactose, inform your doctor before taking this medication.

Avoid sudden changes from a lying/sitting position to a standing position in order to prevent dizziness, and in extreme cases, fainting.

If you are about to undergo any type of surgery or heart examination which includes injecting dipyridamole— inform the doctor about using this medicine.

Drug interactions:

If you are taking any other medications including non-prescription drugs and nutrition supplements, or if you have recently finished treatment with another drug, please notify your doctor in order to prevent any potential risks or a potential reduction in efficacy that may result from drug interactions. In particular, medicines of the following therapeutic groups may interact with your medicine: adenosine, antacids, aspirin and salicylates, anticoagulants (e.g. warfarin), thrombolytic medicines, medicines which prevent platelet aggregation, medicines to lower blood pressure, cholinesterase inhibitors (e.g. medicines for the treatment of myasthenia gravis), non-steroidal anti-inflammatory drugs.

Side effects:

In addition to the desired effect of the medicine, side effects may also appear during the course of taking this medicine. These may include: skin rash, nausea/vomiting, headache, dizziness, weakness, abdominal pain, indigestion, diarrhea, muscle pain, hot flushes, rapid heartbeat, low blood pressure.

These side effects usually pass within a short time following the period of adaptation to the medicine.

If these side effects persist and/or are bothersome, please consult your doctor.

The following side effects require special attention:

Hypersensitivity (may be manifested as edema, runny nose, shortness of breath, and as rash and itching), fainting, irregular heart rate, bruises/wounds, prolonged bleeding of wounds or following surgery, discomfort or pain in the chest (rare) – Stop the treatment and refer to your doctor immediately!

If at any time you experience side effects that are not mentioned in this leaflet or if you feel an adverse change in your general health, consult your doctor immediately!

Dosage:

This medicine is usually not intended for infants and children.

Administer according to the doctor's instructions only.

Do not exceed the recommended dose.

This medicine should be administered at set intervals according to your doctor's instructions.

If you forget to take your medicine at the proper time, take the missed dose as soon as you remember. If it is time to take the next dose, skip the missed dose. Never take two doses at the same time!

Directions for use:

Do not chew or crush the tablet! Swallow the medicine with a glass of water.

It is recommended to take the medicine before meal.

Do not retain the medicine in your mouth beyond the time necessary to swallow it.

Do not cut the tablet in half since there is no scored line.

How can you contribute to the success of the treatment?

Complete the entire course of treatment as instructed by your doctor.

Even if your condition improves do not stop taking this drug without consulting your doctor.

Avoid poisoning!

This medicine, and all other medicines, must be stored in a safe place out of the reach of children and/or infants, to avoid poisoning.

If you have taken an overdose, or if a child has accidentally swallowed the medicine, proceed immediately to a hospital emergency room and bring the package of the medicine with you.

Do not induce vomiting unless explicitly instructed to do so by a doctor!

This medicine was prescribed for the treatment of your ailment; in another patient it may cause harm. Do not give this medicine to your relatives, neighbors or acquaintances.

Do not take medicines in the dark! Carefully check the label and dose each time you take the medicine.

Wear your glasses if you need them.

Storage:

This medicine should be stored below 25°C.

Even if stored properly according to the manufacturer's recommendations, all medicines have a limited shelf life. Please pay attention to the expiry date of your medicine! If you have any doubts, please consult the dispensing pharmacist.

Do not store different medicines in the same container.

Drug registration number: 0271921428

Manufacturer and registration holder:



Rafa Laboratories Ltd.,
P.O.Box 405, Jerusalem 91003

The format of this leaflet was determined by the Ministry of Health that checked and approved its content in May 2012.

Cardoxin Forte PIL PB0612-03