

ספטמבר 2023

Iomeron 300, 350, 400 mg, Solution for Injection

צוות רפואי נכבד,

חברת דקסל בע"מ מבקשת להודיעכם על עדכון בעלון לרופא של התכשירים: אימרון 300, 350, 400.

בהודעה זו מפורטים העדכונים המהווים החמרה במידע הבטיחותי בלבד. למידע מלא, יש לעיין בעלון. העלון לרופא נשלח לפרסום במאגר התרופות שבאתר משרד הבריאות וניתן לקבלו מודפס ע"י פנייה לבעל הרישום: דקסל בע"מ, רח' דקסל 1, אור עקיבא 3060000, ישראל, טל': 04-6364000.

הרכב התכשיר:

Each 100 ml of solution contains iomeprol 61.24 g, 71.44 g, 81.65 g respectively.

התוויות מאושרות:

Contrast medium for diagnostic radiology.

Iomeron 300: Intravenous urography (in adults and paediatrics), peripheral phlebography, CT (brain and body), cavernosography, intravenous DSA, conventional angiography, intraarterial DSA, angiocardiology (in adults and paediatrics), conventional selective coronary arteriography, interventional coronary arteriography, ERCP, arthrography, hysterosalpingography, fistulography, discography, galactography, cholangiography, dacryocystography, sialography, retrograde urethrography, retrograde pyelo-ureterography.

Iomeron 350: Intravenous urography (in adults and paediatrics), CT (body), intravenous DSA, conventional angiography, intraarterial DSA, angiocardiology (in adults and paediatrics), conventional selective coronary arteriography, interventional coronary arteriography, arthrography, hysterosalpingography, fistulography, galactography, retrograde cholangiography, dacryocystography, sialography.

Iomeron 400: Intravenous urography (in adults including those with renal impairment or diabetes), CT (body), conventional angiography, intraarterial DSA, angiocardiology (in adults and paediatrics), conventional selective coronary arteriography, interventional coronary arteriography, fistulography, galactography, dacryocystography, sialography.

העלון לרופא עודכן בספטמבר 2023. להלן העדכונים המהווים החמרה במידע הבטיחותי (מסומנים באדום):

4.4 Special warnings and precautions for use

...

Disturbed thyroid function

Thyroid function tests indicative of hypothyroidism or transient thyroid suppression have been reported following iodinated contrast media administration to adult and paediatric patients, including infants. Some patients were treated for hypothyroidism. See also section on Special population-Children.

Special population

...

Children:

...

Special attention should be paid to paediatric patients below 3 years of age because an incident underactive thyroid during early life may be harmful for motor, hearing, and cognitive development and may require transient T4 replacement therapy. The incidence of hypothyroidism in patients younger than 3 years of age exposed to iodinated contrast media has been reported between 1.3% and 15% depending on the age of the subjects and the dose of the iodinated contrast agent and is more commonly observed in neonates and premature infants. Neonates may also be exposed through the mother during pregnancy. Thyroid function should be evaluated in all paediatric patients younger than 3 years of age following exposure to iodinated contrast media. If hypothyroidism is detected, the need for treatment should be considered and thyroid function should be monitored until normalized.

...

4.6 Fertility, pregnancy and lactation

...

Pregnancy

In neonates who have been exposed to iodinated contrast media in utero, it is recommended to monitor thyroid function (see section 4.4).

...