

אוקטובר 2023

רופא/ה, רוקח/ת נכבד/ה,
Enbrel 50 mg - ו Enbrel 25 mg solution for injection בrzinno להודיע על עדכון בעלונים לרופא ולצריך של :solution for injection

Etanercept 50 mg/ml

Solution for injection

המרכיב הפעיל:

צורת מינון:

התויה:

Rheumatoid arthritis

Enbrel is indicated for the treatment of active rheumatoid arthritis in adults when the response to disease-modifying antirheumatic drugs (DMARDs) including methotrexate (unless contraindicated) has been inadequate.

Enbrel can be used in combination with methotrexate in patients who do not respond adequately to methotrexate alone.

Reducing signs and symptoms and inhibiting the progression of structural damage in patients with moderately to severely active rheumatoid arthritis.

Enbrel, alone or in combination with methotrexate, has been shown to reduce the rate of progression of joint damage as measured by X-ray and to improve physical function.

Juvenile idiopathic arthritis

Treatment of polyarthritis (rheumatoid factor positive or negative) and extended oligoarthritis in children and adolescents from the age of 2 years who have had an inadequate response to, or who have proved intolerant of, methotrexate.

Treatment of psoriatic arthritis in adolescents from the age of 12 years who have had an inadequate response to, or who have proved intolerant of, methotrexate.

Treatment of enthesitis-related arthritis in adolescents from the age of 12 years who have had an inadequate response to, or who have proved intolerant of, conventional therapy.

Enbrel has not been studied in children aged less than 2 years.

Psoriatic arthritis

Treatment of active and progressive psoriatic arthritis in adults when the response to previous disease-modifying antirheumatic drug therapy has been inadequate. *Enbrel has been shown to improve physical function in patients with psoriatic arthritis, and to reduce the rate of progression of peripheral joint damage as measured by X-ray in patients with polyarticular symmetrical subtypes of the disease.*

Axial spondyloarthritis

Ankylosing spondylitis (AS)

Treatment of adults with severe active ankylosing spondylitis who have had an inadequate response to conventional therapy.

Non-radiographic axial spondyloarthritis

Treatment of adults with severe non-radiographic axial spondyloarthritis with objective signs of inflammation as indicated by elevated C-reactive protein (CRP) and/or magnetic resonance imaging (MRI) evidence, who have had an inadequate response to nonsteroidal anti-inflammatory drugs (NSAIDs).

Plaque psoriasis

Treatment of adults patients (18 years or older) with moderate to severe plaque psoriasis who are candidates for systemic therapy or phototherapy.

Paediatric plaque psoriasis

Treatment of chronic severe plaque psoriasis in children and adolescents from the age of 6 years who are inadequately controlled by, or are intolerant to, other systemic therapies or phototherapies.

להלן העדכוניים העיקריים בעלון לרופא:

4.6 FERTILITY, PREGNANCY AND LACTATION

...

Breast-feeding

~~Etanercept has been reported to be excreted in human milk following subcutaneous administration. In lactating rats following subcutaneous administration, etanercept was excreted in the milk and detected in the serum of pups. Because immunoglobulins, in common with many medicinal products, can be excreted. Limited information from the published literature indicates etanercept has been detected at low levels in human milk; a decision must be made whether to discontinue. Etanercept could be considered for use during breast-feeding or to discontinue Enbrel therapy, taking into account the benefit of breast-feeding for the child and the benefit of therapy for the woman.~~

~~While systemic exposure in a breastfed infant is expected to be low because etanercept is largely degraded in the gastrointestinal tract, limited data regarding systemic exposure in the breastfed infant are available.~~

~~Therefore, the administration of live vaccines (e.g., BCG) to a breastfed infant when the mother is receiving etanercept could be considered 16 weeks after stopping breast-feeding (or at an earlier timepoint if the infant etanercept serum levels are undetectable).~~

השינויים המודגשים ברקע צהוב מהווים החמרה. כמו כן, בוצעו שינויים נוספים הכוללים תוספת מידע, השמטת מידע ועדכוני נוסח שאינם מהווים החמרה.

העלונים המעודכנים נשלחו למשרד הבריאות לצורך פרסום במאגר התרופות שבאתר משרד הבריאות:
[חיפוש במאגר התרופות | משרד הבריאות \(www.gov.il\)](http://www.gov.il)

לחילופין, קיבלת עלון מלא מודפס ניתן לפנות לחברת פייזר פרמצטטיקה ישראל בע"מ, שנקר 9, ת.ד. 12133, הרצליה פיתוחה, 46725.

ברכה,
מרגריטה פולישצ'יק
רוקחת ממונה