TOUJEO (INSULIN GLARGINE 300 U/ML)

Toujeo SoloStar pre-filled pen contains 450 Units in 1.5 mL Toujeo DoubleStar pre-filled pen contains 900 Units in 3 mL


GUIDE FOR PATIENTS AND/OR CARERS:

- If you are prescribed Toujeo you must be trained on how to use the Toujeo pen by your healthcare professional and you must read the full Instructions For Use leaflet accompanying the pen.
- This document is provided as a guide only.
- Start with the dose in units written on your medical prescription.

Important information during the initial weeks when prescribed Toujeo

- Insulin glargine 300 U/mL (Toujeo) and insulin glargine 100 U/mL are **not bioequivalent and are therefore not directly interchangeable.**
 - **Dose adjustment may be required** (this means the drugs are not the same and you must not switch between them without adjusting the dose by your healthcare provider).
- The switch between insulin therapies should only be done when prescribed by your healthcare provider. The dose recommended by your healthcare provider should always be followed.
- Your blood glucose needs to be monitored closely during the switch and the initial weeks thereafter.
- Insulin glargine 300 U/mL is available in two different presentations:





The dose increment in Toujeo SoloStar is 1 U

The dose increment in Toujeo DoubleStar is 2 U

The dose delivered is the one shown in the dose window.

Please consult your healthcare professional for further information.

Before injecting Toujeo read the Instructions for Use that come with the package leaflet.

If you get any medication error or side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet.

The format and content of this Patient guide was checked and approved by the Ministry of Health in September 2023. Suspected adverse reactions could be reported to the Ministry of Health by using an online form https://sideeffects.health.gov.il. Also could be reported directly to Sanofi Company 09-8633081

