

## **Veterinary medicine package leaflet**

This medicine is supplied on veterinary prescription only  
For use in animals only

### **1. Name, form, and strength of the veterinary medicine**

**Canidryl 20 mg Tablets for Dogs Veterinary**  
**Canidryl 100 mg Tablets for Dogs Veterinary**

### **2. Active ingredient**

Each oral grilled-meat flavored tablet contains:

Canidryl 20 mg tablets for dogs Veterinary contains 20 mg carprofen

Canidryl 100 mg tablets for dogs Veterinary contains 100 mg carprofen

Inactive ingredients and allergens: See Section 13 'Additional Information'.

### **3. What is this medicine intended for**

This medicine is intended for use in dogs, to reduce inflammation and pain caused by musculoskeletal disorders and degenerative joint disease.

As a follow up to parenteral analgesia in the management of post-operative pain.

**Therapeutic group:** non-steroid anti-inflammatory (NSAIDs).

### **4. Contraindications**

Do not administer this medicine in cases of hypersensitivity to the active ingredient or to any of the inactive ingredients.

Do not give this medicine to cats. Mistakenly using the medicine in cats can bring them to a life threatening situation.

Do not exceed the recommended dose.

Never give this medicine to dogs with heart, liver, or kidney diseases where there is a possibility of bleeding or ulcers in the digestive tract, or if there is evidence of impaired blood composition (blood dyscrasia).

Do not give this medicine to puppies younger than 4 months.

Do not use in pregnant or lactating bitches.

### **5. Side effects**

Undesirable effects associated with NSAIDs have been reported such as vomiting, soft stool/diarrhea, occult blood in the stool, loss of appetite, and exhaustion. These side effects usually occur within the first week of treatment and, in most cases, disappear following termination of the treatment. However, in very rare cases these effects may be serious or fatal. If side effects occur, stop using the drug and consult a veterinarian.

As with other NSAIDs, there is a risk of rare kidney or idiosyncratic liver side effects.

Side effects can be reported to the Ministry of Health by pressing the link 'Reporting Side Effects of Medication' on the Ministry of Health home page ([www.health.gov.il](http://www.health.gov.il)) which links to an online form for reporting side effects, or by entering the link:

<https://sideeffects.health.gov.il>

### **6. Target animals**

Dogs

### **7. Dose and administration**

Recommended dose:

2 to 4 mg carprofen per kg body weight per day.

Starting dose of 4 mg per kg per day, once a day or divided into two equal doses.

Depending on the clinical response, after 7 days the dose can be reduced to 2 mg carprofen per kg body weight per day, administered in a single dose.

Canidryl 20 mg tablets for dogs veterinary: the tablets can be divided into equal halves.

Canidryl 100 mg tablets for dogs veterinary: the tablets can be divided into equal halves or quarters.

## 8. How to use this medicine

Duration of treatment with this medicine depends on the response to treatment.

For oral administration only.

Do not exceed the recommended dose.

## 9. Withdrawal period

Not applicable

## 10. Warnings

- Special warnings about using the medicine in the target animal

There may be additional risks when treating elderly dogs with this medicine. If using this medicine is unavoidable, these dogs may need a lower dose and meticulous medical monitoring.

Avoid use in dogs that are dehydrated, hypovolemic, or have low blood pressure because there is a possible risk of increased kidney toxicity.

NSAID medicines can cause inhibition of phagocytosis. So, when treating inflammations associated with a bacterial infection, add a suitable antibacterial treatment.

These chewable tablets contain flavorings. To avoid animals taking this medicine accidentally, keep the tablets out of the animals' reach.

- Special safety precautions for the person administering the medicine

If a person accidentally swallows a tablet, seek medical advice and show the doctor this medicine's leaflet. Wash hands after contact with this medicine.

- Gestating and lactating animal patients

Studies in laboratory animals (rats and rabbits) have shown evidence that the drug is foetotoxic at a dose close to the therapeutic dose.

The safety of the medicine has not been proven during pregnancy or lactation.

Do not give this medicine to gestating or lactating bitches.

Do not use in bitches during the estrus period.

- Interaction with other medicinal products and other forms of interaction:

Carprofen is highly bound to plasma proteins and competes with other medicines that are also highly bound to plasma proteins, which can increase their respective toxic effects. Do not use Canidryl together with, or at intervals shorter than 24 hours from, other NSAID medicines. Do not use Canidryl together with glucocorticoids.

Avoid giving this medicine together with other medicines that are potentially nephrotoxic (toxic to kidneys).

Do not give Canidryl together with anticoagulants.

- Overdose

No signs of toxicity were observed in dogs given a dose of up to 6 mg/kg twice a day for 8 days (3 times the maximum recommended dose of 4 mg/kg/day) and 6 mg/kg once a day for 7 additional days (1.5 times the maximum recommended dose of 4 mg/kg/day).

There is no antidote to overdose of carprofen. General supportive care must be given, as in cases of NSAIDs overdose.

Serious side effects may occur when taking large amounts. If you suspect that your dog has consumed a quantity of tablets that exceeds the recommended dose, consult your veterinarian.

## 11. Storage instructions

- Prevent poisoning! To prevent poisoning, keep this, and all other medicines, in a closed place out of reach and sight of children and/or infants.
- Do not use this medicine after the expiry date (exp. Date) stated on the label and carton. The expiry date refers to the last day of that month.
- Do not store above 25°C.
- Store in a dry place in the original package.
- Protect from light.
- Canidryl 20 mg tablets for dogs veterinary: halved tablets that were not used must be discarded immediately.
- Canidryl 100 mg tablets for dogs veterinary: divided tablets should be stored back in the blister and used within 72 hours.

## 12. Instructions for destroying the medicine/any remaining medicine after use:

Destroy as toxic waste any unused veterinary medicinal product or any excess veterinary medicinal product. Do not dispose of the medicine in the sewage.

**13. Additional information**

- In addition to the active ingredients this medicine also contains:  
Lactose monohydrate, microcrystalline cellulose, colloidal silicon dioxide, magnesium stearate, grilled meat flavor.
- What the medicine looks like and what is the content of the package:  
Canidryl 20 mg tablets for dogs veterinary: A creamy-white round tablet, with a beveled edge and a score line on one side.  
Canidryl 100 mg tablets for dogs veterinary: A white to off-white round shape tablet with a cross break line on one side.  
Each package contains 100 tablets in blisters.

**Registration holder and importer:**

A.L. Medi-Market Ltd., 3 Hakatif Street, Emek Hefer Industrial Park, 3877701.

**Manufacturer Name:**

CHANELLE Pharmaceuticals Ltd., Loughrea, Co. Galway, Ireland.

Revised in August 2023 according to MOHs guidelines.

**Registration number of the medicine in the Ministry of Health National Drug Registry:**

**Canidryl 20 mg tablets for dogs Veterinary: 160-20-35099**

**Canidryl 100 mg tablets for dogs Veterinary: 153-51-33972**