

Atazanavir Teva 150 mg

Atazanavir Teva 200 mg

Atazanavir Teva 300 mg

(ATAZANAVIR as SULFATE) CAPSULES

Per os

1. INDICATIONS AND USAGE

Atazanavir Teva (atazanavir as sulfate) is indicated in combination with other antiretroviral agents for the treatment of HIV-1 infection.

This indication is based on analyses of plasma HIV-1 RNA levels and CD4+ cell counts from controlled studies of 96 weeks duration in antiretroviral-naïve and 48 weeks duration in antiretroviral-treatment-experienced adult and pediatric patients at least 6 years of age.

The following points should be considered when initiating therapy with Atazanavir Teva:

- In Study AI424-045, atazanavir /ritonavir and lopinavir/ritonavir were similar for the primary efficacy outcome measure of time-averaged difference in change from baseline in HIV RNA level. This study was not large enough to reach a definitive conclusion that atazanavir /ritonavir and lopinavir/ritonavir are equivalent on the secondary efficacy outcome measure of proportions below the HIV RNA lower limit of detection [see Clinical Studies (13.2)].

The number of baseline primary protease inhibitor mutations affects the virologic response to atazanavir /ritonavir [see Clinical Pharmacology (11.4)].

2. DOSAGE AND ADMINISTRATION

2.1. overview

- Atazanavir Teva Capsules must be taken with food.
- Do not open the capsules.
- The recommended oral dosage of Atazanavir Teva depends on the treatment history of the patient and the use of other coadministered drugs. When coadministered with H₂-receptor antagonists or proton-pump inhibitors, dose separation may be required [see Dosage and Administration (2.3,2.4,2.5 and 2.6) and drug interactions (7)].
- When coadministered with didanosine buffered or enteric-coated formulations, Atazanavir Teva should be given (with food) 2 hours before or 1 hour after didanosine.
- Atazanavir Teva without ritonavir is not recommended for treatment-experienced adult or pediatric patients with prior virologic failure [see Clinical Studies (13)].
- Efficacy and safety of Atazanavir Teva with ritonavir when ritonavir is administered in doses

greater than 100 mg once daily have not been established. The use of higher ritonavir doses may alter the safety profile of atazanavir (cardiac effects, hyperbilirubinemia) and, therefore, is not recommended. Prescribers should consult the complete prescribing information for ritonavir when using ritonavir.

2.2. Testing Prior to Initiation and During Treatment with Atazanavir Teva

Renal laboratory testing should be performed in all patients prior to initiation of Atazanavir Teva and continued during treatment with Atazanavir Teva. Renal laboratory testing should include serum creatinine, estimated creatinine clearance, and urinalysis with microscopic examination [see Warnings and Precautions (5.5, 5.6)].

Hepatic laboratory testing should be performed in patients with underlying liver disease prior to initiation of Atazanavir Teva and continued during treatment with Atazanavir Teva [see Warnings and Precautions (5.4)].

2.3. Dosage of Atazanavir Teva in Adult patients

Table 1 summarizes the recommended Atazanavir Teva dosing regimen in adults. All Atazanavir Teva dosing regimens are to be administered as a single dose with food.

Table 1: Atazanavir Teva Dosing Regimens

Treatment-Naive Patients	Atazanavir Teva 300 mg with ritonavir 100 mg once daily
If unable to tolerate ritonavir	Atazanavir Teva 400 mg once daily
When combined with any of the following: Tenofovir H2-receptor antagonist Proton-pump inhibitor	Atazanavir Teva 300 mg with ritonavir 100 mg once daily
<ul style="list-style-type: none"> The H2-receptor antagonist dose should not exceed a dose comparable to famotidine 40 mg twice daily. Administer Atazanavir Teva and ritonavir simultaneously with, and/or at least 10 hours after the H2-receptor antagonist 	
<ul style="list-style-type: none"> If unable to tolerate ritonavir, administer Atazanavir Teva 400 mg once daily at least 2 hours before and at least 10 hours after the H2-receptor antagonist. No single dose of the H2-receptor antagonist should exceed a dose comparable to famotidine 20 mg and the total daily dose should not exceed a dose comparable to famotidine 40 mg. 	
<ul style="list-style-type: none"> The proton-pump inhibitor dose should not exceed a dose comparable to omeprazole 20 mg daily and must be taken approximately 12 hours prior to Atazanavir Teva and ritonavir. 	

When combined with efavirenz	Atazanavir Teva 400 mg with ritonavir 100 mg once daily
<ul style="list-style-type: none"> Efavirenz should be administered on an empty stomach, preferably at bedtime. 	
Treatment-Experienced Patients	Atazanavir Teva 300 mg with ritonavir 100 mg once daily
Do not coadminister with proton-pump inhibitors or efavirenz in treatment-experienced patients.	
When given with an H2-receptor antagonist	Atazanavir Teva 300 mg with ritonavir 100 mg once daily
<ul style="list-style-type: none"> The H2-receptor antagonist dose should not exceed a dose comparable to famotidine 20 mg twice daily. Administer Atazanavir Teva and ritonavir simultaneously with, and/or at least 10 hours after the H2-receptor antagonist. 	
When given with both tenofovir and an H2-receptor antagonist	Atazanavir Teva 400 mg with ritonavir 100 mg once daily
The H2-receptor antagonist dose should not exceed a dose comparable to famotidine 20 mg twice daily. Administer Atazanavir Teva and ritonavir simultaneously with, and/or at least 10 hours after the H2-receptor antagonist.	

[For these drugs and other antiretroviral agents for which dosing modification may be appropriate, see Drug Interactions (7).]

2.4. Dosage of Atazanavir Teva in pediatric patients

The recommended daily dosage of Atazanavir Teva for pediatric patients (6 to less than 18 years of age) is based on body weight and should not exceed the recommended adult dosage. Atazanavir Teva Capsules must be taken with food. The data are insufficient to recommend dosing of Atazanavir Teva for any of the following: (1) patients less than 6 years of age, (2) without ritonavir in any pediatric patient less than 13 years of age, and (3) patients less than 40 kg receiving concomitant tenofovir, H2-receptor antagonists, or proton-pump inhibitors.

The recommended dosage of Atazanavir Teva with ritonavir in pediatric patients at least 6 years of age is shown in Table 2.

Table 2: Dosage for Pediatric Patients (6 to less than 18 years of age) for Atazanavir Teva Capsules with ritonavir^a

Body Weight	Atazanavir Teva dose	ritonavir dose
15 kg to less than 20 kg	150 mg	100 mg

20 kg to less than 40 kg	200 mg	100 mg
at least 40 kg	300 mg	100 mg

^a The Atazanavir Teva and ritonavir dose should be taken together once daily with food.

For treatment-naïve patients at least 13 years of age and at least 40 kg, who are unable to tolerate ritonavir, the recommended dose is Atazanavir Teva 400 mg (without ritonavir) once daily with food. For patients at least 13 years of age and at least 40 kg receiving concomitant tenofovir, H2-receptor antagonists, or proton-pump inhibitors, Atazanavir Teva should not be administered without ritonavir.

Pregnancy Dosing During and the Postpartum Period:

- Atazanavir Teva should not be administered without ritonavir.
- Atazanavir Teva should only be administered to pregnant women with HIV-1 strains susceptible to atazanavir.
- For pregnant patients, no dose adjustment is required for Atazanavir Teva with the following exceptions:
 - For treatment-experienced pregnant women during the second or third trimester, when Atazanavir Teva is coadministered with either an H2-receptor antagonist or tenofovir, Atazanavir Teva 400 mg with ritonavir 100 mg once daily is recommended. There are insufficient data to recommend a Atazanavir Teva dose for use with both an H2-receptor antagonist and tenofovir in treatment-experienced pregnant women.
- No dose adjustment is required for postpartum patients. However, patients should be closely monitored for adverse events because atazanavir exposures could be higher during the first 2 months after delivery. [See Use in Specific Populations (8.1) and Clinical Pharmacology (11.3)]

2.5. Renal Impairment

For patients with renal impairment, including those with severe renal impairment who are not managed with hemodialysis, no dose adjustment is required for Atazanavir Teva. Treatment-naïve patients with end stage renal disease managed with hemodialysis should receive Atazanavir Teva 300 mg with ritonavir 100 mg. Atazanavir Teva should not be administered to HIV-treatment- experienced patients with end stage renal disease managed with hemodialysis. [See Use in Specific Populations (8.7).]

2.6. Hepatic Impairment

Atazanavir Teva should be used with caution in patients with mild-to-moderate hepatic impairment. For patients with moderate hepatic impairment (Child-Pugh Class B) who have not experienced prior virologic

failure, a dose reduction to 300 mg once daily should be considered. Atazanavir Teva should not be used in patients with severe hepatic impairment (Child-Pugh Class C).

Atazanavir Teva/ritonavir has not been studied in subjects with hepatic impairment and is not recommended. [See Warnings and Precautions (5.5) and Use in Specific Populations (8.8).]

3. DOSAGE FORMS AND STRENGTHS

Atazanavir Teva Capsules:

- 150 mg capsule –non transparent capsule, with dark blue cap and black mark 150 on light blue body.
- 200 mg capsule- non transparent capsule, with blue cap and black mark 200 on blue body.
- 300 mg capsule- non transparent capsule, with red cap and black mark 300 on blue body.

4. CONTRAINDICATIONS

Atazanavir Teva is contraindicated:

- Hypersensitivity to the active substance or to any of the excipients list in section description (11)
- in patients with previously demonstrated clinically significant hypersensitivity (eg, Stevens-Johnson syndrome, erythema multiforme, or toxic skin eruptions) to any of the components of Atazanavir Teva capsules [see Warnings and Precautions (5)].
- when coadministered with drugs that are highly dependent on CYP3A or UGT1A1 for clearance, and for which elevated plasma concentrations of the interacting drugs are associated with serious and/or life-threatening events (see Table 6).
- when coadministered with drugs that strongly induce CYP3A and may lead to lower exposure and loss of efficacy of Atazanavir Teva (see Table 6).

Table 6 displays drugs that are contraindicated with Atazanavir Teva.

Table 6: Drugs Contraindicated with Atazanavir Teva (Information in the table applies to Atazanavir Teva with or without ritonavir, unless otherwise indicated)

Drug Class	Drugs within class that are contraindicated with Atazanavir Teva
Alpha 1- Adrenoreceptor Antagonist	Alfuzosin
Antiarrhythmics	Amiodarone (with ritonavir), quinidine (with ritonavir)
Anticonvulsants	Carbamazepine, phenobarbital, phenytoin
Antimycobacterials	Rifampin
Antineoplastics	Apalutamide, encorafenib, irinotecan , ivosidenib

Antipsychotics	Lurasidone (with ritonavir), pimozide
Benzodiazepines	Orally administered midazolam ^a , triazolam
Ergot Derivatives	Dihydroergotamine, ergotamine, ergonovine, methylergonovine
GI Motility Agent	Cisapride
Hepatitis C Direct-Acting Antivirals	Elbasvir/grazoprevir; glecaprevir/pibrentasvir
Herbal Products	St. John's wort (<i>Hypericum perforatum</i>)
Lipid-Modifying Agents	Lovastatin, simvastatin, lomitapide
Phosphodiesterase-5 (PDE-5) Inhibitor	Sildenafil ^b when dosed as Atazanavir Teva for the treatment of pulmonary arterial hypertension
Protease Inhibitors	Indinavir
Non-nucleoside Reverse Transcriptase Inhibitors	Nevirapine

^a See Drug Interactions, Table 16 (7) for parenterally administered midazolam.

^b See Drug Interactions, Table 16 (7) for sildenafil* when dosed as VIAGRA® for erectile dysfunction.

5. WARNINGS AND PRECAUTIONS

5.1. Cardiac Conduction Abnormalities

Atazanavir Teva has been shown to prolong the PR interval of the electrocardiogram in some subjects. In healthy subjects and in subjects with HIV-1 infection treated with atazanavir, abnormalities in atrioventricular (AV) conduction were asymptomatic and generally limited to first-degree AV block. There have been reports of second-degree AV block and other conduction abnormalities [see Adverse Reactions (6.2) and Overdosage (9)]. In clinical trials that included electrocardiograms, asymptomatic first-degree AV block was observed in 5.9% of atazanavir-treated subjects (n=920), 5.2% of lopinavir/ritonavir-treated subjects (n=252), 10.4% of nelfinavir-treated subjects (n=48), and 3.0% of efavirenz-treated subjects (n=329). In Study AI424-045, asymptomatic first-degree AV block was observed in 5% (6/118) of atazanavir with ritonavir-treated subjects and 5% (6/116) of lopinavir/ritonavir-treated subjects who had on-study electrocardiogram measurements. Because of limited clinical experience in those with preexisting conduction system disease (eg, marked first-degree AV block or second- or third-degree AV block). ECG monitoring should be considered in these patients [see Clinical Pharmacology (11.2)].

5.2. Severe Skin Reactions

In controlled clinical trials, rash (all grades, regardless of causality) occurred in approximately 20% of

subjects with HIV-1 infection treated with atazanavir . The median time to onset of rash in clinical studies was 7.3 weeks and the median duration of rash was 1.4 weeks. Rashes were generally mild-to- moderate maculopapular skin eruptions. Treatment-emergent adverse reactions of moderate or severe rash (occurring at a rate of $\geq 2\%$) are presented for the individual clinical studies [see Adverse Reactions (6.1)]. Dosing with atazanavir was often continued without interruption in patients who developed rash. The discontinuation rate for rash in clinical trials was $< 1\%$. Cases of Stevens-Johnson syndrome, erythema multiforme, and toxic skin eruptions, including drug rash, eosinophilia, and systemic symptoms (DRESS) syndrome, have been reported in patients receiving atazanavir [see Contraindications (4) and Adverse Reactions (6.1)]. Atazanavir Teva should be discontinued if severe rash develops.

5.3. Hepatotoxicity

Patients with underlying hepatitis B or C viral infections or marked elevations in transaminases before treatment may be at increased risk for developing further transaminase elevations or hepatic decompensation. In these patients, hepatic laboratory testing should be conducted prior to initiating therapy with Atazanavir Teva and during treatment [see Dosage and Administration (2.2), Adverse Reactions (6.1), and Use in Specific Populations (8.8)].

5.4. Chronic Kidney Disease

Chronic kidney disease in patients with HIV-1 infection treated with atazanavir, with or without ritonavir, has been reported during postmarketing surveillance. Reports included biopsy-proven cases of granulomatous interstitial nephritis associated with the deposition of atazanavir drug crystals in the renal parenchyma. Consider alternatives to Atazanavir Teva in patients at high risk for renal disease or with preexisting renal disease. Renal laboratory testing (including serum creatinine, estimated creatinine clearance, and urinalysis with microscopic examination) should be conducted in all patients prior to initiating therapy with Atazanavir Teva and continued during treatment with Atazanavir Teva. Expert consultation is advised for patients who have confirmed renal laboratory abnormalities while taking Atazanavir Teva. In patients with progressive kidney disease, discontinuation of Atazanavir Teva may be considered [see Dosage and Administration (2.2 and 2.7) and Adverse Reactions (6.2)].

5.5. Nephrolithiasis and Cholelithiasis

Cases of nephrolithiasis and/or cholelithiasis have been reported during postmarketing surveillance in patients with HIV-1 infection receiving atazanavir therapy. Some patients required hospitalization for additional management and some had complications. Because these events were reported voluntarily during clinical practice, estimates of frequency cannot be made. If signs or symptoms of nephrolithiasis

and/or cholelithiasis occur, temporary interruption or discontinuation of therapy may be considered [see Adverse Reactions (6.2)].

5.6. Risk of Serious Adverse Reactions Due to Drug Interactions

Initiation of Atazanavir Teva with ritonavir, a CYP3A inhibitor, in patients receiving medications metabolized by CYP3A or initiation of medications metabolized by CYP3A in patients already receiving Atazanavir Teva with ritonavir, may increase plasma concentrations of medications metabolized by CYP3A. Initiation of medications that inhibit or induce CYP3A may increase or decrease concentrations of Atazanavir Teva with ritonavir, respectively. These interactions may lead to:

- clinically significant adverse reactions potentially leading to severe, life threatening, or fatal events from greater exposures of concomitant medications.
- clinically significant adverse reactions from greater exposures of Atazanavir Teva with ritonavir.
- loss of therapeutic effect (virologic response) of Atazanavir Teva with ritonavir and possible development of resistance.

See Table 16 for steps to prevent or manage these possible and known significant drug interactions, including dosing recommendations [see Drug Interactions (7)]. Consider the potential for drug interactions prior to and during therapy containing Atazanavir Teva with ritonavir; and monitor for the adverse reactions associated with the concomitant medications [see Contraindications (4) and Drug Interactions (7)].

5.7. Hyperbilirubinemia

Most patients taking Atazanavir Teva experience asymptomatic elevations in indirect (unconjugated) bilirubin related to inhibition of UDP-glucuronosyl transferase (UGT). This hyperbilirubinemia is reversible upon discontinuation of Atazanavir Teva. Hepatic transaminase elevations that occur with hyperbilirubinemia should be evaluated for alternative etiologies. No long-term safety data are available for patients experiencing persistent elevations in total bilirubin >5 times the upper limit of normal (ULN). Alternative antiretroviral therapy to Atazanavir Teva may be considered if jaundice or scleral icterus associated with bilirubin elevations presents cosmetic concerns for patients. Dose reduction of atazanavir is not recommended since long-term efficacy of reduced doses has not been established [see Adverse Reactions (6.1)].

5.8. Diabetes Mellitus/Hyperglycemia

New-onset diabetes mellitus, exacerbation of preexisting diabetes mellitus, and hyperglycemia have been

reported during postmarketing surveillance in patients with HIV-1 infection receiving protease inhibitor therapy. Some patients required either initiation or dose adjustments of insulin or oral hypoglycemic agents for treatment of these events. In some cases, diabetic ketoacidosis has occurred. In those patients who discontinued protease inhibitor therapy, hyperglycemia persisted in some cases. Because these events have been reported voluntarily during clinical practice, estimates of frequency cannot be made and a causal relationship between protease inhibitor therapy and these events has not been established [see Adverse Reactions (6.2)].

5.9. Immune Reconstitution Syndrome

Immune reconstitution syndrome has been reported in patients treated with combination antiretroviral therapy, including Atazanavir Teva. During the initial phase of combination antiretroviral treatment, patients whose immune system responds may develop an inflammatory response to indolent or residual opportunistic infections (such as *Mycobacterium avium* infection, cytomegalovirus, *Pneumocystis jirovecii* pneumonia, or tuberculosis), which may necessitate further evaluation and treatment.

Autoimmune disorders (such as Graves' disease, polymyositis, and Guillain-Barre syndrome and autoimmune hepatitis) have also been reported to occur in the setting of immune reconstitution; however, the time to onset is more variable, and can occur many months after initiation of treatment.

5.10. Fat Redistribution

Redistribution/accumulation of body fat including central obesity, dorsocervical fat enlargement (buffalo hump), peripheral wasting, facial wasting, breast enlargement, and “cushingoid appearance” have been observed in patients receiving antiretroviral therapy. The mechanism and long-term consequences of these events are currently unknown. A causal relationship has not been established.

5.11. Hemophilia

There have been reports of increased bleeding, including spontaneous skin hematomas and hemarthrosis, in patients with hemophilia type A and B treated with protease inhibitors. In some patients, additional factor VIII was given. In more than half of the reported cases, treatment with protease inhibitors was continued or reintroduced. A causal relationship between protease inhibitor therapy and these events has not been established.

5.12. Resistance/Cross-Resistance

Various degrees of cross-resistance among protease inhibitors have been observed. Resistance to Atazanavir Teva 150/200/300mg KL SPC Notification 12-2023

atazanavir may not preclude the subsequent use of other protease inhibitors [see Microbiology (11.4)].

5.13. Excipients:

Lactose - Patients with rare hereditary problems of galactose intolerance, total lactase deficiency or glucosegalactose malabsorption should not take this medicine.

6. ADVERSE REACTIONS

The following adverse reactions are discussed in greater detail in other sections of the labeling:

- cardiac conduction abnormalities [see Warnings and Precautions (5.1)]
- rash [see Warnings and Precautions (5.2)]
- hyperbilirubinemia [see Warnings and Precautions (5.7)]
- chronic kidney disease [see Warnings and Precautions (5.4)]
- nephrolithiasis and cholelithiasis [see Warnings and Precautions (5.5)]

6.1. Clinical Trial Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

Adverse Reactions in Treatment-Naive Adult subjects

The safety profile of atazanavir in treatment-naive adults is based on 1625 subjects with HIV-1 infection in clinical trials. 536 subjects received atazanavir 300 mg with ritonavir 100 mg and 1089 subjects received atazanavir 400 mg or higher (without ritonavir).

The most common adverse reactions were nausea, jaundice/scleral icterus, and rash.

Selected clinical adverse reactions of moderate or severe intensity reported in $\geq 2\%$ of treatment-naive subjects receiving combination therapy including atazanavir 300 mg with ritonavir 100 mg and atazanavir 400 mg (without ritonavir) are presented in Tables 7 and 8, respectively.

Table 7: Selected Adverse Reactions^a of Moderate or Severe Intensity Reported in $\geq 2\%$ of Adult Treatment-Naive subjects with HIV-1 infection,^b Study AI424- 138

	96 weeks ^c atazanavir 300 mg with ritonavir 100 mg (once daily) and tenofovir DF / emtricitabine ^d (n=441)	96 weeks ^c Lopinavir/ ritonavir ^d 400 mg / 100 mg (twice daily) and tenofovir DF / emtricitabine ^e (n=437)
Digestive System		

Nausea	4%	8%
Jaundice/scleral icterus	5%	*
Diarrhea	2%	12%
Skin and Appendages		
Rash	3%	2%

* None reported in this treatment arm.

^a Includes events of possible, probable, certain, or unknown relationship to treatment regimen.

^b Based on the regimen containing atazanavir.

^c Median time on therapy.

^d Administered as a fixed-dose

^e As a fixed-dose product: 300 mg tenofovir DF, 200 mg emtricitabine once daily.

Table 8: Selected Adverse Reactions^a of Moderate or Severe Intensity Reported in $\geq 2\%$ of Adult Treatment-Naive subjects with HIV-1 infection,^b Studies AI424-034, AI424-007, and AI424-008

	Study AI424-034		Studies AI424-007, -008	
	64 weeks ^c atazanavir 400 mg once daily + lamivudine / zidovudinee (n=404)	64 weeks ^c efavirenz 600 mg once daily + lamivudine/zidovu dinee (n=401)	120 weeksc, ^d atazanavir 400 mg once daily + stavudine / lamivudine or didanosine (n=279)	73 weeksc, ^d nelfinavir 750 mg TID or 1250 mg BID + stavudine / lamivudine or didanosine (n=191)
Body as a Whole				
Headache	6%	6%	1%	2%
Digestive System				
Nausea	14%	12%	6%	4%
Jaundice/scleral icterus	7%	*	7%	*
Vomiting	4%	7%	3%	3%
Abdominal pain	4%	4%	4%	2%
Diarrhea	1%	2%	3%	16%
Nervous System				
Insomnia	3%	3%	<1%	*
Dizziness	2%	7%	<1%	*
Peripheral neurologic symptoms	<1%	1%	4%	3%
Skin and Appendages				
Rash	7%	10%	5%	1%

* None reported in this treatment arm.

^a Includes events of possible, probable, certain, or unknown relationship to treatment regimen.

^b Based on regimens containing atazanavir.

^c Median time on therapy.

^d Includes long-term follow-up.

^e As a fixed-dose product: 150 mg lamivudine/ 300 mg zidovudine twice daily.

Adverse Reactions in Treatment-Experienced Adult Subjects

The safety profile of atazanavir in treatment-experienced adults with HIV-1 infection is based on 119 subjects with HIV-1 infection in clinical trials.

The most common adverse reactions are jaundice/scleral icterus and myalgia.

Selected clinical adverse reactions of moderate or severe intensity reported in $\geq 2\%$ of treatment-experienced subjects receiving atazanavir with ritonavir are presented in Table 9.

Table 9: Selected Adverse Reactions^a of Moderate or Severe Intensity Reported in $\geq 2\%$ of Adult Treatment-Experienced subjects with HIV-1 infection,^b Study AI424- 045

	48 weeks^c Atazanavir with ritonavir 300/100 mg (once daily) and tenofovir DF and NRTI (n=119)	48 weeks^c lopinavir/ritonavir 400/100 mg (twice daily^d) and tenofovir DF and NRTI (n=118)
Body as a Whole		
Fever	2%	*
Digestive System		
Jaundice/scleral icterus	9%	*
Diarrhea	3%	11%
Nausea	3%	2%
Nervous System		
Depression	2%	<1%
Musculoskeletal System		
Myalgia	4%	*

* None reported in this treatment arm.

^a Includes events of possible, probable, certain, or unknown relationship to treatment regimen.

^b Based on the regimen containing atazanavir.

^c Median time on therapy.

^d As a fixed-dose product.

Laboratory Abnormalities in Treatment-Naive subjects

The percentages of adult treatment-naive subjects with HIV-1 infection treated with combination therapy,

including atazanavir 300 mg with ritonavir 100 mg or atazanavir 400 mg (without ritonavir) with Grade 3-4 laboratory abnormalities are presented in Tables 10 and 11, respectively.

Table 10: Grade 3-4 Laboratory Abnormalities Reported in $\geq 2\%$ of Adult Treatment-Naive Subjects with HIV-1 infection,^a Study AI424-138

Variable	Limit ^e	96 weeks ^b atazanavir 300 mg with ritonavir 100 mg (once daily) and tenofovir DF / emtricitabine ^c (n=441)	96 weeks ^b lopinavir 400 mg / ritonavir 100 mgc (twice daily) and tenofovir DF /emtricitabined (n=437)
Chemistry	High		
SGOT/AST	≥ 5.1 x ULN	3%	1%
SGPT/ALT	≥ 5.1 x ULN	3%	2%
Total Bilirubin	≥ 2.6 x ULN	44%	<1%
Lipase	≥ 2.1 x ULN	2%	2%
Creatine Kinase	≥ 5.1 x ULN	8%	7%
Total Cholesterol	≥ 240 mg/dL	11%	25%
Hematology	Low		
Neutrophils	<750 cells/mm ³	5%	2%

^a Based on the regimen containing atazanavir.

^b Median time on therapy.

^c Administered as a fixed-dose product

^d As a fixed-dose product: 300 mg tenofovir DF, 200 mg emtricitabine once daily.

^e ULN = upper limit of normal.

Table 11: Grade 3-4 Laboratory Abnormalities Reported in $\geq 2\%$ of Adult Treatment-Naive Subjects with HIV-1 infection,^a Studies AI424-034, AI424-007, and AI424- 008

Variable	Limit ^d	Study AI424-034		Studies AI424-007, -008	
		64 weeks ^b Atazanavir 400 mg once daily + lamivudine /zidovudine ^e (n=404)	64 weeks ^b efavirenz 600 mg once daily + lamivudine / zidovudine ^e (n=401)	120 weeks ^{b,c} atazanavir 400 mg once daily with stavudine + lamivudine or with stavudine + didanosine (n=279)	73 weeks ^{b,c} nelfinavir 750 mg TID or 1250 mg BID with stavudine + lamivudine or with stavudine + didanosine (n=191)
Chemistry	High				
SGOT/AST	≥ 5.1 x ULN	2%	2%	7%	5%
SGPT/ALT	≥ 5.1 x ULN	4%	3%	9%	7%

Total Bilirubin	≥2.6 x ULN	35%	<1%	47%	3%
Amylase	≥2.1 x ULN	*	*	14%	10%
Lipase	≥2.1 x ULN	<1%	1%	4%	5%
Creatine Kinase	≥5.1 x ULN	6%	6%	11%	9%
Total Cholesterol	≥240 mg/dL	6%	24%	19%	48%
Triglycerides	≥751 mg/dL	<1%	3%	4%	2%
Hematology	Low				
Hemoglobin	<8.0 g/dL	5%	3%	<1%	4%
Neutrophils	<750 cells/mm ³	7%	9%	3%	7%

* None reported in this treatment arm.

^a Based on regimen(s) containing atazanavir.

^b Median time on therapy.

^c Includes long-term follow-up.

^d ULN = upper limit of normal.

^e As a fixed-dose product: 150 mg lamivudine, 300 mg zidovudine twice daily.

Change in Lipids from Baseline in Treatment-Naive Subjects with HIV-1 infection

For Study AI424-138 and Study AI424-034, changes from baseline in LDL-cholesterol, HDL- cholesterol, total cholesterol, and triglycerides are shown in Tables 12 and 13, respectively.

Table 12: Lipid Values, Mean Change from Baseline, Study AI424-138

	Atazanavir/ritonavir ^{a,b}					Lopinavir/ritonavir ^{b,c}				
	Baseline	Week 48		Week 96		Baseline	Week 48		Week 96	
	mg/dL (n=428 ^e)	mg/dL (n=372 ^e)	Change ^d (n=372 ^e)	mg/dL (n=342 ^e)	Change ^d (n=342 ^e)	mg/dL (n=424 ^e)	mg/dL (n=335 ^e)	Change ^d (n=335 ^e)	mg/dL (n=291 ^e)	Change ^d (n=291 ^e)
LDL-	92	105	+14%	105	+14%	93	111	+19%	110	+17%
HDL-	37	46	+29%	44	+21%	36	48	+37%	46	+29%
Total	149	169	+13%	169	+13%	150	187	+25%	186	+25%
Triglycerides ^f	126	145	+15%	140	+13%	129	194	+52%	184	+50%

a. atazanavir 300 mg with ritonavir 100 mg once daily with the fixed-dose product: 300 mg tenofovir DF, 200 mg emtricitabine once daily.

b. Values obtained after initiation of serum lipid-reducing agents were not included in these analyses. At baseline, serum lipid-reducing agents were used in 1% in the lopinavir/ritonavir treatment arm and 1% in the atazanavir /ritonavir arm. Through Week 48, serum lipid-reducing agents were used in 8% in the lopinavir/ritonavir treatment arm and 2% in the atazanavir /ritonavir arm. Through Week 96, serum lipid- reducing agents were used in 10% in the lopinavir/ritonavir treatment arm and 3% in the atazanavir /ritonavir arm.

c. Lopinavir 400 mg with ritonavir 100 mg twice daily with the fixed-dose product 300 mg tenofovir DF, 200 mg emtricitabine once daily.

- d. The change from baseline is the mean of within-subject changes from baseline for subjects with both baseline and Week 48 or Week 96 values and is not a simple difference of the baseline and Week 48 or Week 96 mean values, respectively.
- e. Number of subjects with LDL-cholesterol measured.
- f. Fasting.

Table 13: Lipid Values, Mean Change from Baseline, Study AI424-034

	atazanavir ^{a,b}			efavirenz ^{b,c}		
	Baseline mg/dL (n=383 ^e)	Week 48 mg/dL (n=283 ^e)	Week 48 Changed (n=272 ^e)	Baseline mg/dL (n=378 ^e)	Week 48 mg/dL (n=264 ^e)	Week 48 Changed (n=253 ^e)
LDL-Cholesterol ^f	98	98	+1%	98	114	+18%
HDL-Cholesterol	39	43	+13%	38	46	+24%
Total Cholesterol	164	168	+2%	162	195	+21%
Triglycerides ^f	138	124	-9%	129	168	+23%

- a. Atazanavir 400 mg once daily with the fixed-dose product: 150 mg lamivudine, 300 mg zidovudine twice daily.
- b. Values obtained after initiation of serum lipid-reducing agents were not included in these analyses. At baseline, serum lipid-reducing agents were used in 0% in the efavirenz treatment arm and <1% in the atazanavir arm. Through Week 48, serum lipid-reducing agents were used in 3% in the efavirenz treatment arm and 1% in the atazanavir arm.
- c. Efavirenz 600 mg once daily with the fixed-dose product: 150 mg lamivudine, 300 mg zidovudine twice daily.
- d. The change from baseline is the mean of within-patient changes from baseline for patients with both baseline and Week 48 values and is not a simple difference of the baseline and Week 48 mean values.
- e. Number of subjects with LDL-cholesterol measured.
- f. Fasting.

Laboratory Abnormalities in Treatment-Experienced Subjects with HIV-1 Infection

The percentages of adult treatment-experienced subjects with HIV-1 infection treated with combination therapy including atazanavir /ritonavir having Grade 3-4 laboratory abnormalities are presented in Table 14.

Table 14: Grade 3-4 Laboratory Abnormalities Reported in ≥2% of Adult Treatment-Experienced Subjects with HIV-1 Infection, Study AI424-045^a

Variable	Limit ^c	48 weeks ^b atazanavir /ritonavir 300/100 mg (once daily) + tenofovir DF + NRTI (n=119)	48 weeks ^b lopinavir/ritonavir 400/100 mg (twice daily ^d) + tenofovir DF+ NRTI (n=118)
Chemistry	High		
SGOT/AST	≥5.1 x ULN	3%	3%
SGPT/ALT	≥5.1 x ULN	4%	3%

Total Bilirubin	≥2.6 x ULN	49%	<1%
Lipase	≥2.1 x ULN	5%	6%
Creatine Kinase	≥5.1 x ULN	8%	8%
Total Cholesterol	≥240 mg/dL	25%	26%
Triglycerides	≥751 mg/dL	8%	12%
Glucose	≥251 mg/dL	5%	<1%
Hematology	Low		
Platelets	<50,000 cells/mm ³	2%	3%
Neutrophils	<750 cells/mm ³	7%	8%

a Based on regimen(s) containing atazanavir.

b Median time on therapy.

c ULN = upper limit of normal.

d As a fixed-dose product.

Change in Lipids from Baseline in Treatment-Experienced Subjects with HIV-1 Infection

For Study AI424-045, changes from baseline in LDL-cholesterol, HDL-cholesterol, total cholesterol, and triglycerides are shown in Table 15. The observed magnitude of dyslipidemia was less with atazanavir /ritonavir than with lopinavir/ritonavir. However, the clinical impact of such findings has not been demonstrated.

Table 15: Lipid Values, Means Change from Baseline, Study AI424-045

	atazanavir /ritonavir ^{a,b}			lopinavir/ritonavir ^{b,c}		
	Baseline mg/dL (n=111 ^e)	Week 48 mg/dL (n=75 ^e)	Week 48 Change ^d (n=74 ^e)	Baseline mg/dL (n=108 ^e)	Week 48 mg/dL (n=76 ^e)	Week 48 Change ^d (n=73 ^e)
LDL-Cholesterol ^f	108	98	-10%	104	103	+1%
HDL-Cholesterol	40	39	-7%	39	41	+2%
Total Cholesterol	188	170	-8%	181	187	+6%
Triglycerides ^f	215	161	-4%	196	224	+30%

a. atazanavir 300 mg once daily + ritonavir + tenofovir DF + 1 NRTI.

b. Values obtained after initiation of serum lipid-reducing agents were not included in these analyses. At baseline, serum lipid-reducing agents were used in 4% in the lopinavir/ritonavir treatment arm and 4% in the atazanavir /ritonavir arm. Through Week 48, serum lipid-reducing agents were used in 19% in the lopinavir/ritonavir treatment arm and 8% in the atazanavir /ritonavir arm.

c. Lopinavir/ritonavir (400/100 mg) as a fixed dose regimen, BID + tenofovir DF + 1 NRTI.

d. The change from baseline is the mean of within-subjects changes from baseline for subjects with both baseline and Week 48 values and is not a simple difference of the baseline and Week 48 mean values.

e. Number of subjects with LDL-cholesterol measured.

f. Fasting.

Adverse Reactions in Pediatric Subjects with HIV-1 Infection: atazanavir Capsules

The safety and tolerability of atazanavir Capsules with and without ritonavir have been established in pediatric subjects with HIV-1 infection, at least 6 years of age from the open-label, multicenter clinical trial PACTG 1020A.

The safety profile of atazanavir in pediatric subjects with HIV-1 infection (6 to less than 18 years of age) taking the capsule formulation was generally similar to that observed in clinical studies of atazanavir in adults. The most common Grade 2-4 adverse events ($\geq 5\%$, regardless of causality) reported in pediatric subjects were cough (21%), fever (18%), jaundice/scleral icterus (15%), rash (14%), vomiting (12%), diarrhea (9%), headache (8%), peripheral edema (7%), extremity pain (6%), nasal congestion (6%), oropharyngeal pain (6%), wheezing (6%), and rhinorrhea (6%). Asymptomatic second-degree atrioventricular block was reported in $<2\%$ of subjects. The most common Grade 3-4 laboratory abnormalities occurring in pediatric subjects taking the capsule formulation were elevation of total bilirubin (≥ 3.2 mg/dL, 58%), neutropenia (9%), and hypoglycemia (4%). All other Grade 3-4 laboratory abnormalities occurred with a frequency of less than 3%.

Adverse Reactions in subjects with HIV-1 Infection Co-Infected with Hepatitis B and/or Hepatitis C Virus

In Study AI424-138, 60 subjects administered Atazanavir /ritonavir 300 mg/100 mg once daily, and 51 subjects treated with lopinavir/ritonavir 400 mg/100 mg (as fixed-dose product) twice daily, each with fixed dose tenofovir DF-emtricitabine, were seropositive for hepatitis B and/or C at study entry. ALT levels >5 times ULN developed in 10% (6/60) of the subjects administered atazanavir /ritonavir and 8% (4/50) of the subjects treated with lopinavir/ritonavir. AST levels >5 times ULN developed in 10% (6/60) of the subjects administered atazanavir /ritonavir and none (0/50) of the subjects treated with lopinavir/ritonavir.

In Study AI424-045, 20 subjects administered atazanavir /ritonavir 300 mg/100 mg once daily, and 18 subjects treated with lopinavir/ritonavir 400 mg/100 mg twice daily (as fixed-dose product), were seropositive for hepatitis B and/or C at study entry. ALT levels >5 times ULN developed in 25% (5/20) of the subjects administered atazanavir /ritonavir and 6% (1/18) of the subjects treated with lopinavir/ritonavir-treated. AST levels >5 times ULN developed in 10% (2/20) of the subjects administered atazanavir /ritonavir and 6% (1/18) of the subjects treated with lopinavir/ritonavir.

In Studies AI424-008 and AI424-034, 74 subjects treated with Atazanavir 400 mg once daily, 58 who received efavirenz, and 12 who received nelfinavir were seropositive for hepatitis B and/or C at study entry. ALT levels >5 times ULN developed in 15% of the subjects treated with atazanavir -, 14% of the subjects treated with efavirenz-, and 17% of the subjects treated with nelfinavir. AST levels >5 times ULN developed in 9% of the subjects treated with atazanavir , 5% of the subjects treated with efavirenz, and

Atazanavir Teva 150/200/300mg KL SPC Notification 12-2023

17% of the subjects treated with nelfinavir. Within atazanavir and control regimens, no difference in frequency of bilirubin elevations was noted between seropositive and seronegative subjects [see Warnings and Precautions (5.7)].

6.2. Postmarketing Experience

The following events have been identified during postmarketing use of Atazanavir. Because these reactions are reported voluntarily from a population of unknown size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Body as a Whole: edema

Cardiovascular System: second-degree AV block, third-degree AV block, left bundle branch block, QTc prolongation [see Warnings and Precautions (5.1)]

Gastrointestinal System: pancreatitis

Hepatic System: hepatic function abnormalities

Hepatobiliary Disorders: cholelithiasis [see Warnings and Precautions (5.6)], cholecystitis, cholestasis

Metabolic System and Nutrition Disorders: diabetes mellitus, hyperglycemia [see Warnings and Precautions (5.9)]

Musculoskeletal System: arthralgia

Renal System: nephrolithiasis [see Warnings and Precautions (5.6)], interstitial nephritis, granulomatous interstitial nephritis, chronic kidney disease [see Warnings and Precautions (5.4)]

Skin and Appendages: alopecia, maculopapular rash [see *Contraindications (4) and Warnings and Precautions (5.2)*], pruritus, angioedema

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product.

Any suspected adverse events should be reported to the Ministry of Health according to the National Regulation by using an online form <https://sideeffects.health.gov.il>

7. DRUG INTERACTIONS

7.1. Potential for Atazanavir Teva to Affect Other Drugs

Atazanavir is an inhibitor of CYP3A and UGT1A1. Coadministration of Atazanavir Teva and drugs primarily metabolized by CYP3A or UGT1A1 may result in increased plasma concentrations of the other drug that could increase or prolong its therapeutic and adverse effects.

Atazanavir is a weak inhibitor of CYP2C8. Use of Atazanavir Teva without ritonavir is not recommended

when coadministered with drugs highly dependent on CYP2C8 with narrow therapeutic indices (eg, paclitaxel, repaglinide). When Atazanavir Teva with ritonavir is coadministered with substrates of CYP2C8, clinically significant interactions are not expected [see Clinical Pharmacology, Table 21 (11.3)]. The magnitude of CYP3A-mediated drug interactions on coadministered drug may change when Atazanavir Teva is coadministered with ritonavir. See the complete prescribing information for ritonavir for information on drug interactions with ritonavir.

7.2. Potential for Other Drugs to Affect Atazanavir Teva

Atazanavir is a CYP3A4 substrate; therefore, drugs that induce CYP3A4 may decrease atazanavir plasma concentrations and reduce Atazanavir Teva's therapeutic effect.

Atazanavir solubility decreases as pH increases. Reduced plasma concentrations of atazanavir are expected if proton-pump inhibitors, antacids, buffered medications, or H2-receptor antagonists are administered with Atazanavir Teva [see Dosage and Administration (2.3, 2.4, 2.5 and 2.6)] .

7.3. Established and Other Potentially Significant Drug Interactions

Table 16 provides dosing recommendations in adults as a result of drug interactions with Atazanavir Teva. These recommendations are based on either drug interaction studies or predicted interactions due to the expected magnitude of interaction and potential for serious events or loss of efficacy.

Table 16: Established and Other Potentially Significant Drug Interactions: Alteration in Dose or Regimen May Be Recommended Based on Drug Interaction Studies or Predicted Interactions (Information in the table applies to Atazanavir Teva with or without ritonavir, unless otherwise indicated)

Concomitant Drug Class: Specific Drugs	Effect on Concentration of Atazanavir or Concomitant Drug	Clinical Comment
<i>HIV Antiviral Agents</i>		
<i>Nucleoside Reverse Transcriptase Inhibitors (NRTIs):</i> didanosine buffered formulations enteric-coated (EC) capsules	↓ atazanavir ↓ didanosine	It is recommended that Atazanavir Teva be given (with food) 2 h before or 1 h after didanosine buffered formulations. Simultaneous administration of didanosine EC and Atazanavir Teva with food results in a decrease in didanosine. Thus, Atazanavir Teva and didanosine EC should be administered at different times.
<i>Nucleotide Reverse Transcriptase Inhibitors:</i> tenofovir disoproxil fumarate (DF)	↓ atazanavir ↑ tenofovir	When coadministered with tenofovir DF in adults, it is recommended that Atazanavir Teva 300 mg be given with ritonavir 100 mg and tenofovir DF 300 mg (all as a single daily dose with food)The mechanism of this interaction is

		unknown. Higher tenofovir concentrations could potentiate tenofovir -associated adverse reactions, including renal disorders. Patients receiving Atazanavir Teva and tenofovir DF should be monitored for tenofovir-associated adverse reactions. For pregnant women taking AtazanavirTeva with ritonavir and tenofovir DF, see <i>Dosage and Administration (2.6)</i> .
<i>Non-nucleoside Reverse Transcriptase Inhibitors (NNRTIs):</i> efavirenz	↓atazanavir	<i>In HIV-treatment-naive adult patients:</i> If Atazanavir Teva is combined with efavirenz, Atazanavir Teva 400 mg (two 200-mg capsules) should be administered with ritonavir 100 mg simultaneously once daily with food, and efavirenz 600 mg should be administered once daily on an empty stomach, preferably at bedtime. <i>In HIV treatment-experienced adult patients:</i> Coadministration of Atazanavir Teva with efavirenz is not recommended.
nevirapine	↓atazanavir ↑ nevirapine	Coadministration of Atazanavir Teva with nevirapine is contraindicated due to the potential loss of virologic response and development of resistance, as well as potential risk for nevirapine-associated adverse reactions [see <i>Contraindications (4)</i>].
<i>Protease Inhibitors:</i> saquinavir (soft gelatin capsules)	↑saquinavir	Appropriate dosing recommendations for this combination, with or without ritonavir, with respect to efficacy and safety have not been established. In a clinical study, saquinavir 1200 mg coadministered with atazanavir 400 mg and tenofovir DF 300 mg (all given once daily) plus nucleoside analogue reverse transcriptase inhibitors did not provide adequate efficacy [see <i>Clinical Studies (13.2)</i>].
indinavir		Coadministration of Atazanavir Teva with indinavir is contraindicated. Both Atazanavir Teva and indinavir are associated with indirect (unconjugated) hyperbilirubinemia [see <i>Contraindications (4)</i>].
Ritonavir	↑ atazanavir	If Atazanavir Teva is coadministered with ritonavir, it is recommended that Atazanavir Teva 300 mg once daily be given with ritonavir 100 mg once daily with food in adults. See the complete prescribing information for ritonavir for information on drug interactions with ritonavir.
Others	↑other protease inhibitor	Coadministration with other protease inhibitors is not recommended.

Hepatitis C Antiviral Agents

<i>Concomitant Drug Class: Specific Drugs</i>	<i>Effect on Concentration of Atazanavir or Concomitant Drug</i>	<i>Clinical Comment</i>
elbasvir/grazoprevir	↑grazoprevir	Coadministration of Atazanavir Teva with grazoprevir is

		contraindicated due to the potential increased risk of ALT elevations [see <i>Contraindications (4)</i>].
glecaprevir/pibrentasvir	↑glecaprevir ↑ pibrentasvir	Coadministration of Atazanavir Teva with glecaprevir/pibrentasvir is contraindicated due to the potential for increased the risk of ALT elevations [see <i>Contraindications (4)</i>].
voxilaprevir/sofosbuvir/velpatasvir	↑voxilaprevir	Coadministration with Atazanavir Teva is not recommended

Other Agents

Concomitant Drug Class: Specific Drugs	Effect on Concentration of Atazanavir or Concomitant Drug	Clinical Comment
Alpha 1-Adrenoreceptor Antagonist: alfuzosin	↑alfuzosin	Coadministration of Atazanavir Teva with alfuzosin is contraindicated due to risk for hypotension [see <i>Contraindications (4)</i>].
<i>Antacids and buffered medications</i>	↓ atazanavir	Atazanavir should be administered 2 hours before or 1 hour after antacids and buffered medications.
Antiarrhythmics: amiodarone, quinidine amiodarone, bepridil, lidocaine (systemic), quinidine	↑amiodarone, bepridil, lidocaine (systemic), quinidine	Concomitant use of Atazanavir Teva with ritonavir and either quinidine or amiodarone is contraindicated due to the potential for serious or life-threatening reactions such as cardiac arrhythmias [see <i>Contraindications (4)</i>]. Coadministration with Atazanavir Teva without ritonavir has the potential to produce serious and/or life-threatening adverse events but has not been studied. Caution is warranted and therapeutic concentration monitoring of these drugs is recommended if they are used concomitantly with Atazanavir without ritonavir.
Anticoagulants: warfarin	↑warfarin	Coadministration with Atazanavir Teva has the potential to produce serious and/or life-threatening bleeding and has not been studied. It is recommended that International Normalized Ratio (INR) be monitored.
Direct-Acting Oral Anticoagulants: betrixaban, dabigatran, edoxaban	↑ betrixaban ↑dabigatran ↑edoxaban	Concomitant use of Atazanavir Teva with ritonavir, a strong CYP3A4/P-gp inhibitor, may result in increased risk of bleeding. Refer to the respective DOAC prescribing information regarding dosing instructions for coadministration with P-gp inhibitors.
rivaroxaban	Atazanavir <i>with ritonavir</i> ↑ rivaroxaban Atazanavir ↑ rivaroxaban	Coadministration of Atazanavir Teva with ritonavir, a strong CYP3A4/P-gp inhibitor, and rivaroxaban is not recommended, as it may increase risk of bleeding. Coadministration of Atazanavir Teva, a CYP3A4 inhibitor, and rivaroxaban may result in increased increase. Close monitoring is recommended when Atazanavir Teva is coadministered with rivaroxaban.

Concomitant Drug Class: Specific Drugs	Effect on Concentration of Atazanavir or Concomitant Drug	Clinical Comment
apixaban	Atazanavir <i>with ritonavir</i> ↑ apixaban Atazanavir ↑ apixaban	Concomitant use of Atazanavir Teva with ritonavir, a strong CYP3A4/P-gp inhibitor, may result in increased risk of bleeding. Refer to apixaban dosing instructions for coadministration with strong CYP3A4 and P-gp inhibitors in the apixaban prescribing information. Concomitant use of Atazanavir, a CYP3A4 inhibitor, and apixaban may result in increased risk of bleeding. Close monitoring is recommended when apixaban is coadministered with Atazanavir Teva.
Antidepressants: tricyclic antidepressants	↑tricyclic antidepressants	Coadministration with Atazanavir Teva has the potential to produce serious and/or life-threatening adverse events and has not been studied. Concentration monitoring of these drugs is recommended if they are used concomitantly with Atazanavir Teva.
trazodone	↑trazodone	Nausea, dizziness, hypotension, and syncope have been observed following coadministration of trazodone with ritonavir. If trazodone is used with a CYP3A4 inhibitor such as Atazanavir Teva, the combination should be used with caution and a lower dose of trazodone should be considered.
Antiepileptics: carbamazepine	↓Atazanavir ↑carbamazepine	Coadministered with Atazanavir Teva (without or without ritonavir) with carbamazepine is contraindicated due to the risk for loss of virologic response and development of resistance [see Contraindications (4)].
phenytoin, phenobarbital	↓atazanavir ↓phenytoin ↓phenobarbital	Coadministration Atazanavir Teva (without or without ritonavir) with phenytoin or phenobarbital is contraindicated due to the risk for loss of virologic response and development of resistance [see Contraindications (4)].
Lamotrigine	↓lamotrigine	Coadministration of lamotrigine and Atazanavir Teva with ritonavir may require dosage adjustment of lamotrigine. No dose adjustment of lamotrigine is required when coadministered with Atazanavir Teva without ritonavir.
Antifungals: ketoconazole, itraconazole	Atazanavir <i>with ritonavir</i> : ↑ketoconazole ↑itraconazole	Coadministration of ketoconazole has only been studied with Atazanavir Teva without ritonavir (negligible increase in atazanavir AUC and C _{max}). Due to the effect of ritonavir on ketoconazole, high doses of ketoconazole and itraconazole (>200 mg/day) should be used cautiously when administering Atazanavir Teva with ritonavir.
Voriconazole	Atazanavir Teva <i>with ritonavir in</i>	The use of voriconazole in patients receiving Atazanavir Teva/ritonavir is not recommended unless an assessment

Concomitant Drug Class: Specific Drugs	Effect on Concentration of Atazanavir or Concomitant Drug	Clinical Comment
	<p><i>subjects with a functional CYP2C19 allele:</i> ↓ voriconazole ↓ atazanavir</p> <p><i>Atazanavir Teva with ritonavir in subjects without a functional CYP2C19 allele:</i> ↑ voriconazole ↓ atazanavir</p>	of the benefit/risk to the patient justifies the use of voriconazole. Patients should be carefully monitored for voriconazole- associated adverse reactions and loss of either voriconazole or atazanavir efficacy during the coadministration of voriconazole and Atazanavir Teva /ritonavir. Coadministration of voriconazole with Atazanavir Teva (without ritonavir) may affect atazanavir concentrations; however, no data are available.
Antigout: colchicine	↑ colchicine	<p>The coadministration of Atazanavir Teva with colchicine in patients with renal or hepatic impairment is not recommended.</p> <p>Recommended adult dosage of colchicine when administered with Atazanavir Teva: Treatment of gout flares: 0.6 mg (1 tablet) for 1 dose, followed by 0.3 mg (half tablet) 1 hour later. Not to be repeated before 3 days. Prophylaxis of gout flares: If the original regimen was 0.6 mg twice a day, the regimen should be adjusted to 0.3 mg once a day. If the original regimen was 0.6 mg once a day, the regimen should be adjusted to 0.3 mg once every other day. Treatment of familial Mediterranean fever (FMF): Maximum daily dose of 0.6 mg (may be given as 0.3 mg twice a day).</p>
Antimycobacterials: rifampin	↓ atazanavir	Coadministration of Atazanavir Teva with rifampin is contraindicated due to risk for loss of virologic response and development of resistance [<i>see Contraindications (4)</i>].
rifabutin	↑ rifabutin	A rifabutin dose reduction of up to 75% (eg, 150 mg every other day or 3 times per week) is recommended. Increased monitoring for rifabutin-associated adverse reactions including neutropenia is warranted.
Antineoplastics: irinotecan	↑ irinotecan	Coadministration of Atazanavir Teva with irinotecan is contraindicated. Atazanavir inhibits UGT1A1 and may interfere with the metabolism of irinotecan, resulting in increased irinotecan toxicities [<i>see Contraindications (4)</i>].
apalutamide	↓ atazanavir	Coadministration of Atazanavir Teva (with or without ritonavir) and apalutamide is contraindicated due to the

Concomitant Drug Class: Specific Drugs	Effect on Concentration of Atazanavir or Concomitant Drug	Clinical Comment
ivosidenib	↓ atazanavir ↑ ivosidenib	potential for subsequent loss of virologic response and possible resistance to the class of protease inhibitors [<i>see Contraindications (4)</i>]. Coadministration of ivosidenib with Atazanavir Teva (with or without ritonavir) is contraindicated due to the potential for loss of virologic response and risk of serious adverse events such as QT interval prolongation.
encorafenib	↓ atazanavir ↑ encorafenib	Coadministration of encorafenib with Atazanavir Teva (with or without ritonavir) is contraindicated due to the potential for the loss of virologic response and risk of serious adverse events such as QT interval prolongation.
<i>Antiplatelets</i>		
ticagrelor	↑ ticagrelor	Coadministration with ticagrelor is not recommended due to potential increase in the risk of dyspnea, bleeding and other adverse events associated with ticagrelor.
clopidogrel	↓ clopidogrel active metabolite	Coadministration of Atazanavir Teva (with or without ritonavir) and clopidogrel is not recommended. This is due to the potential reduction of the antiplatelet activity of clopidogrel.
<i>Antipsychotics:</i> pimozide	↑ pimozide	Coadministration of Atazanavir Teva with pimozide is contraindicated. This is due to the potential for serious and/or life-threatening reactions such as cardiac arrhythmias [<i>see Contraindications (4)</i>]
lurasidone	<i>Atazanavir Teva with ritonavir</i> ↑ lurasidone	<i>Atazanavir Teva with ritonavir</i> Coadministration of lurasidone with <i>Atazanavir Teva</i> with ritonavir is contraindicated. This is due to the potential for serious and/or life-threatening reactions [<i>see Contraindications (4)</i>].
	<i>Atazanavir Teva</i> ↑ lurasidone	<i>Atazanavir Teva without ritonavir</i> If coadministration is necessary, reduce the lurasidone dose. Refer to the lurasidone prescribing information for concomitant use with moderate CYP3A4 inhibitors.
quetiapine	↑ quetiapine	<i>Initiation of Atazanavir Teva with ritonavir in patients taking quetiapine:</i> Consider alternative antiretroviral therapy to avoid increases in quetiapine exposures. If coadministration is necessary, reduce the quetiapine dose to 1/6 of the current dose and monitor for quetiapine-associated adverse reactions. Refer to the quetiapine prescribing information for recommendations on adverse reaction monitoring.

Concomitant Drug Class: Specific Drugs	Effect on Concentration of Atazanavir or Concomitant Drug	Clinical Comment
		Initiation of quetiapine in patients taking Atazanavir Teva with ritonavir: Refer to the quetiapine prescribing information for initial dosing and titration of quetiapine.
Benzodiazepines: Midazolam (oral) triazolam	↑ midazolam ↑ triazolam	Coadministration of Atazanavir Teva with either orally administered midazolam or triazolam is contraindicated. Triazolam and orally administered midazolam are extensively metabolized by CYP3A4, and coadministration with Atazanavir Teva can lead to the potential for serious and/or life-threatening events such as prolonged or increased sedation or respiratory depression [<i>see Contraindications (4)</i>].
parenterally administered midazolam ^b	↑ midazolam	. Coadministration parenteral with midazolam should be done in a setting which ensures close clinical monitoring and appropriate medical management in case of respiratory depression and/or prolonged sedation. Dosage reduction for midazolam should be considered, especially if more than a single dose of midazolam is administered.
<i>Calcium channel blockers:</i> diltiazem	↑ diltiazem and desacetyl-diltiazem	Caution is warranted. A dose reduction of diltiazem by 50% should be considered. ECG monitoring is recommended. Coadministration of diltiazem and atazanavir with ritonavir has not been studied.
felodipine, nifedipine, nicardipine, and verapamil	↑ calcium channel blocker	Caution is warranted. Dose titration of the calcium channel blocker should be considered. ECG monitoring is recommended.
<i>Corticosteroids:</i> dexamethasone and other corticosteroids (all routes of administration)	↓ atazanavir ↑ corticosteroids	Coadministration with dexamethasone or other corticosteroids that induce CYP3A may result in loss of therapeutic effect of REYATAZ and development of resistance to atazanavir and/or ritonavir. Alternative corticosteroids should be considered. Coadministration with corticosteroids (all routes of administration) that are metabolized by CYP3A, particularly for long-term use, may increase the risk for development of systemic corticosteroid effects including Cushing's syndrome and adrenal suppression. Consider the potential benefit of treatment versus the risk of systemic corticosteroid effects. For coadministration of cutaneously administered corticosteroids sensitive to CYP3A inhibition, refer to the prescribing information of the corticosteroid for additional information

Concomitant Drug Class: Specific Drugs	Effect on Concentration of Atazanavir or Concomitant Drug	Clinical Comment
Endothelin receptor antagonists: Bosentan	Atazanavir ↓ atazanavir Atazanavir with ritonavir ↑ bosentan	Coadministration of bosentan and Atazanavir Teva without ritonavir is not recommended. For adult patients who have been receiving Atazanavir Teva with ritonavir for at least 10 days, start bosentan at 62.5 mg once daily or every other day based on individual tolerability. For adult patients who have been receiving bosentan, discontinue bosentan at least 36 hours before starting Atazanavir Teva with ritonavir. At least 10 days after starting Atazanavir Teva with ritonavir, resume bosentan at 62.5 mg once daily or every other day based on individual tolerability.
Ergot derivatives: dihydroergotamine, ergotamine, ergonovine, methylergonovine	↑ ergot derivatives	Coadministration of Atazanavir Teva with ergot derivatives is contraindicated. This is due to the potential for serious and/or life-threatening events such as acute ergot toxicity characterized by peripheral vasospasm and ischemia of the extremities and other tissues [<i>see Contraindications (4)</i>].
GI Motility Agents: cisapride	↑ cisapride	Coadministration of Atazanavir Teva with cisapride is contraindicated. This is due to the potential for serious and/or life-threatening reactions such as cardiac arrhythmias [<i>see Contraindications (4)</i>].
Gonadotropin-releasing hormone Receptor (GnRH) Antagonists: elagolix	↓ atazanavir ↑ elagolix	Coadministration of elagolix and REYATAZ with or without ritonavir is not recommended due to the potential of loss of virologic response and the potential risk of adverse events such as bone loss and hepatic transaminase elevations associated with elagolix. In the event coadministration is necessary, limit concomitant use of elagolix 200mg twice daily with REYATAZ with or without ritonavir for up to 1 month or limit concomitant use of elagolix 150 mg once daily with REYATAZ (with or without ritonavir) for up to 6 months and monitor virologic response.
Herbal Products: St. John's wort (Hypericum perforatum)	↓ atazanavir	Coadministration of products containing St. John's wort with Atazanavir Teva is contraindicated. This may result in loss of therapeutic effect of Atazanavir Teva and the development of resistance [<i>see Contraindications (4)</i>].
Kinase inhibitors: fostamatinib	↑ R406 (active metabolite of fostamatinib)	When coadministering fostamatinib with REYATAZ (with or without ritonavir), monitor for toxicities of R406 exposure resulting in dose-related adverse events such as hepatotoxicity and neutropenia. Fostamatinib dose reduction may be required.

Concomitant Drug Class: Specific Drugs	Effect on Concentration of Atazanavir or Concomitant Drug	Clinical Comment
Lipid-modifying agents HMG-CoA reductase inhibitors: lovastatin, simvastatin	↑ Lovastatin ↑ simvastatin	Coadministration of Atazanavir Teva with lovastatin or simvastatin is contraindicated. This is due to the potential for serious reactions such as myopathy, including rhabdomyolysis [<i>see Contraindications (4)</i>].
atorvastatin, rosuvastatin	↑ atorvastatin ↑ rosuvastatin	Titrate atorvastatin dose carefully and use the lowest necessary dose. Rosuvastatin dose should not exceed 10 mg/day. The risk of myopathy, including rhabdomyolysis, may be increased when HIV protease inhibitors, including Atazanavir Teva, are used in combination with these drugs.
Other Lipid Modifying Agents: lomitapide	↑ lomitapide	Coadministration of Atazanavir Teva with lomitapide is contraindicated. This is due to the potential for risk of markedly increased transaminase levels and hepatotoxicity associated with increased plasma concentrations of lomitapide. The mechanism of interaction is CYP3A4 inhibition by atazanavir and/or ritonavir [<i>see Contraindications (4)</i>].
H2-Receptor antagonists	↓ atazanavir	Coadministration may result in loss of virologic response and development of resistance. <i>In HIV treatment-naive adult patients:</i> Atazanavir Teva 300 mg with ritonavir 100 mg once daily with food should be administered simultaneously with, and/or at least 10 hours after, a dose of the H2-receptor antagonist (H2RA). An H2RA dose comparable to famotidine 20 mg once daily up to a dose comparable to famotidine 40 mg twice daily can be used with Atazanavir Teva 300 mg with ritonavir 100 mg in treatment-naive patients. OR For patients unable to tolerate ritonavir, Atazanavir Teva 400 mg once daily with food should be administered at least 2 hours before and at least 10 hours after a dose of the H2RA. No single dose of the H2RA should exceed a dose comparable to famotidine 20 mg, and the total daily dose should not exceed a dose comparable to famotidine 40 mg. The use of Atazanavir Teva without ritonavir in pregnant women is not recommended. <i>In treatment-experienced adult patients:</i> Whenever an H2RA is given to a patient receiving Atazanavir Teva with ritonavir, the H2RA dose should not exceed a dose comparable to famotidine 20 mg twice daily, and the Atazanavir Teva and ritonavir doses should be administered simultaneously with, and/or at least 10 hours after, the dose of the H2RA.

Concomitant Drug Class: Specific Drugs	Effect on Concentration of Atazanavir or Concomitant Drug	Clinical Comment
		<ul style="list-style-type: none"> • Atazanavir Teva 300 mg with ritonavir 100 mg once daily (all as a single dose with food) if taken with an H2RA. • Atazanavir Teva 400 mg with ritonavir 100 mg once daily (all as a single dose with food) if taken with both tenofovir DF and an H2RA. • Atazanavir Teva 400 mg with ritonavir 100 mg once daily (all as a single dose with food) if taken with both tenofovir DF and an H2RA. • Atazanavir Teva 400 mg with ritonavir 100 mg once daily (all as a single dose with food) if taken with either tenofovir DF or an H2RA for pregnant patient during the second and third trimester. Atazanavir Teva is not recommended for pregnant patients during the second and third trimester taking Atazanavir Teva with both tenofovir DF and an H2RA.
Hormonal contraceptives: ethinyl estradiol and norgestimate or norethindrone	<p>↓ethinyl estradiol ↑norgestimate^c</p> <p>↑ ethinyl estradiol ↑norethindrone^d</p>	<p>Use caution if considering coadministration of oral contraceptives with Atazanavir Teva or Atazanavir Teva with ritonavir.</p> <p>If Atazanavir with ritonavir is coadministered with an oral contraceptive, it is recommended that the oral contraceptive contain at least 35 mcg of ethinyl estradiol. If Atazanavir Teva is administered without ritonavir, the oral contraceptive should contain no more than 30 mcg of ethinyl estradiol.</p> <p>Potential safety risks include substantial increases in progesterone exposure. The long-term effects of increases in concentration of the progestational agent are unknown and could increase the risk of insulin resistance, dyslipidemia, and acne.</p> <p>Coadministration of atazanavir or atazanavir with ritonavir and other hormonal contraceptives (eg, contraceptive patch, contraceptive vaginal ring, or injectable contraceptives) or oral contraceptives containing progestogens other than norethindrone or norgestimate, or less than 25 mcg of ethinyl estradiol, has not been studied; therefore, alternative methods of contraception are recommended.</p>
<i>Immunosuppressants:</i> cyclosporine, sirolimus, tacrolimus	↑ immunosuppressants	Therapeutic concentration monitoring is recommended for these immunosuppressants when coadministered with Atazanavir Teva.
<i>Inhaled beta agonist:</i> Salmeterol	↑ salmeterol	Coadministration of salmeterol with Atazanavir Teva is not recommended. Concomitant use of salmeterol and Atazanavir Teva may result in increased risk of cardiovascular adverse reactions

Concomitant Drug Class: Specific Drugs	Effect on Concentration of Atazanavir or Concomitant Drug	Clinical Comment
		associated with salmeterol, including QT prolongation, palpitations, and sinus tachycardia.
<i>Inhaled nasal steroid:</i> fluticasone	<i>Atazanavir</i> ↑ fluticasone <i>Atazanavir with ritonavir</i> ↑fluticasone	Concomitant use of fluticasone propionate and Atazanavir Teva without ritonavir should be used with caution. Consider alternatives to fluticasone propionate, particularly for long term use. With concomitant use of fluticasone propionate and Atazanavir Teva with ritonavir systemic corticosteroid effects, including Cushing's syndrome and adrenal suppression, have been reported during postmarketing use in patients receiving ritonavir and inhaled or intranasally administered fluticasone propionate. Coadministration of fluticasone propionate and Atazanavir Teva with ritonavir is not recommended unless the potential benefit to the patient outweighs the risk of systemic corticosteroid side effects [<i>see Warnings and Precautions (5.1)</i>].
<i>Macrolide antibiotics:</i> clarithromycin	↑ clarithromycin ↓ 14-OH clarithromycin ↑ atazanavir	Increased concentrations of clarithromycin may cause QTc prolongations; therefore, a dose reduction of clarithromycin by 50% should be considered when it is coadministered with Atazanavir Teva. In addition, concentrations of the active metabolite 14-OH clarithromycin are significantly reduced; consider alternative therapy for indications other than infections due to Mycobacterium avium complex. Coadministration of Atazanavir /ritonavir with clarithromycin has not been studied.
<i>Opioids:</i> Buprenorphine	Atazanavir Teva or Atazanavir with ritonavir ↑ buprenorphine ↑ norbuprenorphine Atazanavir ↓ atazanavir	Coadministration of Atazanavir Teva plus ritonavir with buprenorphine warrants clinical monitoring for sedation and cognitive effects. A dose reduction of buprenorphine may be considered. Coadministration of buprenorphine and Atazanavir Teva with ritonavir is not expected to decrease atazanavir plasma concentrations. The coadministration of Atazanavir Teva and buprenorphine without ritonavir is not recommended.
<i>PDE5 inhibitors:</i> sildenafil, tadalafil, vardenafil	↑ sildenafil ↑ tadalafil ↑ vardenafil	Coadministration with atazanavir has not been studied but may result in an increase in PDE5 inhibitor-associated adverse reactions, including hypotension, syncope, visual disturbances, and priapism. <i>Use of PDE5 inhibitors for pulmonary arterial hypertension (PAH):</i> contraindicated of Atazanavir Teva with REVATIO® (sildenafil) for the treatment of pulmonary hypertension (PAH) is contraindicated [<i>see Contraindications (4)</i>].

Concomitant Drug Class: Specific Drugs	Effect on Concentration of Atazanavir or Concomitant Drug	Clinical Comment
		<p>The following dose adjustments are recommended for the use of ADCIRCA® (tadalafil) with Atazanavir Teva:</p> <p>Coadministration of ADCIRCA® in patients on Atazanavir (with or without ritonavir):</p> <ul style="list-style-type: none"> For patients receiving Atazanavir Teva (with or without ritonavir) for at least one week, start ADCIRCA® * at 20 mg once daily. Increase to 40 mg once daily based on individual tolerability. <p>Coadministration of Atazanavir Teva (with or without ritonavir) in patients on ADCIRCA®:</p> <ul style="list-style-type: none"> Avoid the use of ADCIRCA® when starting Atazanavir Teva (with or without ritonavir). Stop ADCIRCA® at least 24 hours before starting Atazanavir Teva (with or without ritonavir). At least one week after starting Atazanavir Teva (with or without ritonavir), resume ADCIRCA® at 20 mg once daily. Increase to 40 mg once daily based on individual tolerability. <p>Use of PDE5 inhibitors for erectile dysfunction: Use VIAGRA® (sildenafil) with caution at reduced doses of 25 mg every 48 hours with increased monitoring for adverse events. Use CIALIS® (tadalafil) with caution at reduced doses of 10 mg every 72 hours with increased monitoring for adverse events.</p> <p>Atazanavir Teva /ritonavir: Use vardenafil with caution at reduced doses of no more than 2.5 mg every 72 hours with increased monitoring for adverse reactions.</p> <p>Atazanavir Teva: Use vardenafil with caution at reduced doses of no more than 2.5 mg every 24 hours with increased monitoring for adverse reactions.</p>
Proton-pump inhibitors: omeprazole	↓ atazanavir	<p>Coadministration Atazanavir Teva with or without ritonavir and omeprazole may result in loss of virologic response and development of resistance.</p> <p>In HIV- treatment-naïve adult patients: The proton-pump inhibitor (PPI) dose should not exceed a dose comparable to omeprazole 20 mg and must be taken approximately 12 hours prior to the Atazanavir Teva 300 mg with ritonavir 100 mg dose.</p>

Concomitant Drug Class: Specific Drugs	Effect on Concentration of Atazanavir or Concomitant Drug	Clinical Comment
		<i>In HIV-treatment-experienced adult patients:</i> Coadministration Atazanavir Teva PPIs is not recommended.

^a For magnitude of interactions see Clinical Pharmacology, Tables 20 and 21 (11.3).

^b See Contraindications (4), Table 6 for orally administered midazolam.

^c In combination with atazanavir 300 mg with ritonavir 100 mg once daily.

^d In combination with atazanavir 400 mg once daily

7.4. Drugs with No Observed Interactions with Atazanavir Teva

No clinically significant drug interactions were observed when Atazanavir Teva was coadministered with methadone, fluconazole, acetaminophen, atenolol, or the nucleoside reverse transcriptase inhibitors lamivudine or zidovudine [see Clinical Pharmacology, Tables 20 and 21 (11.3)].

8. USE IN SPECIFIC POPULATIONS

8.1. Pregnancy

Pregnancy Exposure Registry

There is a pregnancy exposure registry that monitors pregnancy outcomes in women exposed to Atazanavir Teva during pregnancy.

Risk Summary

Atazanavir has been evaluated in a limited number of women during pregnancy. Available human and animal data suggest that atazanavir does not increase the risk of major birth defects overall compared to the background rate [see Data]. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2-4% and 15-20%, respectively. No treatment-related malformations were observed in rats and rabbits, for which the atazanavir exposures were 0.7-1.2 times of those at the human clinical dose (300 mg/day atazanavir boosted with 100 mg/day ritonavir). When atazanavir was administered to rats during pregnancy and throughout lactation, reversible neonatal growth retardation was observed [see Data].

Clinical Considerations

Dose Adjustments during Pregnancy and the Postpartum Period

- Atazanavir Teva must be administered with ritonavir in pregnant women.
- For pregnant patients, no dosage adjustment is required for Atazanavir Teva with the following exceptions:

- For treatment-experienced pregnant women during the second or third trimester, when Atazanavir Teva is coadministered with either an H2-receptor antagonist or tenofovir DF, Atazanavir Teva 400 mg with ritonavir 100 mg once daily is recommended. There are insufficient data to recommend a Atazanavir Teva dose for use with both an H2-receptor antagonist and tenofovir DF in treatment-experienced pregnant women.
- No dosage adjustment is required for postpartum patients. However, patients should be closely monitored for adverse events because atazanavir exposures could be higher during the first 2 months after delivery [see Dosage and Administration (2.6) and Clinical Pharmacology (11.3)].

Maternal Adverse Reactions

Cases of lactic acidosis syndrome, sometimes fatal, and symptomatic hyperlactatemia have occurred in pregnant women using Atazanavir Teva in combination with nucleoside analogues, which are associated with an increased risk of lactic acidosis syndrome.

Hyperbilirubinemia occurs frequently in patients who take Atazanavir Teva [see Warnings and Precautions (5.7)], including those who are pregnant [*see Data*].

Advise pregnant women of the potential risks of lactic acidosis syndrome and hyperbilirubinemia.

Fetal/Neonatal Adverse Reactions

All infants, including neonates exposed to Atazanavir Teva in utero, should be monitored for the development of severe hyperbilirubinemia during the first few days of life [*see Data*].

Data

Human Data

In study AI424-182, atazanavir /ritonavir (300/100 mg or 400/100 mg) coadministered with lamivudine/zidovudine (150 mg/ 300 mg, as fixed-dose product) was administered to 41 pregnant women with HIV-1 infection, during the second or third trimester. Among the 39 women who completed the study, 38 women achieved an HIV-1 RNA less than 50 copies/mL at time of delivery. Six of 20 (30%) women on atazanavir /ritonavir 300/100 mg and 13 of 21 (62%) women on atazanavir /ritonavir 400/100 mg experienced hyperbilirubinemia (total bilirubin greater than or equal to 2.6 times ULN). There were no cases of lactic acidosis observed in clinical trial AI424-182.

Atazanavir drug concentrations in fetal umbilical cord blood were approximately 12% to 19% of maternal concentrations. Among the 40 infants born to 40 pregnant women with HIV-1 infection, all had test results that were negative for HIV-1 DNA at the time of delivery and/or during the first 6 months postpartum. All 40 infants received antiretroviral prophylactic treatment containing zidovudine. No evidence of severe hyperbilirubinemia (total bilirubin levels greater than 20 mg/dL) or acute or chronic bilirubin encephalopathy was observed among neonates in this study. However, 10/36 (28%) infants (6 greater than

or equal to 38 weeks gestation and 4 less than 38 weeks gestation) had bilirubin levels of 4 mg/dL or greater within the first day of life.

Lack of ethnic diversity was a study limitation. In the study population, 33/40 (83%) infants were Black/African American, who have a lower incidence of neonatal hyperbilirubinemia than Caucasians and Asians. In addition, women with Rh incompatibility were excluded, as well as women who had a previous infant who developed hemolytic disease and/or had neonatal pathologic jaundice (requiring phototherapy). Additionally, of the 38 infants who had glucose samples collected in the first day of life, 3 had adequately collected serum glucose samples with values of less than 40 mg/dL that could not be attributed to maternal glucose intolerance, difficult delivery, or sepsis.

Based on prospective reports from the APR of approximately 1600 live births following exposure to atazanavir-containing regimens (including 1037 live births in infants exposed in the first trimester and 569 exposed in second/third trimesters), there was no difference between atazanavir and overall birth defects compared with the background birth defect rate. In the U.S. general population, the estimated background risk of major birth defects in clinically recognized pregnancies is 2-4%.

Animal Data

In animal reproduction studies, there was no evidence of mortality or teratogenicity in offspring born to animals at systemic drug exposure levels (AUC) 0.7 (in rabbits) to 1.2 (in rats) times those observed at the human clinical dose (300 mg/day atazanavir boosted with 100 mg/day ritonavir). In pre- and postnatal development studies in the rat, atazanavir caused neonatal growth retardation during lactation that reversed after weaning. Maternal drug exposure at this dose was 1.3 times the human exposure at the recommended clinical exposure. Minimal maternal toxicity occurred at this exposure level.

8.2. Lactation

Risk Summary

The Centers for Disease Control and Prevention recommend that patients with HIV-1 infection, not breastfeed their infants to avoid risking postnatal transmission of HIV-1. Atazanavir has been detected in human milk. No data are available regarding atazanavir effects on milk production. Atazanavir was present in the milk of lactating rats and was associated with neonatal growth retardation that reversed after weaning.

Because of both the potential for HIV-1 transmission and the potential for serious adverse reactions in breastfed infants, advise women not to breastfeed.

8.3. Pediatric Use

Atazanavir Teva is indicated in combination with other antiretroviral agents for the treatment of pediatric

patients with HIV-1 infection, in pediatric patients 6 years of age and older. Atazanavir Teva is not recommended for use in pediatric patients below the age of 3 months due to the risk of kernicterus [see *Indications and Usage (1)*]. All Atazanavir Teva contraindications, warnings, and precautions apply to pediatric patients [see *Contraindications (4) and Warnings and Precautions (5)*].

The safety, pharmacokinetic profile, and virologic response of atazanavir in pediatric patients at least 3 months of age and older weighing at least 5 kg were established in three open-label, multicenter clinical trials: PACTG 1020A, AI424-451, and AI424-397 [see *Clinical Pharmacology (11.3) and Clinical Studies (13.3)*]. The safety profile in pediatric patients was generally similar to that observed in adults [see *Adverse Reactions (6.1)*]. See *Dosage and Administration (2.4,2.5)* for dosing recommendations for the use of Atazanavir Teva capsules in pediatric patients.

8.4. Geriatric Use

Clinical studies of atazanavir did not include sufficient numbers of patients aged 65 and over to determine whether they respond differently from younger patients. Based on a comparison of mean single-dose pharmacokinetic values for C_{max} and AUC, a dose adjustment based upon age is not recommended. In general, appropriate caution should be exercised in the administration and monitoring of atazanavir in elderly patients reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy.

8.5. Age/Gender

A study of the pharmacokinetics of atazanavir was performed in young (n=29; 18-40 years) and elderly (n=30; ≥65 years) healthy subjects. There were no clinically significant pharmacokinetic differences observed due to age or gender.

8.6. Impaired Renal Function

Atazanavir Teva is not recommended for use in treatment-experienced patients with HIV-1 infection, who have end-stage renal disease managed with hemodialysis [see *Dosage and Administration (2.7) and Clinical Pharmacology (11.3)*].

8.7. Impaired Hepatic Function

Atazanavir Teva is not recommended for use in patients with severe hepatic impairment. Atazanavir Teva/ritonavir is not recommended in patients with any degree of hepatic impairment [see *Dosage and Administration (2.8) and Clinical Pharmacology (11.3)*].

9. OVERDOSAGE


Human experience of acute overdose with atazanavir is limited. Single doses up to 1200 mg (three times the 400 mg maximum recommended dose) have been taken by healthy subjects without symptomatic untoward effects. A single self-administered overdose of 29.2 g of atazanavir in a patient with HIV-1 infection (73 times the 400-mg recommended dose) was associated with asymptomatic bifascicular block and PR interval prolongation. These events resolved spontaneously. At atazanavir doses resulting in high atazanavir exposures, jaundice due to indirect (unconjugated) hyperbilirubinemia (without associated liver function test changes) or PR interval prolongation may be observed [see Warnings and Precautions (5.1, 5.7) and Clinical Pharmacology (11.2)].

Treatment of overdosage with atazanavir should consist of general supportive measures, including monitoring of vital signs and ECG, and observations of the patient's clinical status. If indicated, elimination of unabsorbed atazanavir should be achieved by emesis or gastric lavage. Administration of activated charcoal may also be used to aid removal of unabsorbed drug. There is no specific antidote for overdose with atazanavir. Since atazanavir is extensively metabolized by the liver and is highly protein bound, dialysis is unlikely to be beneficial in significant removal of this medicine.

10. DESCRIPTION

The active ingredient in Atazanavir Tevacapsules is atazanavir sulfate, which is an HIV-1 protease inhibitor.

The chemical name for atazanavir sulfate is (3S,8S,9S,12S)-3,12-Bis(1,1-dimethylethyl)-8-hydroxy-4,11-dioxo-9-(phenylmethyl)-6-[[4-(2-pyridinyl)phenyl]methyl]-2,5,6,10,13-pentaazatetradecanedioic acid dimethyl ester, sulfate (1:1). Its molecular formula is $C_{38}H_{52}N_6O_7 \cdot H_2SO_4$, which corresponds to a molecular weight of 802.9 (sulfuric acid salt). The free base molecular weight is 704.9. Atazanavir sulfate has the following structural formula:



Atazanavir sulfate is a white to pale-yellow crystalline powder. It is slightly soluble in water (4-5 mg/mL, free base equivalent) with the pH of a saturated solution in water being about 1.9 at $24 \pm 3^\circ\text{C}$.

Atazanavir Teva Capsules are available for oral administration in strengths of 150 mg, 200 mg, or 300 mg of atazanavir, which are equivalent to 170.8 mg, 227.8 mg, or 341.69 mg of atazanavir sulfate, respectively.

The capsules also contain the following inactive ingredients: Lactose monohydrate, Crospovidone type A, magnesium stearate

The hard gelatin capsule shells contain: gelatin, titanium dioxide (E171), Indigotine (FD&C Blue No. 2).

The 300 mg capsule shells also contain: red iron oxide (E172) and yellow iron oxide (E172).

The capsules are printed with ink containing shellac glaze, iron oxide black, n-butyl alcohol, purified water, propylene glycol, dehydrated ethanol, isopropyl alcohol, ammonia solution.

11. CLINICAL PHARMACOLOGY

11.1. Mechanism of Action

Atazanavir is an HIV-1 antiretroviral drug [see *Microbiology (11.4)*].

11.2. Pharmacodynamics

Cardiac Electrophysiology

Concentration- and dose-dependent prolongation of the PR interval in the electrocardiogram has been observed in healthy subjects receiving atazanavir. In a placebo-controlled study AI424-076, the mean (\pm SD) maximum change in PR interval from the predose value was 24 (\pm 15) msec following oral dosing with 400 mg of atazanavir (n=65) compared to (\pm 11) msec following dosing with placebo (n=67). The PR interval prolongations in this study were asymptomatic. There is limited information on the potential for a pharmacodynamic interaction in humans between atazanavir and other drugs that prolong the PR interval of the electrocardiogram [see *Warnings and Precautions (5.1)*].

Electrocardiographic effects of atazanavir were determined in a clinical pharmacology study of 72 healthy subjects. Oral doses of 400 mg (maximum recommended dosage) and 800 mg (twice the maximum recommended dosage) were compared with placebo; there was no concentration-dependent effect of atazanavir on the QTc interval (using Fridericia's correction). In 1793 subjects with HIV-1 infection receiving antiretroviral regimens, QTc prolongation was comparable in the atazanavir and comparator regimens. No atazanavir-treated healthy subject or subjects with HIV-1 infection in clinical trials had a QTc interval $>$ 500 msec [see *Warnings and Precautions (5.1)*].

11.3. Pharmacokinetics

The pharmacokinetics of atazanavir were evaluated in healthy adult subjects who either were healthy, or with HIV infection, after administration of atazanavir 400 mg once daily and after administration of

atazanavir 300 mg with ritonavir 100 mg once daily (see Table 17).

Table 17: Steady-State Pharmacokinetics of atazanavir in Healthy Subjects or Subjects with HIV-1 Infection in the Fed State


Parameter	400 mg once daily		300 mg with ritonavir 100 mg once daily	
	Healthy Subjects (n=14)	Subjects with HIV-1 Infection (n=13)	Healthy Subjects (n=28)	Subjects with HIV-1 Infection (n=10)
C_{max} (ng/mL)				
Geometric mean (CV%)	5199 (26)	2298 (71)	6129 (31)	4422 (58)
Mean (SD)	5358 (1371)	3152 (2231)	6450 (2031)	5233 (3033)
T_{max} (h)				
Median	2.5	2.0	2.7	3.0
AUC (ng*h/mL)				
Geometric mean (CV%)	28132 (28)	14874 (91)	57039 (37)	46073 (66)
Mean (SD)	29303 (8263)	22262 (20159)	61435 (22911)	53761 (35294)
T-half (h)				
Mean (SD)	7.9 (2.9)	6.5 (2.6)	18.1 (6.2) ^a	8.6 (2.3)
C_{min} (ng/mL)				
Geometric mean (CV%)	159 (88)	120 (109)	1227 (53)	636 (97)
Mean (SD)	218 (191)	273 (298) ^b	1441 (757)	862 (838)

a. n=26

b. n=12

Figure 1 displays the mean plasma concentrations of atazanavir at steady state after atazanavir 400 mg once daily (as two 200-mg capsules) with a light meal and after atazanavir 300 mg (as two 150-mg capsules) with ritonavir 100 mg once daily with a light meal in adult subjects with HIV-1 infection.

Figure 1 : Mean (SD) Steady-State Plasma Concentrations of Atazanavir 400 mg (n=13) and 300 mg with Ritonavir (n=10) for HIV-Infected Adult subjects with HIV-1 Infection



Absorption

Atazanavir is rapidly absorbed with a T_{max} of approximately 2.5 hours. Atazanavir demonstrates nonlinear pharmacokinetics with greater than dose-proportional increases in AUC and C_{max} values over the dose range of 200 to 800 mg once daily. Steady state is achieved between Days 4 and 8, with an accumulation of approximately 2.3 fold.

Food Effect

Administration of atazanavir with food enhances bioavailability and reduces pharmacokinetic variability. Administration of a single 400-mg dose of atazanavir with a light meal (357 kcal, 8.2 g fat, 10.6 g protein) resulted in a 70% increase in AUC and 57% increase in C_{max} relative to the fasting state. Administration of a single 400-mg dose of atazanavir with a high-fat meal (721 kcal, 37.3 g fat, 29.4 g protein) resulted in a mean increase in AUC of 35% with no change in C_{max} relative to the fasting state. Administration of atazanavir with either a light meal or high-fat meal decreased the coefficient of variation of AUC and C_{max} by approximately one-half compared to the fasting state.

Coadministration of a single 300-mg dose of atazanavir and a 100-mg dose of ritonavir with a light meal (336 kcal, 5.1 g fat, 9.3 g protein) resulted in a 33% increase in the AUC and a 40% increase in both the C_{max} and the 24-hour concentration of atazanavir relative to the fasting state. Coadministration with a high-fat meal (951 kcal, 54.7 g fat, 35.9 g protein) did not affect the AUC of atazanavir relative to fasting conditions and the C_{max} was within 11% of fasting values. The 24-hour concentration following a high-fat meal was increased by approximately 33% due to delayed absorption; the median T_{max} increased from 2.0 to 5.0 hours.

Coadministration of Atazanavir with ritonavir with either a light or a high-fat meal decreased the coefficient of variation of AUC and C_{max} by approximately 25% compared to the fasting state.

Distribution

Atazanavir is 86% bound to human serum proteins and protein binding is independent of concentration. Atazanavir binds to both alpha-1-acid glycoprotein (AAG) and albumin to a similar extent (89% and 86%, respectively). In a multiple-dose study in subjects with HIV-1 infection dosed with atazanavir 400 mg once daily with a light meal for 12 weeks, atazanavir was detected in the cerebrospinal fluid and semen. The cerebrospinal fluid/plasma ratio for atazanavir (n=4) ranged between 0.0021 and 0.0226 and seminal fluid/plasma ratio (n=5) ranged between 0.11 and 4.42.

Metabolism

Atazanavir is extensively metabolized in humans. The major biotransformation pathways of atazanavir in

humans consisted of monooxygenation and dioxygenation. Other minor biotransformation pathways for atazanavir or its metabolites consisted of glucuronidation, N-dealkylation, hydrolysis, and oxygenation with dehydrogenation. Two minor metabolites of atazanavir in plasma have been characterized. Neither metabolite demonstrated in vitro antiviral activity. In vitro studies using human liver microsomes suggested that atazanavir is metabolized by CYP3A.

Elimination

Following a single 400-mg dose of ¹⁴C-atazanavir, 79% and 13% of the total radioactivity was recovered in the feces and urine, respectively. Unchanged drug accounted for approximately 20% and 7% of the administered dose in the feces and urine, respectively. The mean elimination half-life of atazanavir in healthy subjects (n=214) and adult subjects with HIV-1 infection (n=13) was approximately 7 hours at steady state following a dose of 400 mg daily with a light meal.

Specific Populations

Renal Impairment

In healthy subjects, the renal elimination of unchanged atazanavir was approximately 7% of the administered dose. Atazanavir has been studied in adult subjects with severe renal impairment (n=20), including those on hemodialysis, at multiple doses of 400 mg once daily. The mean atazanavir C_{max} was 9% lower, AUC was 19% higher, and C_{min} was 96% higher in subjects with severe renal impairment not undergoing hemodialysis (n=10), than in age-, weight-, and gender-matched subjects with normal renal function. In a 4-hour dialysis session, 2.1% of the administered dose was removed. When atazanavir was administered either prior to, or following hemodialysis (n=10), the geometric means for C_{max}, AUC, and C_{min} were approximately 25% to 43% lower compared to subjects with normal renal function. The mechanism of this decrease is unknown. Atazanavir is not recommended for use in treatment-experienced patients with HIV-1 who have end stage renal disease managed with hemodialysis [see Dosage and Administration (2.7)].

Hepatic Impairment

Atazanavir has been studied in adult subjects with moderate-to-severe hepatic impairment (14 Child-Pugh B and 2 Child-Pugh C subjects) after a single 400-mg dose. The mean ALC_(0-∞) was 42% greater in subjects with impaired hepatic function than in healthy subjects. The mean half-life of atazanavir in hepatically impaired subjects was 12.1 hours compared to 6.4 hours in healthy subjects. A dose reduction to 300 mg is recommended for patients with moderate hepatic impairment (Child-Pugh Class B) who have not experienced prior virologic failure as increased concentrations of atazanavir are expected. atazanavir is

not recommended for use in patients with severe hepatic impairment. The pharmacokinetics of atazanavir in combination with ritonavir has not been studied in subjects with hepatic impairment; thus, coadministration of atazanavir with ritonavir is not recommended for use in patients with any degree of hepatic impairment [see Dosage and Administration (2.8)].

Pediatrics

The pharmacokinetic parameters for atazanavir at steady state in pediatric subjects taking the capsule formulation were predicted by a population pharmacokinetic model and are summarized in Table 18 by weight ranges that correspond to the recommended doses [see Dosage and Administration (2.4).]

Table 18: Predicted Steady-State Pharmacokinetics of Atazanavir (capsule formulation) with Ritonavir in Pediatric Subjects with HIV-1 Infection

Body Weight (range in kg)	atazanavir/ritonavir Dose (mg)	C_{max} ng/mL Geometric Mean (CV%)	AUC ng*h/mL Geometric Mean (CV%)	C_{min} ng/mL Geometric Mean (CV%)
15 to <35	200/100	3303 (86%)	37235 (84%)	538 (99%)
≥35	300/100	2980 (82%)	37643 (83%)	653 (89%)

Pregnancy

The pharmacokinetic data from pregnant women with HIV-1 infection receiving atazanavir Capsules with ritonavir are presented in Table 19.

Table 19: Steady-State Pharmacokinetics of Atazanavir with Ritonavir in Pregnant Women with HIV-1 Infection in the Fed State

Pharmacokinetic Parameter	Atazanavir 300 mg with ritonavir 100 mg		
	2nd Trimester (n=5^a)	3rd Trimester (n=20)	Postpartum^b (n=34)
C _{max} ng/mL	3078.85	3291.46	5721.21
Geometric mean (CV%)	(50)	(48)	(31)
AUC ng*h/mL	27657.1	34251.5	61990.4
Geometric mean (CV%)	(43)	(43)	(32)
C _{min} ng/mL ^c	538.70	668.48	1462.59
Geometric mean (CV%)	(46)	(50)	(45)

- a. Available data during the 2nd trimester are limited.
- b. Atazanavir peak concentrations and AUCs were found to be approximately 28% to 43% higher during the postpartum period (4-12 weeks) than those observed historically in non-pregnant patients with HIV-1 infection. Atazanavir plasma trough concentrations were approximately 2.2-fold higher during the postpartum period when compared to those observed historically in non-

- pregnant patients with HIV-1 infection.
 c. C_{min} is concentration 24 hours post-dose.

Drug Interaction Data

Atazanavir is a metabolism-dependent CYP3A inhibitor, with a K_{inact} value of 0.05 to 0.06 min^{-1} and K_i value of 0.84 to 1.0 μM . Atazanavir is also a direct inhibitor for UGT1A1 ($K_i=1.9 \mu\text{M}$) and CYP2C8 ($K_i=2.1 \mu\text{M}$).

Atazanavir has been shown in vivo not to induce its own metabolism nor to increase the biotransformation of some drugs metabolized by CYP3A. In a multiple-dose study, atazanavir decreased the urinary ratio of endogenous 6 β -OH cortisol to cortisol versus baseline, indicating that CYP3A production was not induced.

Clinically significant interactions are not expected between atazanavir and substrates of CYP2C19, CYP2C9, CYP2D6, CYP2B6, CYP2A6, CYP1A2, or CYP2E1. Clinically significant interactions are not expected between atazanavir when administered with ritonavir and substrates of CYP2C8. See the complete prescribing information for ritonavir for information on other potential drug interactions with ritonavir.

Based on known metabolic profiles, clinically significant drug interactions are not expected between Atazanavir Teva and dapsone, trimethoprim/sulfamethoxazole, azithromycin, or erythromycin. Atazanavir Teva does not interact with substrates of CYP2D6 (eg, nortriptyline, desipramine, metoprolol).

Drug interaction studies were performed with atazanavir and other drugs likely to be coadministered and some drugs commonly used as probes for pharmacokinetic interactions. The effects of coadministration of atazanavir on the AUC, C_{max} , and C_{min} are summarized in Tables 20 and 21. Neither didanosine EC nor diltiazem had a significant effect on atazanavir exposures (see Table 21 for effect of atazanavir on didanosine EC or diltiazem exposures). Atazanavir did not have a significant effect on the exposures of didanosine (when administered as the buffered tablet), stavudine, or fluconazole. For information regarding clinical recommendations, see *Drug Interactions (7)*.

Table 20: Drug Interactions: Pharmacokinetic Parameters for Atazanavir in the Presence of Coadministered Drugs^a

Coadministered Drug	Coadministered Drug Dose/Schedule	Atazanavir Dose/Schedule	Ratio (90% Confidence Interval) of Atazanavir Pharmacokinetic Parameters with/without Coadministered Drug; No Effect = 1.00		
			C_{max}	AUC	C_{min}
atenolol	50 mg QD, d 7-11 (n=19) and d 19-23	400 mg QD, d 1-11 (n=19)	1.00 (0.89, 1.12)	0.93 (0.85, 1.01)	0.74 (0.65, 0.86)

Coadministered Drug	Coadministered Drug Dose/Schedule	Atazanavir Dose/Schedule	Ratio (90% Confidence Interval) of Atazanavir Pharmacokinetic Parameters with/without Coadministered Drug; No Effect = 1.00		
			C _{max}	AUC	C _{min}
clarithromycin	500 mg BID, d 7-10 (n=29) and d 18-21	400 mg QD, d 1-10 (n=29)	1.06 (0.93, 1.20)	1.28 (1.16, 1.43)	1.91 (1.66, 2.21)
didanosine (ddI) (buffered tablets) plus stavudine (d4T) ^b	ddI: 200 mg x 1 dose, d4T: 40 mg x 1 dose (n=31)	400 mg x 1 dose simultaneously with ddI and d4T (n=31)	0.11 (0.06, 0.18)	0.13 (0.08, 0.21)	0.16 (0.10, 0.27)
	ddI: 200 mg x 1 dose, d4T: 40 mg x 1 dose (n=32)	400 mg x 1 dose 1 h after ddI + d4T (n=32)	1.12 (0.67, 1.18)	1.03 (0.64, 1.67)	1.03 (0.61, 1.73)
efavirenz	600 mg QD, d 7-20 (n=27)	400 mg QD, d 1-20 (n=27)	0.41 (0.33, 0.51)	0.26 (0.22, 0.32)	0.07 (0.05, 0.10)
	600 mg QD, d 7-20 (n=13)	400 mg QD, d 1-6 (n=23) then 300 mg/ritonavir 100 mg QD, 2 h before efavirenz, d 7-20 (n=13)	1.14 (0.83, 1.58)	1.39 (1.02, 1.88)	1.48 (1.24, 1.76)
	600 mg QD, d 11-24 (pm) (n=14)	300 mg QD/ritonavir 100 mg QD, d 1-10 (pm) (n=22), then 400 mg QD/ritonavir 100 mg QD, d 11-24 (pm), (simultaneously with efavirenz) (n=14)	1.17 (1.08, 1.27)	1.00 (0.91, 1.10)	0.58 (0.49, 0.69)
famotidine	40 mg BID, d 7-12 (n=15)	400 mg QD, d 1-6 (n=45), d 7-12 (simultaneous administration) (n=15)	0.53 (0.34, 0.82)	0.59 (0.40, 0.87)	0.58 (0.37, 0.89)
	40 mg BID, d 7-12 (n=14)	400 mg QD (pm), d 1-6 (n=14), d 7-12 (10 h after, 2 h before famotidine) (n=14)	1.08 (0.82, 1.41)	0.95 (0.74, 1.21)	0.79 (0.60, 1.04)
	40 mg BID, d 11-20 (n=14) ^c	300 mg QD/ritonavir 100 mg QD, d 1-10 (n=46), d 11-20d (simultaneous administration) (n=14)	0.86 (0.79, 0.94)	0.82 (0.75, 0.89)	0.72 (0.64, 0.81)
	20 mg BID, d 11-17 (n=18)	300 mg QD/ritonavir 100 mg QD/tenofovir DF 300 mg QD, d 1-10 (am) (n=39), d 11-17 (am) (simultaneous	0.91 (0.84, 0.99)	0.90 (0.82, 0.98)	0.81 (0.69, 0.94)

Coadministered Drug	Coadministered Drug Dose/Schedule	Atazanavir Dose/Schedule	Ratio (90% Confidence Interval) of Atazanavir Pharmacokinetic Parameters with/without Coadministered Drug; No Effect = 1.00		
			C _{max}	AUC	C _{min}
		administration with am famotidine) (n=18) ^{d,e}			
	40 mg QD (pm), d 18-24 (n=20)	300 mg QD/ritonavir 100 mg QD/tenofovir DF 300 mg QD, d 1-10 (am) (n=39), d 18-24 (am) (12 h after pm famotidine) (n=20) ^e	0.89 (0.81, 0.97)	0.88 (0.80, 0.96)	0.77 (0.63, 0.93)
	40 mg BID, d 18-24 (n=18)	300 mg QD/ritonavir 100 mg QD/tenofovir DF 300 mg QD, d 1-10 (am) (n=39), d 18-24 (am) (10 h after pm famotidine and 2 h before am famotidine) (n=18) ^e	0.74 (0.66, 0.84)	0.79 (0.70, 0.88)	0.72 (0.63, 0.83)
	40 mg BID, d 11-20 (n=15)	300 mg QD/ritonavir 100 mg QD, d 1-10 (am) (n=46), then 400 mg QD/ritonavir 100 mg QD, d 11-20 (am) (n=15)	1.02 (0.87, 1.18)	1.03 (0.86, 1.22)	0.86 (0.68, 1.08)
grazoprevir/ elbasvir	grazoprevir 200 mg QD d 1 - 35 (n = 11)	300 mg QD/ritonavir 100 mg QD, d 1- 35 (n = 11)	1.12 (1.01, 1.24)	1.43 (1.30, 1.57)	1.23 (1.13, 1.34)
	elbasvir 50 mg QD d 1 - 35 (n = 8)	300 mg QD/ritonavir 100 mg QD, d 1 - 35 (n = 8)	1.02 (0.96, 1.08)	1.07 (0.98,1.17)	1.15 (1.02, 1.29)
ketoconazole	200 mg QD, d 7-13 (n=14)	400 mg QD, d 1-13 (n=14)	0.99 (0.77, 1.28)	1.10 (0.89, 1.37)	1.03 (0.53, 2.01)
nevirapine ^{f,g}	200 mg BID, d 1-23 (n=23)	300 mg QD/ritonavir 100 mg QD, d 4-13, then	0.72 (0.60, 0.86)	0.58 (0.48, 0.71)	0.28 (0.20, 0.40)
		400 mg QD/ritonavir 100 mg QD, d 14-23 (n=23) ^h	1.02 (0.85, 1.24)	0.81 (0.65, 1.02)	0.41 (0.27, 0.60)
omeprazole	40 mg QD, d 7-12 (n=16) ⁱ	400 mg QD, d 1-6 (n=48), d 7-12 (n=16)	0.04 (0.04, 0.05)	0.06 (0.05, 0.07)	0.05 (0.03, 0.07)
	40 mg QD, d 11-20 (n=15) ⁱ	300 mg QD/ritonavir 100 mg QD, d 1-20 (n=15)	0.28 (0.24, 0.32)	0.24 (0.21, 0.27)	0.22 (0.19, 0.26)

Coadministered Drug	Coadministered Drug Dose/Schedule	Atazanavir Dose/Schedule	Ratio (90% Confidence Interval) of Atazanavir Pharmacokinetic Parameters with/without Coadministered Drug; No Effect = 1.00		
			C _{max}	AUC	C _{min}
	20 mg QD, d 17-23 (am) (n=13)	300 mg QD/ritonavir 100 mg QD, d 7-16 (pm) (n=27), d 17-23 (pm) (n=13) ^{j,k}	0.61 (0.46, 0.81)	0.58 (0.44, 0.75)	0.54 (0.41, 0.71)
	20 mg QD, d 17-23 (am) (n=14)	300 mg QD/ritonavir 100 mg QD, d 7-16 (am) (n=27), then 400 mg QD/ritonavir 100 mg QD, d 17-23 (am) (n=14) ^{l,m}	0.69 (0.58, 0.83)	0.70 (0.57, 0.86)	0.69 (0.54, 0.88)
pitavastatin	4 mg QD for 5 days	300 mg QD for 5 days	1.13 (0.96, 1.32)	1.06 (0.90, 1.26)	NA
rifabutin	150 mg QD, d 15-28 (n=7)	400 mg QD, d 1-28 (n=7)	1.34 (1.14, 1.59)	1.15 (0.98, 1.34)	1.13 (0.68, 1.87)
rifampin	600 mg QD, d 17-26 (n=16)	300 mg QD/ritonavir 100 mg QD, d 7-16 (n=48), d 17-26 (n=16)	0.47 (0.41, 0.53)	0.28 (0.25, 0.32)	0.02 (0.02, 0.03)
ritonavir ⁿ	100 mg QD, d 11-20 (n=28)	300 mg QD, d 1-20 (n=28)	1.86 (1.69, 2.05)	3.38 (3.13, 3.63)	11.89 (10.23, 13.82)
tenofovir DF ^o	300 mg QD, d 9-16 (n=34)	400 mg QD, d 2-16 (n=34)	0.79 (0.73, 0.86)	0.75 (0.70, 0.81)	0.60 (0.52, 0.68)
	300 mg QD, d 15-42 (n=10)	300 mg/ritonavir 100 mg QD, d 1-42 (n=10)	0.72 ^p (0.50, 1.05)	0.75 ^p (0.58, 0.97)	0.77 ^p (0.54, 1.10)
voriconazole (Subjects with at least one functional CYP2C19 allele)	200 mg BID, d 2-3, 22-30; 400 mg BID, d 1, 21 (n=20)	300 mg/ritonavir 100 mg QD, d 11-30 (n=20)	0.87 (0.80, 0.96)	0.88 (0.82, 0.95)	0.80 (0.72, 0.90)
voriconazole (Subjects without a functional CYP2C19 allele)	50 mg BID, d 2-3, 22-30; 100 mg BID, d 1, 21 (n=8)	300 mg/ritonavir 100 mg QD, d 11-30 (n=8)	0.81 (0.66, 1.00)	0.80 (0.65, 0.97)	0.69 (0.54, 0.87)

- Data provided are under fed conditions unless otherwise noted.
- All drugs were given under fasted conditions.
- Atazanavir 300 mg with ritonavir 100 mg once daily coadministered with famotidine 40 mg twice daily resulted in atazanavir geometric mean C_{max} that was similar and AUC and C_{min} values that were 1.79- and 4.46-fold higher relative to Atazanavir 400 mg once daily alone.
- Similar results were noted when famotidine 20 mg BID was administered 2 hours after and 10 hours before atazanavir 300 mg with ritonavir 100 mg and tenofovir DF 300 mg.
- coadministration of Atazanavir with ritonavir 100 mg and tenofovir DF was administered after a

- light meal.
- f. Study was conducted in subjects with HIV-1 infection.
 - g. Compared with atazanavir 400 mg historical data without nevirapine (n=13), the ratio of geometric means (90% confidence intervals) for C_{max} , AUC, and C_{min} were 1.42 (0.98, 2.05), 1.64 (1.11, 2.42), and 1.25 (0.66, 2.36), respectively, for atazanavir/ritonavir 300/100 mg; and 2.02 (1.42, 2.87), 2.28 (1.54, 3.38), and 1.80 (0.94, 3.45), respectively, for atazanavir/ritonavir 400/100 mg.
 - h. Parallel group design; n=23 for atazanavir/ritonavir and nevirapine, n=22 for atazanavir 300 mg/ritonavir 100 mg without nevirapine. Subjects were treated with nevirapine prior to study entry.
 - i. Omeprazole 40 mg was administered on an empty stomach 2 hours before atazanavir.
 - j. Omeprazole 20 mg was administered 30 minutes prior to a light meal in the morning and atazanavir 300 mg with ritonavir 100 mg in the evening after a light meal, separated by 12 hours from omeprazole.
 - k. Atazanavir 300 mg with ritonavir 100 mg once daily separated by 12 hours from omeprazole 20 mg daily resulted in increases in atazanavir geometric mean AUC (10%) and C_{min} (2.4-fold), with a decrease in C_{max} (29%) relative to atazanavir 400 mg once daily in the absence of omeprazole (study days 1-6).
 - l. Omeprazole 20 mg was given 30 minutes prior to a light meal in the morning and atazanavir 400 mg with ritonavir 100 mg once daily after a light meal, 1 hour after omeprazole. Effects on atazanavir concentrations were similar when atazanavir 400 mg with ritonavir 100 mg was separated from omeprazole 20 mg by 12 hours.
 - m. Atazanavir 400 mg with ritonavir 100 mg once daily administered with omeprazole 20 mg once daily resulted in increases in atazanavir geometric mean AUC (32%) and C_{min} (3.3-fold), with a decrease in C_{max} (26%) relative to atazanavir 400 mg once daily in the absence of omeprazole (study days 1-6).
 - n. Compared with atazanavir 400 mg QD historical data, administration of atazanavir/ritonavir 300/100 mg QD increased the atazanavir geometric mean values of C_{max} , AUC, and C_{min} by 18%, 103%, and 671%, respectively.
 - o. Note that similar results were observed in studies where administration of tenofovir DF and atazanavir was separated by 12 hours.
 - p. Ratio of atazanavir with ritonavir and tenofovir DF to atazanavir with ritonavir. Atazanavir 300 mg with ritonavir 100 mg results in higher atazanavir exposure than atazanavir 400 mg (see footnote o). The geometric mean values of atazanavir pharmacokinetic parameters when coadministered with ritonavir and tenofovir DF were: C_{max} = 3190 ng/mL, AUC = 34459 ng*h/mL, and C_{min} = 491 ng/mL. Study was conducted in subjects with HIV-1 infection.
- NA = not available.

Table 21: Drug Interactions: Pharmacokinetic Parameters for Coadministered Drugs in the Presence of Atazanavir^a

Coadministered Drug	Coadministered Drug Dose/Schedule	Atazanavir Dose/Schedule	Ratio (90% Confidence Interval) of Coadministered Drug Pharmacokinetic Parameters with/without atazanavir; No Effect = 1.00		
			C_{max}	AUC	C_{min}
acetaminophen	1 gm BID, d 1-20 (n=10)	300 mg QD/ritonavir 100 mg QD, d 11-	0.87 (0.77, 0.99)	0.97 (0.91, 1.03)	1.26 (1.08, 1.46)

Coadministered Drug	Coadministered Drug Dose/Schedule	Atazanavir Dose/Schedule	Ratio (90% Confidence Interval) of Coadministered Drug Pharmacokinetic Parameters with/without atazanavir; No Effect = 1.00		
			C _{max}	AUC	C _{min}
		20 (n=10)			
atenolol	50 mg QD, d 7-11 (n=19) and d 19-23	400 mg QD, d 1-11 (n=19)	1.34 (1.26, 1.42)	1.25 (1.16, 1.34)	1.02 (0.88, 1.19)
clarithromycin	500 mg BID, d 7-10 (n=21) and d18-21	400 mg QD, d 1-10 (n=21)	1.50 (1.32, 1.71) OH-clarithromycin: 0.28 (0.24, 0.33)	1.94 (1.75, 2.16) OH-clarithromycin: 0.30 (0.26, 0.34)	2.60 (2.35, 2.88) OH-clarithromycin: 0.38 (0.34, 0.42)
ddI (enteric-coated [EC] capsules) ^b	400 mg d 1 (fasted), d 8 (fed) (n=34)	400 mg QD, d 2-8 (n=34)	0.64 (0.55, 0.74)	0.66 (0.60, 0.74)	1.13 (0.91, 1.41)
	400 mg d 1 (fasted), d 19 (fed) (n=31)	300 mg QD/ritonavir 100 mg QD, d 9-19 (n=31)	0.62 (0.52, 0.74)	0.66 (0.59, 0.73)	1.25 (0.92, 1.69)
diltiazem	180 mg QD, d 7-11 (n=28) and d 19-23	400 mg QD, d 1-11 (n=28)	1.98 (1.78, 2.19) desacetyl-diltiazem: 2.72 (2.44, 3.03)	2.25 (2.09, 2.16) desacetyl-diltiazem: 2.65 (2.45, 2.87)	2.42 (2.14, 2.73) desacetyl-diltiazem: 2.21 (2.02, 2.42)
ethinyl estradiol & norethindrone ^c	Ortho-Novum [®] 7/7/7 QD, d 1-29 (n=19)	400 mg QD, d16-29 (n=19)	ethinyl estradiol: 1.15 (0.99, 1.32) norethindrone: 1.67 (1.42, 1.96)	ethinyl estradiol: 1.48 (1.31, 1.68) norethindrone: 2.10 (1.68, 2.62)	ethinyl estradiol: 1.91 (1.57, 2.33) norethindrone: 3.62 (2.57, 5.09)
ethinyl estradiol & norgestimate ^d	Ortho Tri-Cyclen [®] QD, d 1-28 (n=18), then Ortho Tri-Cyclen [®] LO QD, d 29-42 ^e (n=14)	300 mg QD/ritonavir 100 mg QD, d29-42 (n=14)	ethinyl estradiol: 0.84 (0.74, 0.95) 17-deacetyl norgestimate: ^f 1.68 (1.51, 1.88)	ethinyl estradiol: 0.81 (0.75, 0.87) 17-deacetyl norgestimate: ^f 1.85 (1.67, 2.05)	ethinyl estradiol: 0.63 (0.55, 0.71) 17-deacetyl norgestimate: ^f 2.02 (1.77, 2.31)
glecaprevir/pibrentasvir	300 mg glecaprevir (n=12)	300 mg QD/ritonavir 100 mg QD (n=12)	≥4.06 ^g (3.15, 5.23)	≥6.53 ^g (5.24, 8.14)	≥14.3 ^g (9.85, 20.7)
	120 mg pibrentasvir (n=12)	300 mg QD/ritonavir 100 mg QD	≥1.29 ^g (1.15, 1.45)	≥1.64 ^g (1.48, 1.82)	≥2.29 ^g (1.95, 2.68)

Coadministered Drug	Coadministered Drug Dose/Schedule	Atazanavir Dose/Schedule	Ratio (90% Confidence Interval) of Coadministered Drug Pharmacokinetic Parameters with/without atazanavir; No Effect = 1.00		
			C _{max}	AUC	C _{min}
		(n=12)			
grazoprevir/ elbasvir	grazoprevir 200 mg QD d 1 - 35 (n = 12)	300 mg QD/ritonavir 100 mg QD d 1 - 35 (n=12)	6.24 (4.42, 8.81)	10.58 (7.78, 14.39)	11.64 (7.96, 17.02)
	elbasvir 50 mg QD d 1 - 35 (n = 10)	300 mg QD/ritonavir 100 mg QD d 1 - 35 (n=10)	4.15 (3.46, 4.97)	4.76 (4.07, 5.56)	6.45 (5.51 7.54)
methadone	Stable maintenance dose, d 1-15 (n=16)	400 mg QD, d 2-15 (n=16)	(R)-methadone ^h 0.91 (0.84, 1.0) total: 0.85 (0.78, 0.93)	(R)-methadone ^h 1.03 (0.95, 1.10) total: 0.94 (0.87, 1.02)	(R)-methadone ^h 1.11 (1.02, 1.20) total: 1.02 (0.93, 1.12)
nevirapine, ^{i,j}	200 mg BID, d 1-23 (n=23)	300 mg QD/ ritonavir 100 mg QD, d 4-13, then 400 mg QD/ ritonavir 100 mg QD, d 14-23 (n=23)	1.17 (1.09, 1.25) 1.21 (1.11, 1.32)	1.25 (1.17, 1.34) 1.26 (1.17, 1.36)	1.32 (1.22, 1.43) 1.35 (1.25, 1.47)
omeprazole ^k	40 mg single dose, d 7 and d 20 (n=16)	400 mg QD, d 1-12 (n=16)	1.24 (1.04, 1.47)	1.45 (1.20, 1.76)	NA
rifabutin	300 mg QD, d 1-10 then 150 mg QD, d 11-20 (n=3)	600 mg QD, ^l d 11-20 (n=3)	1.18 (0.94, 1.48) 25-O-desacetyl-rifabutin: 8.20 (5.90, 11.40)	2.10 (1.57, 2.79) 25-O-desacetyl-rifabutin: 22.01 (15.97, 30.34)	3.43 (1.98, 5.96) 25-O-desacetyl-rifabutin: 75.6 (30.1, 190.0)
	150 mg twice weekly, d 1-15 (n=7)	300 mg QD/ ritonavir 100 mg QD, d 1-17 (n=7)	2.49 ^m (2.03, 3.06) 25-O-desacetyl-rifabutin: 7.77 (6.13, 9.83)	1.48 ^m (1.19, 1.84) 25-O-desacetyl-rifabutin: 10.90 (8.14, 14.61)	1.40 ^m (1.05, 1.87) 25-O-desacetyl-rifabutin: 11.45 (8.15, 16.10)
pitavastatin	4 mg QD for 5 days	300 mg QD for 5 days	1.60 (1.39, 1.85)	1.31 (1.23, 1.39)	NA
rosiglitazone ⁿ	4 mg single dose, d 1, 7, 17 (n=14)	400 mg QD, d 2-7, then 300 mg QD/ ritonavir 100 mg QD, d 8-17 (n=14)	1.08 (1.03, 1.13) 0.97 (0.91, 1.04)	1.35 (1.26, 1.44) 0.83 (0.77, 0.89)	NA NA
rosuvastatin	10 mg single	300 mg QD/	↑7-fold ^o	↑ 3-fold ^o	NA

Coadministered Drug	Coadministered Drug Dose/Schedule	Atazanavir Dose/Schedule	Ratio (90% Confidence Interval) of Coadministered Drug Pharmacokinetic Parameters with/without atazanavir; No Effect = 1.00		
			C _{max}	AUC	C _{min}
	dose	ritonavir 100 mg QD for 7 days			
saquinavir ^p (soft gelatin capsules)	1200 mg QD, d 1-13 (n=7)	400 mg QD, d 7-13 (n=7)	4.39 (3.24, 5.95)	5.49 (4.04, 7.47)	6.86 (5.29, 8.91)
sofosbuvir/ velpatasvir/ voxilaprevir	400 mg sofosbuvir single dose (n=15)	300 mg/100 mg ritonavir single dose (n=15)	1.29 (1.09, 1.52) sofosbuvir metabolite GS-331007 1.05 (0.99, 1.12)	1.40 (1.25, 1.57) sofosbuvir metabolite GS-331007 1.25 (1.16, 1.36)	NA
	100 mg velpatasvir single dose (n=15)	300 mg/100 mg ritonavir single dose (n=15)	1.29 (1.07, 1.56)	1.93 (1.58, 2.36)	NA
	100 mg voxilaprevir single dose (n=15)	300 mg/100 mg ritonavir single dose (n=15)	4.42 (3.65, 5.35)	4.31 (3.76, 4.93)	NA
Tenofovir DF ^a	300 mg QD, d 9-16 (n=33) and d 24-30 (n=33)	400 mg QD, d 2-16 (n=33)	1.14 (1.08, 1.20)	1.24 (1.21, 1.28)	1.22 (1.15, 1.30)
	300 mg QD, d 1-7 (pm) (n=14) d 25-34 (pm) (n=12)	300 mg QD/ritonavir 100 mg QD, d 25-34 (am) (n=12) ^f	1.34 (1.20, 1.51)	1.37 (1.30, 1.45)	1.29 (1.21, 1.36)
voriconazole (Subjects with at least one functional CYP2C19 allele)	200 mg BID, d 2-3, 22-30; 400 mg BID, d 1, 21 (n=20)	300 mg/ritonavir 100 mg QD, d 11-30 (n=20)	0.90 (0.78, 1.04)	0.67 (0.58, 0.78)	0.61 (0.51, 0.72)
voriconazole (Subjects without a functional CYP2C19 allele)	50 mg BID, d 2-3, 22-30; 100 mg BID, d 1, 21 (n=8)	300 mg/ritonavir 100 mg QD, d 11-30 (n=8)	4.38 (3.55, 5.39)	5.61 (4.51, 6.99)	7.65 (5.71, 10.2)
lamivudine + zidovudine	150 mg lamivudine + 300 mg zidovudine BID, d 1-12 (n=19)	400 mg QD, d 7-12 (n=19)	lamivudine: 1.04 (0.92, 1.16) zidovudine: 1.05 (0.88, 1.24) zidovudine glucuronide: 0.95 (0.88, 1.02)	lamivudine: 1.03 (0.98, 1.08) zidovudine: 1.05 (0.96, 1.14) zidovudine glucuronide: 1.00 (0.97, 1.03)	lamivudine: 1.12 (1.04, 1.21) zidovudine: 0.69 (0.57, 0.84) zidovudine glucuronide: 0.82 (0.62, 1.08)

- a. Data provided are under fed conditions unless otherwise noted.
- b. 400 mg dDI EC and atazanavir were administered together with food on Days 8 and 19.
- c. Upon further dose normalization of ethinyl estradiol 25 mcg with atazanavir relative to ethinyl estradiol 35 mcg without atazanavir, the ratio of geometric means (90% confidence intervals) for C_{max}, AUC, and C_{min} were 0.82 (0.73, 0.92), 1.06 (0.95, 1.17), and 1.35 (1.11, 1.63), respectively.
- d. Upon further dose normalization of ethinyl estradiol 35 mcg with atazanavir/ritonavir relative to ethinyl estradiol 25 mcg without atazanavir/ritonavir, the ratio of geometric means (90% confidence intervals) for C_{max}, AUC, and C_{min} were 1.17 (1.03, 1.34), 1.13 (1.05, 1.22), and 0.88 (0.77, 1.00), respectively.
- e. All subjects were on a 28 day lead-in period; one full cycle of Ortho Tri-Cyclen®. Ortho Tri-Cyclen® contains 35 mcg of ethinyl estradiol. Ortho Tri-Cyclen® LO contains 25 mcg of ethinyl estradiol. Results were dose normalized to an ethinyl estradiol dose of 35 mcg.
- f. 17-deacetyl norgestimate is the active component of norgestimate.
- g. Effect of atazanavir and ritonavir on the first dose of glecaprevir and pibrentasvir is reported.
- h. (R)-methadone is the active isomer of methadone.
- i. Study was conducted in subjects with HIV-1 infection.
- j. Subjects were treated with nevirapine prior to study entry.
- k. Omeprazole was used as a metabolic probe for CYP2C19. Omeprazole was given 2 hours after atazanavir on Day 7; and was given alone 2 hours after a light meal on Day 20.
- l. Not the recommended therapeutic dose of atazanavir.
- m. When compared to rifabutin 150 mg QD alone d1-10 (n=14). Total of rifabutin + 25-O-desacetyl-rifabutin: AUC 2.19 (1.78, 2.69).
- n. Rosiglitazone used as a probe substrate for CYP2C8.
- o. Mean ratio (with/without coadministered drug). ↑ indicates an increase in rosuvastatin exposure.
- p. The combination of atazanavir and saquinavir 1200 mg QD produced daily saquinavir exposures similar to the values produced by the standard therapeutic dosing of saquinavir at 1200 mg TID. However, the C_{max} is about 79% higher than that for the standard dosing of saquinavir (soft gelatin capsules) alone at 1200 mg TID.
- q. Note that similar results were observed in a study where administration of tenofovir DF and atazanavir was separated by 12 hours.
- r. Administration of tenofovir DF and atazanavir was temporally separated by 12 hours.
NA = not available.

11.4. Microbiology

Mechanism of Action

Atazanavir (ATV) is an azapeptide HIV-1 protease inhibitor (PI). The compound selectively inhibits the virus-specific processing of viral Gag and Gag-Pol polyproteins in HIV-1 infected cells, thus preventing formation of mature virions.

Antiviral Activity in Cell Culture

Atazanavir exhibits anti-HIV-1 activity with a mean 50% effective concentration (EC₅₀) in the absence of human serum of 2 to 5 nM against a variety of laboratory and clinical HIV-1 isolates grown in peripheral blood mononuclear cells, macrophages, CEM-SS cells, and MT-2 cells. Atazanavir has activity against

HIV-1 Group M subtype viruses A, B, C, D, AE, AG, F, G, and J isolates in cell culture. Atazanavir has variable activity against HIV-2 isolates (1.9-32 nM), with EC50 values above the EC50 values of failure isolates. Two-drug combination antiviral activity studies with atazanavir showed no antagonism in cell culture with PIs (amprenavir, indinavir, lopinavir, nelfinavir, ritonavir, and saquinavir), NNRTIs (delavirdine, efavirenz, and nevirapine), NRTIs (abacavir, didanosine, emtricitabine, lamivudine, stavudine, tenofovir DF, and zidovudine), the HIV-1 fusion inhibitor enfuvirtide, and two compounds used in the treatment of viral hepatitis, adefovir and ribavirin, without enhanced cytotoxicity.

Resistance

In Cell Culture: HIV-1 isolates with a decreased susceptibility to atazanavir have been selected in cell culture and obtained from patients treated with atazanavir or atazanavir/ritonavir. HIV-1 isolates with 93- to 183-fold reduced susceptibility to atazanavir from three different viral strains were selected in cell culture by 5 months. The substitutions in these HIV-1 viruses that contributed to atazanavir resistance include I50L, N88S, I84V, A71V, and M46I. Changes were also observed at the protease cleavage sites following drug selection. Recombinant viruses containing the I50L substitution without other major PI substitutions were growth impaired and displayed increased susceptibility in cell culture to other PIs (amprenavir, indinavir, lopinavir, nelfinavir, ritonavir, and saquinavir). The I50L and I50V substitutions yielded selective resistance to atazanavir and amprenavir, respectively, and did not appear to be cross-resistant.

Clinical Studies of Treatment-Naive Patients: Comparison of Ritonavir-Boosted atazanavir vs. Unboosted atazanavir : Study AI424-089 compared atazanavir 300 mg once daily with ritonavir 100 mg vs. atazanavir 400 mg once daily when administered with lamivudine and extended- release stavudine in treatment-naive subjects with HIV-1 infection . A summary of the number of virologic failures and virologic failure isolates with atazanavir resistance in each arm is shown in Table 22.

Table 22: Summary of Virologic Failures^a at Week 96 in Study AI424-089: Comparison of Ritonavir Boosted atazanavir vs. Unboosted Atazanavir: Randomized Patients

	Atazanavir 300 mg + ritonavir 100 mg (n=95)	Atazanavir 400 mg (n=105)
Virologic Failure (≥ 50 copies/mL) at Week 96	15 (16%)	34 (32%)
Virologic Failure with Genotypes and Phenotypes Data	5	17
Virologic Failure Isolates with ATV-resistance at Week 96	0/5 (0%) ^b	4/17 (24%) ^b
Virologic Failure Isolates with I50L Emergence at Week 96 ^c	0/5 (0%) ^b	2/17 (12%) ^b

Virologic Failure Isolates with Lamivudine Resistance at Week 96	2/5 (40%) ^b	11/17 (65%) ^b
--	------------------------	--------------------------

^a Virologic failure includes subjects who were never suppressed through Week 96 and on study at Week 96, had virologic rebound or discontinued due to insufficient viral load response.

^b Percentage of Virologic Failure Isolates with genotypic and phenotypic data.

^c Mixture of I50I/L emerged in 2 other atazanavir 400 mg-treated subjects. Neither isolate was phenotypically resistant to atazanavir.

Clinical Studies of Treatment-Naive subjects Receiving atazanavir 300 mg with Ritonavir 100 mg: In Phase 3 Study AI424-138, an as-treated genotypic and phenotypic analysis was conducted on samples from subjects who experienced virologic failure (HIV-1 RNA \geq 400 copies/mL) or discontinued before achieving suppression on atazanavir with ritonavir (n=39; 9%) and lopinavir/ritonavir (n=39; 9%) through 96 weeks of treatment. In the atazanavir with ritonavir arm, one of the virologic failure isolates had a 56-fold decrease in atazanavir susceptibility emerge on therapy with the development of PI resistance-associated substitutions L10F, V32I, K43T, M46I, A71I, G73S, I85I/V, and L90M. The NRTI resistance-associated substitution M184V also emerged on treatment in this isolate conferring emtricitabine resistance. Two atazanavir with ritonavir -virologic failure isolates had baseline phenotypic atazanavir resistance and IAS-defined major PI resistance-associated substitutions at baseline. The I50L substitution emerged on study in one of these failure isolates and was associated with a 17-fold decrease in atazanavir susceptibility from baseline and the other failure isolate with baseline atazanavir resistance and PI substitutions (M46M/I and I84I/V) had additional IAS-defined major PI substitutions (V32I, M46I, and I84V) emerge on atazanavir treatment associated with a 3-fold decrease in atazanavir susceptibility from baseline. Five of the treatment failure isolates in the atazanavir with ritonavir arm developed phenotypic emtricitabine resistance with the emergence of either the M184I (n=1) or the M184V (n=4) substitution on therapy and none developed phenotypic tenofovir disoproxil resistance. In the lopinavir/ritonavir arm, one of the virologic failure subjects isolates had a 69-fold decrease in lopinavir susceptibility emerge on therapy with the development of PI substitutions L10V, V11I, I54V, G73S, and V82A in addition to baseline PI substitutions L10L/I, V32I, I54I/V, A71I, G73G/S, V82V/A, L89V, and L90M. Six lopinavir/ritonavir virologic failure isolates developed the M184V substitution and phenotypic emtricitabine resistance and two developed phenotypic tenofovir disoproxil resistance.

Clinical Studies of Treatment-Naive subjects Receiving atazanavir 400 mg without Ritonavir: atazanavir -resistant clinical isolates from treatment-naive subjects who experienced virologic failure on atazanavir 400 mg treatment without ritonavir often developed an I50L substitution (after an average of 50 weeks of atazanavir therapy), often in combination with an A71V substitution, but also developed one or more other PI substitutions (eg, V32I, L33F, G73S, V82A, I85V, or N88S) with or without the I50L substitution. In treatment-naive subjects, viral isolates that developed the I50L substitution, without other major PI

substitutions, showed phenotypic resistance to atazanavir but retained in cell culture susceptibility to other PIs (amprenavir, indinavir, lopinavir, nelfinavir, ritonavir, and saquinavir); however, there are no clinical data available to demonstrate the effect of the I50L substitution on the efficacy of subsequently administered PIs.

Clinical Studies of Treatment-Experienced subjects: In studies of treatment-experienced subjects treated with atazanavir or atazanavir with ritonavir, most atazanavir -resistant isolates from patients who experienced virologic failure developed substitutions that were associated with resistance to multiple PIs and displayed decreased susceptibility to multiple PIs. The most common protease substitutions to develop in the viral isolates of subjects who failed treatment with atazanavir 300 mg once daily and ritonavir 100 mg once daily (together with tenofovir DF and an NRTI) included V32I, L33F/V/I, E35D/G, M46I/L, I50L, F53L/V, I54V, 71V/T/I, G73S/T/C, V82A/T/L, I85V, and L89V/Q/M/T. Other substitutions that developed on atazanavir with ritonavir treatment including E34K/A/Q, G48V, I84V, N88S/D/T, and L90M occurred in less than 10% of subjects isolates. Generally, if multiple PI resistance substitutions were present in the HIV-1 virus of the subject at baseline, atazanavir resistance developed through substitutions associated with resistance to other PIs and could include the development of the I50L substitution. The I50L substitution has been detected in treatment-experienced subjects experiencing virologic failure after long-term treatment. Protease cleavage site changes also emerged on atazanavir treatment but their presence did not correlate with the level of atazanavir resistance.

Clinical Studies of Pediatric Subjects in AI424-397 (PRINCE I) and AI424-451 (PRINCE II): Treatment-emergent atazanavir with ritonavir resistance-associated amino acid substitution M36I in the protease was detected in the virus of one subject among treatment failures in AI424-397. In addition, three known resistance-associated substitutions for other PIs arose in the viruses from one subject each (L19I/R, H69K/R, and I72I/V). Reduced susceptibility to atazanavir, ritonavir, or atazanavir with ritonavir was not seen with these viruses. In AI424-451, atazanavir with ritonavir resistance-associated substitutions G16E, V82A/I/T, I84V, and/or L90M arose in the viruses of two subjects. The virus population harboring the M46M/V, V82V/I, I84I/V, and L90L/M substitutions acquired phenotypic resistance to ritonavir (ritonavir phenotypic fold-change of 3.5, with a ritonavircutoff of 2.5- fold change). However, these substitutions did not result in phenotypic resistance to atazanavir (atazanavir phenotypic fold-change of <1.8, with an atazanavircutoff of 2.2-fold change). Secondary PI resistance-associated amino acid substitutions also arose in the viruses of one subject each, including V11V/I, D30D/G, E35E/D, K45K/R, L63P/S, and I72I/T. Q61D and Q61E/G emerged in the viruses of two subjects who failed treatment with atazanavir with ritonavir. Viruses from nine subjects in the two studies developed NRTI resistance-associated substitutions: K65K/R (n=1), M184V (n=7), and T215I (n=1).

Cross-Resistance

Cross-resistance among PIs has been observed. Baseline phenotypic and genotypic analyses of clinical isolates from atazanavir clinical trials of PI-experienced subjects showed that isolates crossresistant to multiple PIs were cross-resistant to atazanavir. Greater than 90% of the isolates with substitutions that included I84V or G48V were resistant to atazanavir. Greater than 60% of isolates containing L90M, G73S/T/C, A71V/T, I54V, M46I/L, or a change at V82 were resistant to atazanavir, and 38% of isolates containing a D30N substitution in addition to other changes were resistant to atazanavir. Isolates resistant to atazanavir were also cross-resistant to other PIs with >90% of the isolates resistant to indinavir, lopinavir, nelfinavir, ritonavir, and saquinavir, and 80% resistant to amprenavir. In treatment-experienced subjects, PI-resistant viral isolates that developed the I50L substitution in addition to other PI resistance-associated substitution were also cross-resistant to other PIs.

Baseline Genotype/Phenotype and Virologic Outcome Analyses

Genotypic and/or phenotypic analysis of baseline virus may aid in determining atazanavir susceptibility before initiation of atazanavir with ritonavir therapy. An association between virologic response at 48 weeks and the number and type of primary PI resistance-associated substitutions detected in baseline HIV-1 isolates from antiretroviral-experienced subjects receiving atazanavir with ritonavir once daily or lopinavir / ritonavir (fixed-dose product) twice daily in Study AI424-045 is shown in Table 23. Overall, both the number and type of baseline PI substitutions affected response rates in treatment-experienced subjects. In the atazanavir with ritonavir group, subjects had lower response rates when 3 or more baseline PI substitutions, including a substitution at position 36, 71, 77, 82, or 90, were present compared to subjects with 1-2 PI substitutions, including one of these substitutions.

Table 23: HIV RNA Response by Number and Type of Baseline PI Substitution, Antiretroviral-Experienced Subjects in Study AI424- 045, As-Treated Analysis

Number and Type of Baseline PI Substitutions^a	Virologic Response = HIV RNA <400 copies/mL^b	
	atazanavir with ritonavir (n=110)	opinavir/ritonavir (n=113)
3 or more primary PI substitutions including^d:		
D30N	75% (6/8)	50% (3/6)
M36I/V	19% (3/16)	33% (6/18)
M46I/L/T	24% (4/17)	23% (5/22)

I54V/L/T/M/A	31% (5/16)	31% (5/16)
A71V/T/I/G	34% (10/29)	39% (12/31)
G73S/A/C/T	14% (1/7)	38% (3/8)
V77I	47% (7/15)	44% (7/16)
V82A/F/T/S/I	29% (6/21)	27% (7/26)
I84V/A	11% (1/9)	33% (2/6)
N88D	63% (5/8)	67% (4/6)
L90M	10% (2/21)	44% (11/25)
Number of baseline primary PI substitutions^a		
All patients, as-treated	58% (64/110)	59% (67/113)
0-2 PI substitutions	75% (50/67)	75% (50/67)
3-4 PI substitutions	41% (14/34)	43% (12/28)
5 or more PI substitutions	0% (0/9)	28% (5/18)

^a Primary substitutions include any change at D30, V32, M36, M46, I47, G48, I50, I54, A71, G73, V77, V82, I84, N88, and L90.

^b Results should be interpreted with caution because the subgroups were small.

^c Administered as a fixed-dose product.

^d There were insufficient data (n<3) for PI substitutions V32I, I47V, G48V, I50V, and F53L.

The response rates of antiretroviral-experienced subjects in Study AI424-045 were analyzed by baseline phenotype (shift in susceptibility in cell culture relative to reference, Table 24). The analyses are based on a select population with 62% of subjects receiving an NNRTI-based regimen before study entry compared to 35% receiving a PI-based regimen. Additional data are needed to determine clinically relevant break points for atazanavir.

Table 24: Baseline Phenotype by Outcome, Antiretroviral-Experienced Subjects in Study AI424-045, As-Treated Analysis

Baseline Phenotype ^a	Virologic Response = HIV-1 RNA <400 copies/mL ^b	
	Atazanavir with ritonavir (n=111)	lopinavir/ritonavir ^c (n=111)
0-2	71% (55/78)	70% (56/80)
>2-5	53% (8/15)	44% (4/9)
>5-10	13% (1/8)	33% (3/9)
>10	10% (1/10)	23% (3/13)

^a Fold change susceptibility in cell culture relative to the wild-type reference.

^b Results should be interpreted with caution because the subgroups were small.

^c Administered as a fixed-dose product.

12. NONCLINICAL TOXICOLOGY

12.1. Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis

Long-term carcinogenicity studies in mice and rats were carried out with atazanavir for two years. In the mouse study, drug-related increases in hepatocellular adenomas were found in females at 360 mg/kg/day. The systemic drug exposure (AUC) at the NOAEL (no observable adverse effect level) in females, (120 mg/kg/day) was 2.8 times and in males (80 mg/kg/day) was times higher than those in humans at the clinical dose (300 mg/day atazanavir boosted with 100 mg/day ritonavir, non-pregnant patients). In the rat study, no drug-related increases in tumor incidence were observed at doses up to 1200 mg/kg/day, for which AUCs were 1.1 (males) or (females) times those measured in humans at the clinical dose.

Mutagenesis

Atazanavir tested positive in an in vitro clastogenicity test using primary human lymphocytes, in the absence and presence of metabolic activation. Atazanavir tested negative in the in vitro Ames reverse-mutation assay, in vivo micronucleus and DNA repair tests in rats, and in vivo DNA damage test in rat duodenum (comet assay).

Impairment of Fertility

At the systemic drug exposure levels (AUC) 0.9 (in male rats) or 2.3 (in female rats) times that of the human clinical dose, (300 mg/day atazanavir boosted with 100 mg/day ritonavir) significant effects on mating, fertility, or early embryonic development were not observed.

13. CLINICAL STUDIES

13.1. Adult Patients without Prior Antiretroviral Therapy

Study AI424-138: a 96-week study comparing the antiviral efficacy and safety of either atazanavir / or lopinavir/ritonavir, each in combination with fixed-dose tenofovir DF- emtricitabine in treatment-naive subjects with HIV-1 infection . Study AI424-138 (NCT00272779) was a 96-week, open-label, randomized, multicenter study, comparing atazanavir (300 mg once daily) with ritonavir (100 mg once daily) to lopinavir with ritonavir (400/100 mg twice daily as fixed-dose product), each in combination with the fixed-dose product, tenofovir DF with emtricitabine (300/200 mg once daily), in 878 antiretroviral treatment-naive subjects. Subjects had a mean age of 36 years (range: 19-72), 49% were Caucasian, 18% Black, 9% Asian, 23% Hispanic/Mestizo/mixed race, and 68% were male. The median baseline plasma CD4+ cell count was 204 cells/mm³ (range: 2 to 810 cells/mm³) and the mean baseline plasma HIV-1

Atazanavir Teva 150/200/300mg KL SPC Notification 12-2023

RNA level was 4.94 log₁₀ copies/MI (range: 2.60 to 5.88 log₁₀ copies/mL). Treatment response and outcomes through Week 96 are presented in Table 25.

Table 25: Outcomes of Treatment Through Week 96 in Treatment-Naïve Adults (Study AI424-138)

Outcome	Atazanavir 300 mg + ritonavir 100 mg (once daily) and tenofovir DF/emtricitabine (once daily)^a (n=441) 96 Weeks	lopinavir/ritonavir^b 400 mg/100 mg (twice daily) with tenofovir DF/emtricitabine (once daily)^a (n=437) 96 Weeks
Responder ^{c,d,e}	75%	68%
Virologic failure ^f	17%	19%
Rebound	8%	10%
Never suppressed through Week 96	9%	9%
Death	1%	1%
Discontinued due to adverse event	3%	5%
Discontinued for other reasons ^g	4%	7%

^a As a fixed-dose product: 300 mg tenofovir DF/ 200 mg emtricitabine once daily.

^b As a fixed-dose product: 400 mg lopinavir/100 mg ritonavir (twice daily).

^c subjects achieved HIV-1 RNA <50 copies/mL at Week 96. Roche Amplicor®, v1.5 ultra-sensitive assay.

^d Pre-specified ITT analysis at Week 48 using as-randomized cohort: atazanavir with ritonavir 78% and lopinavir/ritonavir 76% (difference estimate: 1.7% [95% confidence interval: -3.8%, 7.1%]).

^e Pre-specified ITT analysis at Week 96 using as-randomized cohort: atazanavir with ritonavir 74% and lopinavir/ritonavir 68% (difference estimate: 6.1% [95% confidence interval: 0.3%, 12.0%]).

^f Includes viral rebound and failure to achieve confirmed HIV-1 RNA <50 copies/mL through Week 96.

^g Includes lost to follow-up, subject's withdrawal, noncompliance, protocol violation, and other reasons

Through 96 weeks of therapy, the proportion of responders among subjects with high viral loads (ie, baseline HIV-1 RNA ≥100,000 copies/mL) was comparable for the atazanavir /ritonavir (165 of 223 subjects, 74%) and lopinavir/ritonavir (148 of 222 patients, 67%) arms. At 96 weeks, the median increase from baseline in CD4+ cell count was 261 cells/mm³ for the atazanavir /ritonavir arm and 273 cells/mm³ for the lopinavir/ritonavir arm.

Study AI424-034: atazanavir once daily compared to efavirenz once daily, each in combination with fixed-dose lamivudine + zidovudine twice daily. Study AI424-034 (NCT00013897) was a randomized, double-blind, multicenter trial comparing atazanavir (400 mg once daily) to efavirenz (600 mg once daily), each in combination with the fixed-dose product of lamivudine/ zidovudine (150 mg/300 mg) given twice daily, in 810 antiretroviral treatment-naïve subjects. Subjects had a mean age of 34 years (range: 18 to 73), 36% were Hispanic, 33% were Caucasian, and 65% were male. The mean baseline CD4+ cell count was 321 cells/mm³ (range: 64 to 1424 cells/mm³) and the mean baseline plasma HIV-1 RNA level was 4.8 log₁₀

copies/mL (range: 2.2 to 5.9 log₁₀ copies/mL). Treatment response and outcomes through Week 48 are presented in Table 26.

Table 26: Outcomes of Randomized Treatment Through Week 48 in Treatment-Naive Adults (Study AI424-034)

Outcome	Atazanavir 400 mg once daily + lamivudine/ zidovudine ^d (n=405)	efavirenz 600 mg once daily + lamivudine/ zidovudine ^d (n=405)
Responder ^a	67% (32%)	62% (37%)
Virologic failure ^b	20%	21%
Rebound	17%	16%
Never suppressed through Week 48	3%	5%
Death	-	<1%
Discontinued due to adverse event	5%	7%
Discontinued for other reasons ^c	8%	10%

^a subjects achieved and maintained confirmed HIV-1 RNA <400 copies/mL (<50 copies/mL) through Week 48. Roche Amplicor® HIV-1 Monitor™ Assay, test version 1.0 or 1.5 as geographically appropriate.

^b Includes viral rebound and failure to achieve confirmed HIV-1 RNA <400 copies/mL through Week 48.

^c Includes lost to follow-up subject's withdrawal, noncompliance, protocol violation, and other reasons.

^d As a fixed-dose product 150 mg lamivudine/300 mg zidovudine twice daily.

Through 48 weeks of therapy, the proportion of responders among subjects with high viral loads (ie, baseline HIV-1 RNA ≥100,000 copies/mL) was comparable for the atazanavir and efavirenz arms. The mean increase from baseline in CD4+ cell count was 176 cells/mm³ for the atazanavir arm and 160 cells/mm³ for the efavirenz arm.

Study AI424-008: atazanavir 400 mg once daily compared to atazanavir 600 mg once daily, and compared to nelfinavir 1250 mg twice daily, each in combination with stavudine and lamivudine twice daily. Study AI424-008 (NCT identifier not available) was a 48-week, randomized, multicenter trial, blinded to dose of atazanavir, comparing atazanavir at two dose levels (400 mg and 600 mg once daily) to nelfinavir (1250 mg twice daily), each in combination with stavudine (40 mg) and lamivudine (150 mg) given twice daily, in 467 antiretroviral treatment-naive subjects. Subjects had a mean age of 35 years (range: 18 to 69), 55% were Caucasian, and 63% were male. The mean baseline

CD4+ cell count was 295 cells/mm³ (range: 4 to 1003 cells/mm³) and the mean baseline plasma HIV-1 RNA level was 4.7 log₁₀ copies/mL (range: 1.8 to 5.9 log₁₀ copies/mL). Treatment response and outcomes through Week 48 are presented in Table 27.

Table 27: Outcomes of Randomized Treatment Through Week 48 in Treatment-Naive Adults (Study AI424-008)

Outcome	Atazanavir 400 mg once daily with lamivudine + stavudine (n=181)	nelfinavir 1250 mg twice daily with lamivudine + stavudine (n=91)
Responder ^a	67% (33%)	59% (38%)
Virologic failure ^b	24%	27%
Rebound	14%	14%
Never suppressed through Week 48	10%	13%
Death	<1%	-
Discontinued due to adverse event	1%	3%
Discontinued for other reasons ^c	7%	10%

^a subjects achieved and maintained confirmed HIV-1 RNA <400 copies/mL (<50 copies/mL) through Week 48. Roche Amplicor® HIV-1 Monitor™ Assay, test version 1.0 or 1.5 as geographically appropriate.

^b Includes viral rebound and failure to achieve confirmed HIV-1 RNA <400 copies/mL through Week 48.

^c Includes lost to follow-up, subject's withdrawal, noncompliance, protocol violation, and other reasons.

Through 48 weeks of therapy, the mean increase from baseline in CD4+ cell count was 234 cells/mm³ for the atazanavir 400-mg arm and 211 cells/mm³ for the nelfinavir arm.

13.2. Adult Subjects with Prior Antiretroviral Therapy

Study AI424-045: atazanavir once daily with ritonavir once daily compared to atazanavir once daily and saquinavir (soft gelatin capsules) once daily, and compared to lopinavir / ritonavir twice daily, each in combination with tenofovir DF and one NRTI. *Study AI424-045 (NCT00035932):* was a randomized, multicenter trial comparing atazanavir (300 mg once daily) with ritonavir (100 mg once daily) to atazanavir (400 mg once daily) with saquinavir soft gelatin capsules (1200 mg once daily), and to lopinavir + ritonavir (400/100 mg twice daily as fixed-dose product), each in combination with tenofovir DF and one NRTI, in 347 (of 358 randomized) subjects who experienced virologic failure on highly active antiretroviral therapy regimens containing PIs, NNRTIs, and NRTIs. The mean time of prior exposure to antiretrovirals was 139 weeks for PIs, 85 weeks for NNRTIs, and 283 weeks for NRTIs. The mean age was 41 years (range: 24 to 74); 60% were Caucasian, and 78% were male. The mean baseline CD4+ cell count was 338 cells/mm³ (range: 14 to 1543 cells/mm³) and the mean baseline plasma HIV-1 RNA level was 4.4 log₁₀ copies/mL (range: 2.6 to 5.88 log₁₀ copies/mL).

Treatment outcomes through Week 48 for the atazanavir /ritonavir and lopinavir/ritonavir treatment arms

are presented in Table 28. atazanavir /ritonavir and lopinavir/ritonavir were similar for the primary efficacy outcome measure of time-averaged difference in change from baseline in HIV-1 RNA level. Study AI424-045 was not large enough to reach a definitive conclusion that atazanavir /ritonavir and lopinavir/ritonavir are equivalent on the secondary efficacy outcome measure of proportions below the HIV-1 RNA lower limit of quantification [see Microbiology, Tables 23 and 24 (12.4)].

Table 28: Outcomes of Treatment Through Week 48 in Study AI424-045 (Subjects with Prior Antiretroviral Experience)

Outcome	Atazanavir 300 mg + ritonavir 100 mg once daily + tenofovir DF+ 1 NRTI (n=119)	lopinavir/ritonavir (400/100 mg) twice daily + tenofovir DF + 1 NRTI (n=118)	Difference ^a (Atazanavir - lopinavir/ritonavir) ^b (CI)
HIV-1 RNA Change from Baseline (log ₁₀ copies/mL) ^c	-1.58	-1.70	+0.12 ^c (-0.17, 0.41)
CD4+ Change from Baseline (cells/mm ³) ^c	116	123	-7 (-67, 52)
Percent of Patients Responding ^e			
HIV-1 RNA <400 copies/mL ^c	55%	57%	-2.2% (-14.8%, 10.5%)
HIV RNA <50 copies/mL ^c	38%	45%	-7.1% (-19.6%, 5.4%)

^a Time-averaged difference through Week 48 for HIV-1 RNA; Week 48 difference in HIV-1 RNA percentages and CD4+ mean changes, atazanavir /ritonavir vs lopinavir/ritonavir; CI = 97.5% confidence interval for change in HIV-1 RNA; 95% confidence interval otherwise.

^b Administered as a fixed-dose product.

^c Roche Amplicor® HIV-1 Monitor™ Assay, test version 1.5.

^d Protocol-defined primary efficacy outcome measure.

^e Based on subjects with baseline and Week 48 CD4+ cell count measurements (atazanavir /ritonavir, n=85; lopinavir/ritonavir, n=93).

^f Subjects achieved and maintained confirmed HIV-1 RNA <400 copies/mL (<50 copies/mL) through Week 48.

No subjects in the atazanavir /ritonavir treatment arm and three subjects in the lopinavir/ritonavir treatment arm experienced a new-onset CDC Category C event during the study.

In Study AI424-045, the mean change from baseline in plasma HIV-1 RNA for atazanavir 400 mg with saquinavir (n=115) was -1.55 log₁₀ copies/mL, and the time-averaged difference in change in HIV-1 RNA levels versus lopinavir/ritonavir was 0.33. The corresponding mean increase in CD4+ cell count was 72 cells/mm³. Through 48 weeks of treatment, the proportion of subjects in this treatment arm with plasma

HIV-1 RNA <400 (<50) copies/mL was 38% (26%).

In this study, coadministration of atazanavir and saquinavir did not provide adequate efficacy [see Drug Interactions (7)].

Study AI424-045 also compared changes from baseline in lipid values. [See Adverse Reactions (6.1).]

Study AI424-043(NCT00028301) : Study AI424-043 was a randomized, open-label, multicenter trial comparing atazanavir (400 mg once daily) to lopinavir/ritonavir (400/100 mg twice daily as fixed-dose product), each in combination with two NRTIs, in 300 subjects who experienced virologic failure to only one prior PI-containing regimen. Through 48 weeks, the proportion of subjects with plasma HIV-1 RNA <400 (<50) copies/mL was 49% (35%) for subjects randomized to atazanavir (n=144) and 69% (53%) for subjects randomized to lopinavir/ritonavir (n=146). The mean change from baseline was -1.59 log₁₀ copies/mL in the atazanavir treatment arm and -2.02 log₁₀ copies/mL in the lopinavir/ritonavir arm. Based on the results of this study, atazanavir without ritonavir was inferior to lopinavir/ritonavir in PI-experienced subjects with prior virologic failure and is not recommended for such patients.

13.3. Pediatric Subjects

Pediatric Trials with atazanavir Capsules

Study AI424-040; PACTG 1020A (NCT00006604): Assessment of the pharmacokinetics, safety, tolerability, and virologic response of atazanavir capsules was based on data from this open-label, multicenter clinical trial which included subjects from 6 years to 21 years of age. In this study, 105 subjects (43 antiretroviral-naive and 62 antiretroviral-experienced) received once daily atazanavir capsule formulation, with or without ritonavir, in combination with two NRTIs.

One-hundred five (105) subjects (6 to less than 18 years of age) treated with the atazanavir capsule formulation, with or without ritonavir, were evaluated. Using an intent-to-treat (ITT) analysis, the overall proportions of antiretroviral-naive and -experienced subjects with HIV-1 RNA <400 copies/mL at Week 96 were 51% (22/43) and 34% (21/62), respectively. The overall proportions of antiretroviral-naive and -experienced subjects with HIV-1 RNA <50 copies/mL at Week 96 were 47% (20/43) and 24% (15/62), respectively. The median increase from baseline in absolute CD4 count at 96 weeks of therapy was 335 cells/mm³ in antiretroviral-naive subjects and 220 cells/mm³ in antiretroviral-experienced subjects.

14. HOW SUPPLIED/STORAGE AND HANDLING

Atazanavir Teva Capsules

Atazanavir Teva (atazanavir) capsules are available in the following strengths and configurations of plastic bottles with child-resistant closures or blister.

Product Strength*	Capsule Shell Color (cap/body)	Markings on Capsule (ink color)	Capsules per Bottle/blister
150 mg	Dark blue/light blue	150 (black)	Bottle- 60 Blister- 60
200 mg	blue/blue	200 (black)	Bottle- 60 Blister- 60
300 mg	red/blue	300 (black)	Bottle- 30 Blister- 30,60,90

Not all pack sizes may be marketed

Atazanavir Teva (atazanavir sulfate) Capsules should be stored below 25°C.

The expiry date of the product is indicated on the packaging materials.

After first opening the bottle:

300 mg use within 2 months

150 mg and 200 mg use within 3 months .

*150 mg atazanavir equivalent to 170.9 mg atazanavir sulfate.

200 mg atazanavir equivalent to 227.8 mg atazanavir sulfate.

300 mg atazanavir equivalent to 341.7 mg atazanavir sulfate.

REGISTRATION NUMBERS

Atazanavir Teva 150 mg: 160-95-35017

Atazanavir Teva 200 mg: 160-96-35018

Atazanavir Teva 300 mg: 160-97-35019

MANUFACTURER & LICENSE HOLDER

Teva Israel LTD

124 Dvora HaNevi'a St., Tel Aviv 6944020 Israel

this leaflet was revised in December 2023 according to MOHs guidelines