



פברואר 2024

Suliqua 100/50

Suliqua 100/33

Suliqua 100/50

Each pre-filled pen contains 300 units of insulin glargine and 150 micrograms lixisenatide in 3 ml solution.
Each ml contains 100 units of insulin glargine and 50 micrograms lixisenatide.
Each dose step contains 1 unit of insulin glargine and 0.5 micrograms of lixisenatide

Suliqua 100/33

Each pre-filled pen contains 300 units of insulin glargine and 100 micrograms Lixisenatide in 3 ml solution.
Each ml contains 100 units of insulin glargine and 33 micrograms lixisenatide.
Each dose step contains 1 unit of insulin glargine and 0.33 micrograms of lixisenatide

ההתוויה המאושרת:

Suliqua is indicated for the treatment of adults with insufficiently controlled type 2 diabetes mellitus to improve glycaemic control as an adjunct to diet and exercise in addition to metformin with or without sodium-glucose co-transporter-2 (SGLT-2) inhibitors.

העדכונים בעלון לרופא הינם:

4.2 Posology and method of administration

#####

Starting dose

Therapy with basal insulin or glucagon-like peptide-1 (GLP-1) receptor agonist or oral glucose lowering medicinal product other than metformin and SGLT-2 inhibitors should be discontinued prior to initiation of Suliqua.

The starting dose of Suliqua is based on previous anti-diabetic treatment, and in order not to exceed the recommended lixisenatide starting dose of 10 mcg:



		Previous therapy		
		Insulin naïve patients (oral anti-diabetic treatment or GLP-1 receptor agonist)	Insulin glargine (100 units/ml)** ≥20 to <30 units	Insulin glargine (100 units/ml)** ≥30 to ≤60 units
Starting dose and pen	Suliqua (10-40) pen	10 dose steps (10 units/5 mcg)*	20 dose steps (20 units/10 mcg)*	
	Suliqua (30-60) pen			30 dose steps (30 units/10 mcg)*

* Units insulin glargine (100 units/ml)/mcg lixisenatide

Patients on less than 20 Units of insulin glargine may be considered similar to insulin naïve patients.

#####

Missed dose

If a dose of Suliqua is missed, it should be injected within the hour prior to the next meal.

Special population

Elderly (≥65 years old)

Suliqua can be used in elderly patients. The dose should be adjusted on an individual basis, based on glucose monitoring. In the elderly, progressive deterioration of renal function may lead to a steady decrease in insulin requirements. For lixisenatide no dose adjustment is required based on age. The therapeutic experience of Suliqua in patients ≥75 years of age is limited.

#####

העלון המעודכן נשלח לפרסום במאגר התרופות שבאתר משרד הבריאות וניתן לקבלו מודפס על ידי פנייה לבעל הרישום - סאנופי ישראל בע"מ, Greenwork Park, מתחם העסקים בקיבוץ יקום, בניין E (קומה 1), 6097600, יקום או בטלפון: 09-8633081.

להלן הקישור לאתר משרד הבריאות:

<https://israeldrugs.health.gov.il/#/byDrug>

בברכה,

חברת סאנופי ישראל בע"מ