

SUMMARY OF PRODUCT CHARACTERISTICS

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Tylo-Vine Veterinary

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each 1.1 g of granules contains

Active substance: 1000 mg of tylosin (equivalent to 1100 mg of tylosin tartrate)

3. PHARMACEUTICAL FORM

Granules for use in drinking water.

4. CLINICAL PARTICULARS

4.1 Target species

Chickens, Turkeys

4.2 Indications for use, specifying the target species

Chickens: Treatment and prevention of chronic respiratory diseases (CRD) caused by *Mycoplasma gallisepticum* and *Mycoplasma synoviae* when the disease has been established in the flock.

Treatment and prevention of necrotic enteritis caused by *Clostridium perfringens* when the disease has been established in the flock

Turkeys: Treatment and prevention of infectious sinusitis caused by *Mycoplasma gallisepticum*. when the disease has been established in the flock

4.3 Contraindications

Do not use in animals with known hypersensitivity to tylosin, other macrolides or any of the excipients listed in section 6.1.

Do not use in cases of known resistance to tylosin or cross-resistance to other macrolides (MLS-resistance).

Do not use in animals vaccinated with tylosin-sensitive vaccines either at the same time or within 1 week previously.

Do not use in animals with hepatic disorders.

4.4 Special warnings for each target species

Due to likely variability (time, geographical) in susceptibility of bacteria to tylosin, bacteriological sampling and susceptibility testing are recommended.

Under-dosing and/or treating for an insufficient length of time are considered to promote the development of resistance in bacteria and should be avoided.

4.5 Special precautions for use

Special precautions for use in animals

Animals with acute infections may have a reduced water and feed consumption and should be treated with a suitable injectable veterinary medicinal product first.

Do not leave or dispose of water containing tylosin tartrate where it may be accessible to either animals not under treatment or wildlife.

Special precautions for the person administering the veterinary medicinal product to animals

Tylosin may induce irritation. Macrolides, such as tylosin, may also cause hypersensitivity (allergy) following injection, inhalation, ingestion or contact with skin or eye. Hypersensitivity to tylosin may lead to cross reactions to other macrolides and vice versa. Allergic reactions to these substances may occasionally be serious and therefore direct contact should be avoided.

To avoid exposure during preparation of the medicated drinking water, wear overalls, safety glasses, impervious gloves and wear either a disposable half mask respirator conforming to European Standard EN149 or a non-disposable respirator to European Standard EN140 with a filter to EN143.

Wash hands after use.

In the event of accidental skin contact, wash thoroughly with soap and water. In case of accidental eye contact, flush the eyes with plenty of clean, running water.

Do not handle the product if you are allergic to ingredients in the product.

If you develop symptoms following exposure, such as skin rash, you should seek medical advice and show the physician this warning. Swelling of the face, lips and eyes or difficulty in breathing are more serious symptoms and require urgent medical attention.

4.6 Adverse reactions (frequency and seriousness)

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Any suspected adverse events should be reported to the Ministry of Health according to the National Regulation by using an online form

<http://forms.gov.it/globaldata/getsequence/getsequence.aspx?formType=AdversEffectMedic@moh.gov.it>

4.7 Use during pregnancy, lactation or lay

Laboratory studies in mice and rats have not produced any evidence of teratogenic, foetotoxic or maternotoxic effects. No studies have been conducted in the target species. Use only according to the benefit/risk assessment by the responsible veterinarian.

4.8 Interaction with other medicinal products and other forms of interaction

Lincosamides and aminoglycoside antibiotics antagonise the activity of tylosin.

4.9 Amounts to be administered and administration route

Oral administration through the drinking water

1.1 gram of the veterinary medicinal product corresponds to 1 gram of tylosin. The dosages are as follows:

Turkeys:

75 – 100 mg tylosin per kg BW per day (corresponding to 82.5 – 110 mg of the veterinary medicinal product per kg BW) for 3 – 5 days.

Chickens:

For the treatment of chronic respiratory disease:

75 – 100 mg tylosin per kg BW per day (corresponding to 82.5 – 110 mg of the veterinary medicinal product per kg BW) for 3 – 5 days.

For the treatment of necrotic enteritis:

20 mg tylosin per kg BW per day (corresponding to 22 mg of the veterinary medicinal product) for 3 days.

For the preparation of the medicated water the body weight of the animals to be treated and their actual daily water consumption should be taken into due account. Consumption may vary depending on factors like age, state of health, breed, husbandry system. To provide the required amount of active substance in mg per litre drinking water/milk/milk-replacer the following calculation should be made:

$$\frac{\text{..... mg tylosin per kg bodyweight per day} \times \text{Average bodyweight (kg) of the animals to be treated}}{\text{Average amount of drinking water / animal (l)}} = \frac{\text{.....mg tylosin}}{\text{l of drinking water}}$$

Sufficient access to the system of water supply should be available for the animals to be treated to ensure adequate water consumption. No other source of drinking water should be available during the medication period.

Should there be no clear response to treatment within 3 days the treatment approach should be reconsidered. After the end of the medication period the water supply system should be cleaned appropriately to avoid intake of sub-therapeutic amounts of the active substance which might support development of resistance.

Medicated water should be replaced every 24 hours.

If individual animals show signs of a serious infection such as a reduced water or feed intake, then alternative treatments should be considered.

4.10 Overdose (symptoms, emergency procedures, antidotes), if necessary

There is no evidence of tylosin toxicity in rats, at dose rates of up to 1000 mg/kg by the oral route.

There is no evidence of tylosin toxicity in chickens or turkeys when administered orally at up to three times the recommended dose.

4.11 Withdrawal period(s)

Turkeys (meat and offal): 2 days

Turkey (eggs): Zero days

Chickens (meat and offal): 1 day

Chicken (eggs): Zero days

5. PHARMACOLOGICAL PROPERTIES

Pharmacotherapeutic group: macrolide antibiotic,

ATC Vet Code: QJ01FA90

5.1 Pharmacodynamic properties

Tylosin is a macrolide antibiotic produced by a strain of *Streptomyces fradiae*. It exerts its antimicrobial effect by inhibiting protein synthesis of susceptible micro-organisms.

The tylosin spectrum of activity includes amongst others Gram-positive bacteria, some Gram – negative strains such as *Pasteurella*, and *Mycoplasma spp.*

5.2 Pharmacokinetic particulars

In most species peak plasma concentrations have been attained 1 to 2 hours after administration of tylosin. Compared to plasma levels clearly higher tissue concentrations have been observed. Tylosin was extensively metabolised.

Environmental properties

Most of the residues are excreted in faeces predominantly consisting of tylosin (factor A), relomycin (factor D) and dihydrodesmycosin.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

None

6.2 Incompatibilities

In the absence of compatibility studies , this veterinary medicinal product must not be mixed with other veterinary medicinal products.

6.3 Shelf life

The expiry date of the product is indicated on the packaging materials

Shelf-life after dilution: medicated water:24 hours

6.4. Special precautions for storage

Store below 25°C.

6.5 Nature and composition of immediate packaging

1 kg Aluminum bag, 300 g Aluminum bag, 100 g HDPE vial.

Not all pack sizes may be marketed.

6.6 Special precautions for the disposal of unused veterinary medicinal product or waste materials derived from the use of such products

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

7. MARKETING AUTHORISATION HOLDER

Abic Veterinary Products Ltd. POB 489 Industrial Zone, Beit Shemesh

8. MARKETING AUTHORISATION NUMBER

082-55-91161

This leaflet format has been determined by the Ministry of Health and the content has been checked and approved in 12.2018