

SUMMARY OF PRODUCT CHARACTERISTICS

1 NAME OF THE MEDICINAL PRODUCT

Lanton 15, Lanton 30

2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Each Lanton 15 capsule contains 15 mg of lansoprazole.

Each Lanton 30 capsule contains 30 mg of lansoprazole.

Excipient(s) with known effect:

Each Lanton 15 capsule contains 110 mg of sucrose.

Each Lanton 30 capsule contains 220 mg of sucrose.

For the full list of excipients, see section 6.1.

3 PHARMACEUTICAL FORM

Capsules

Lanton 15: opaque Yellow hard gelatin capsule containing white spherical microgranules.

Lanton 30: opaque white hard gelatin capsule containing white spherical microgranules.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

- Treatment of duodenal and gastric ulcer
- Treatment of reflux oesophagitis
- Prophylaxis of reflux oesophagitis
- Eradication of *Helicobacter pylori* (*H. pylori*) concurrently given with appropriate antibiotic therapy for treatment of *H.pylori*-associated ulcers
- Treatment of NSAID-associated benign gastric and duodenal ulcers in patients requiring continued NSAID treatment
- Prophylaxis of NSAID-associated gastric ulcers and duodenal ulcers in patients at risk (see section 4.2) requiring continued therapy
- Symptomatic gastroesophageal reflux disease
- Zollinger-Ellison syndrome.

4.2 Posology and method of administration

Posology

Treatment of duodenal ulcers:

The recommended dose is 30 mg once daily for 2 weeks. In patients not fully healed within this time, the medication is continued at the same dose for another two weeks.

Treatment of gastric ulcers:

The recommended dose is 30 mg once daily for 4 weeks. The ulcer usually heals within 4 weeks, but in patients not fully healed within this time, the medication may be continued at the same dose for another 4 weeks.

Reflux oesophagitis:

The recommended dose is 30 mg once daily for 4 weeks. In patients not fully healed within this time, the treatment may be continued at the same dose for another 4 weeks.

Prophylaxis of reflux oesophagitis:

15 mg once daily. The dose may be increased up to 30 mg daily as necessary.

Eradication of *Helicobacter pylori*:

When selecting appropriate combination therapy consideration should be given to official local guidance regarding bacterial resistance, duration of treatment, (most commonly 7 days but sometimes up to 14 days), and appropriate use of antibacterial agents.

The recommended dose is 30 mg of lanton twice daily for 7 days in combination with one of the following:

- clarithromycin 250-500 mg twice daily + amoxicillin 1 g twice daily
- clarithromycin 250 mg twice daily + metronidazole 400-500 mg twice daily

H. pylori eradication rates of up to 90% are obtained when clarithromycin is combined with lanton and amoxicillin or metronidazole.

Six months after successful eradication treatment, the risk of re-infection is low and relapse is therefore unlikely.

Use of a regimen including lanton 30 mg twice daily, amoxicillin 1 g twice daily and metronidazole 400-500 mg twice daily has also been examined. Lower eradication rates were seen using this combination than in regimens involving clarithromycin. It may be suitable for those who are unable to take clarithromycin as part of an eradication therapy, when local resistance rates to metronidazole are low.

Treatment of NSAID associated benign gastric and duodenal ulcers in patients requiring continued NSAID treatment:

30 mg once daily for four weeks. In patients not fully healed the treatment may be continued for another four weeks. For patients at risk or with ulcers that are difficult to heal, a longer course of treatment and/or a higher dose should probably be used.

Prophylaxis of NSAID associated gastric and duodenal ulcers in patients at risk (such as age > 65 or history of gastric or duodenal ulcer) requiring prolonged NSAID treatment:

15 mg once daily. If the treatment fails the dose 30 mg once daily should be used.

Symptomatic gastro-oesophageal reflux disease:

The recommended dose is 15 mg or 30 mg daily. Relief of symptoms is obtained rapidly. Individual adjustment of dosage should be considered. If the symptoms are not relieved within 4 weeks with a daily dose of 30 mg, further examinations are recommended.

Zollinger-Ellison syndrome:

The recommended adult starting dose is 60 mg once daily. Doses should be adjusted to individual patient needs and should continue for as long as clinically indicated.

Special populations

Renal impairment:

There is no need for a dose adjustment in patients with impaired renal function.

Hepatic impairment:

Patients with moderate or severe liver disease should be kept under regular supervision and a 50% reduction of the daily dose is recommended (see section 4.4 and 5.2).

Elderly:

Due to reduced clearance of lanton in the elderly an adjustment of dose may be necessary based on individual requirements. A daily dose of 30 mg should not be exceeded in the elderly unless there are compelling clinical indications.

Paediatric population:

The use of lanton is not recommended in children as clinical data are limited (see also section 5.2) and juvenile animal studies have findings of currently unknown human relevance (see section 5.3), except for the indications of reflux esophagitis and Symptomatic gastro-oesophageal reflux disease. Treatment of small children below one year of age should be avoided as available data have not shown beneficial effects in the treatment of gastro-oesophageal reflux disease.

Paediatric patients 1 to 11 years of age

In clinical studies, lanton was not administered beyond 12 weeks in 1 to 11 year olds. It is not known if lanton is safe and effective if used longer than the recommended duration. Do not exceed the recommended dose and duration of use in children as outlined below.

Therapeutic indication	classification	posology
Reflux esophagitis (Erosive esophagitis)	Short-term treatment	The recommended dose is 15mg once daily for up to 12 weeks for children weighting≤30kg or 30 mg once daily for up to 12 weeks for children weighting>30kg.
Symptomatic Gastroesophageal reflux disease (s-GERD)		

Paediatric patients 12 to 17 years of age

In clinical studies, lanton was not administered beyond 8 weeks in 12- to 17-year-olds. It is not known if lanton is safe and effective if used longer than the recommended duration. Do not exceed the recommended dose and duration of use in children as outlined below.

Therapeutic indication	classification	posology
Reflux esophagitis (Erosive esophagitis)	Short-term treatment	The recommended dose is 30mg once daily for up to 8 weeks for erosive esophagitis.
Symptomatic Gastroesophageal reflux disease (s-GERD)		The recommended dose is 15mg once daily for up to 8 weeks for non-erosive GERD.

Method of administration

For optimal effect lanton should be taken once daily in the morning, except when used for *H. pylori* eradication when treatment should be twice a day, once in the morning and once in the evening. Lanton should be taken at least 30 minutes before food (see section 5.2). Capsules should be swallowed whole with water.

4.3 Contraindications

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

4.4 Special warnings and precautions for use

Gastric malignancy

In common with other anti-ulcer therapies, the possibility of malignant gastric tumour should be excluded when treating a gastric ulcer with lanton because

lanton can mask the symptoms and delay the diagnosis.

Human immunodeficiency virus (HIV) protease inhibitors

Co-administration of lanton is not recommended with HIV protease inhibitors for which absorption is dependent on acidic intragastric pH, such as atazanavir and nelfinavir, due to significant reduction in their bioavailability (see section 4.5). If coadministration of lanton with HIV protease inhibitors is unavoidable, close clinical monitoring is recommended.

Hypomagnesaemia

Severe hypomagnesaemia has been rarely reported in patients treated with PPIs like lanton for at least three months, and in most cases for a year. Serious manifestations of hypomagnesaemia such as fatigue, tetany, delirium, convulsions, dizziness and ventricular arrhythmia can occur but they may begin insidiously and be overlooked. Hypomagnesaemia may lead to hypocalcaemia and/or hypokalaemia (see section 4.8). In most affected patients, hypomagnesaemia (and hypomagnesaemia associated hypocalcaemia and/or hypokalaemia) improved after magnesium replacement and discontinuation of the PPI.

For patients expected to be on prolonged treatment or who take PPIs with digoxin or medicinal products that may cause hypomagnesaemia (e.g., diuretics), healthcare professionals should consider measuring magnesium levels before starting PPI treatment and periodically during treatment.

Interference with laboratory tests

Increased Chromogranin A (CgA) level may interfere with investigations for neuroendocrine tumours. To avoid this interference, lanton treatment should be stopped for at least 5 days before CgA measurements (see section 5.1). If CgA and gastrin levels have not returned to reference range after initial measurement, measurements should be repeated 14 days after cessation of proton pump inhibitor treatment.

Influence on vitamin B12 absorption

Daily treatment with any acid-suppressing medications over a prolonged period of time (several years) may lead to malabsorption of cyanocobalamin (vitamin B12) caused by hypo- or achlorhydria. Cyanocobalamin deficiency should be considered in patients with Zollinger-Ellison syndrome and other pathological hypersecretory conditions requiring long-term treatment, individuals with reduced body stores or risk factors for reduced vitamin B12 absorption (such as the elderly) on long-term therapy or if relevant clinical symptoms are observed.

Hepatic impairment

Lanton should be used with caution in patients with moderate and severe hepatic dysfunction (see sections 4.2 and 5.2).

Gastrointestinal infections caused by bacteria

Lanton, like all proton pump inhibitors (PPIs), might increase the counts of bacteria normally present in the gastrointestinal tract. This may increase the risk of gastrointestinal infections caused by bacteria such as *Salmonella*, *Campylobacter* and, especially in hospitalised patients, *Clostridium difficile*.

In patients suffering from gastro-duodenal ulcers, the possibility of *H.pylori* infection as an etiological factor should be considered.

If lanton is used in combination with antibiotics for eradication therapy of *H.pylori*, then the instructions for the use of these antibiotics should also be followed.

Long term treatment

Because of limited safety data for patients on maintenance treatment for longer than 1 year, regular review of the treatment and a thorough risk/benefit assessment should be regularly performed in these patients.

Gastrointestinal disorders

Very rarely cases of colitis have been reported in patients taking lanton.

Therefore, in the case of severe and/or persistent diarrhoea, discontinuation of therapy should be considered.

With the exception of patients treated for the eradication of *H. pylori* infection, if diarrhoea persists, administration of lanton should be discontinued, due to the possibility of microscopic colitis with thickening of the collagen bundle or infiltration of inflammatory cells noted in the large intestine submucosa. In majority of cases, symptoms of microscopic colitis resolve on discontinuation of lanton.

Co-administration with NSAIDs

The treatment for the prevention of peptic ulceration of patients in need of continuous NSAID treatment should be restricted to high risk patients (e.g. previous gastrointestinal bleeding, perforation or ulcer, advanced age, concomitant use of medication known to increase the likelihood of upper GI adverse events [e.g. corticosteroids or anticoagulants], the presence of a serious co-morbidity factor or the prolonged use of NSAID maximum recommended doses).

Bone fractures

Proton pump inhibitors, especially if used in high doses and over long durations (>1 year), may modestly increase the risk of hip, wrist and spine fracture, predominantly in the elderly or in presence of other recognised risk factors. Observational studies suggest that proton pump inhibitors may increase the overall risk of fracture by 10–

40%. Some of this increase may be due to other risk factors. Patients at risk of osteoporosis should receive care according to current clinical guidelines and they should have an adequate intake of vitamin D and calcium.

Severe cutaneous adverse reactions

Severe cutaneous adverse reactions, including Stevens Johnson syndrome (SJS), toxic epidermal necrolysis (TEN), and drug reaction with eosinophilia and systemic symptoms (DRESS), which can be life-threatening or fatal, have been reported in association with the use of PPIs (see section 4.8). At the time of prescription patients should be advised of the signs and symptoms and monitored closely for skin reactions. If signs and symptoms suggestive of these reactions appear, lanton should be withdrawn immediately.

Subacute cutaneous lupus erythematosus (SCLE)

Proton pump inhibitors are associated with very infrequent cases of SCLE. If lesions occur, especially in sun-exposed areas of the skin, and if accompanied by arthralgia, the patient should seek medical help promptly and the healthcare professional should consider stopping lanton. SCLE after previous treatment with a proton pump inhibitor may increase the risk of SCLE with other proton pump inhibitors (see section 4.8).

Renal impairment

Acute tubulointerstitial nephritis (TIN) has been observed in patients taking lanton and may occur at any point during lanton therapy (see section 4.8). Acute tubulointerstitial nephritis can progress to renal failure. Lanton should be discontinued in case of suspected TIN, and appropriate treatment should be promptly initiated.

Excipients

This medicine contains sucrose. Patients with rare hereditary problems of fructose intolerance, glucose-galactose malabsorption or sucrase-isomaltase insufficiency should not take this medicine.

4.5 Interaction with other medicinal products and other forms of interaction

Effects of lanton on other medicinal products

Medicinal products with pH dependent absorption:

Lanton may interfere with the absorption of drugs where gastric pH is an important determinant of oral bioavailability.

HIV Protease Inhibitors:

Co-administration of lanton is not recommended with HIV protease inhibitors for which absorption is dependent on acidic intragastric pH, such as atazanavir and nelfinavir, due to significant reduction in their bioavailability (see section 4.4).

A study has shown that co-administration of lanton (60 mg once daily) with atazanavir 400 mg to healthy volunteers resulted in a substantial reduction in atazanavir exposure (approximately 90% decrease in AUC and C_{max}).

Ketoconazole and itraconazole:

The absorption of ketoconazole and itraconazole from the gastrointestinal tract is enhanced by the presence of gastric acid. Administration of lanton may result in sub-therapeutic concentrations of ketoconazole and itraconazole and the combination should be avoided.

Digoxin:

Co-administration of lanton and digoxin may lead to increased digoxin plasma levels. The plasma levels of digoxin should therefore be monitored and the dose of digoxin adjusted if necessary when initiating and ending lanton treatment.

Medicinal products metabolised by P450 enzymes:

Lanton may increase plasma concentrations of drugs that are metabolised by CYP3A4. Caution is advised when combining lanton with medicinal products which are metabolised by this enzyme and have a narrow therapeutic window.

Warfarin:

There have been reports of increased INR and prothrombin time in patients receiving PPIs and warfarin concomitantly. Increases in INR and prothrombin time may lead to abnormal bleeding and even death. Patients treated with lanton and warfarin concomitantly may need to be monitored for increase in INR and prothrombin time.

Theophylline:

Lanton reduces the plasma concentration of theophylline, which may

decrease the expected clinical effect at the dose. Patient monitoring should be taken in co-administration of lanton with theophylline.

Tacrolimus:

Co-administration of lanton increases the plasma concentrations of tacrolimus (a CYP3A and P-gp substrate). Lanton exposure increased the mean exposure of tacrolimus by up to 81%. Monitoring of tacrolimus plasma concentrations is advised when concomitant treatment with lanton is initiated or ended.

Medicinal products transported by P-glycoprotein:

Lanton has been observed to inhibit the transport protein, P-glycoprotein (P-gp) *in vitro*. The clinical relevance of this is unknown.

Effects of other drugs on lanton

Medicinal products which inhibit CYP2C19

Fluvoxamine:

A dose reduction may be considered when combining lanton with the CYP2C19 inhibitor fluvoxamine. The plasma concentrations of lanton increase up to 4-fold.

Medicinal products which induces CYP2C19 and CYP3A4

Enzyme inducers affecting CYP2C19 and CYP3A4 such as rifampicin, and St John's wort (*Hypericum perforatum*) can markedly reduce the plasma concentrations of lanton.

Others:

Methotrexate:

Concomitant use with high-dose methotrexate may elevate and prolong serum levels of methotrexate and/or its metabolite, possibly leading to methotrexate toxicities.

Sucralfate/Antacids:

Sucralfate/Antacids may decrease the bioavailability of lanton.

Therefore lanton should be taken at least 1 hour after taking these medicinal products.

Non-steroidal anti-inflammatory medicinal products:

No clinically significant interactions of lanton with nonsteroidal anti-inflammatory drugs have been demonstrated, although no formal interactions studies have been performed.

4.6 Fertility, pregnancy and lactation

Pregnancy:

For lanton no clinical data on exposed pregnancies are available. Animal studies do not indicate direct or indirect harmful effects with respect to pregnancy, embryonal/foetal development, parturition or postnatal development.

Therefore, the use of lanton during pregnancy is not recommended.

Breast-feeding:

It is not known whether lanton is excreted in human breast milk. Animal studies have shown excretion of lanton in milk.

A decision on whether to continue/discontinue breast-feeding or to continue/discontinue therapy with lanton should be made taking into account the benefit of breastfeeding to the child and the benefit of lanton therapy to the woman.

Fertility:

No human data on the effect of lanton on fertility is available. Reproductive studies in pregnant rats and rabbits revealed no lanton-related impairment of fertility.

4.7 Effects on ability to drive and use machines

Adverse drug reactions such as dizziness, vertigo, visual disturbances and somnolence may occur (see section 4.8). Under these conditions the ability to react may be decreased.

4.8 Undesirable effects

Frequencies are defined as common ($\geq 1/100$, $< 1/10$); uncommon ($\geq 1/1,000$, $< 1/100$); rare ($\geq 1/10,000$, $< 1/1,000$); very rare ($< 1/10,000$).

	Common	Uncommon	Rare	Very Rare	Not known
Blood and lymphatic system disorders		Thrombocytopenia*, eosinophilia, leucopenia*	anaemia	Agranulocytosis*, pancytopenia*	
Immune System disorders				Anaphylactic shock*	
Metabolism and nutrition disorders				Hyponatremia*	Hypomagnesemia‡, hypocalcemia*#, Hypokalemia*#
Psychiatric disorders		depression	Insomnia, hallucination, confusion		Visual hallucinations
Nervous system disorders	Headache, dizziness		Restlessness, vertigo, paraesthesia, somnolence, tremor		
Eye disorders			Visual disturbances		
Gastrointestinal disorders	Nausea, diarrhoea, stomach ache, constipation, vomiting, flatulence, dry mouth or throat, fundic gland polyps (benign)		Glossitis, Candidiasis of the oesophagus, pancreatitis, taste disturbances	Colitis*‡, Stomatitis	
Hepatobiliary disorders	Increase in liver enzyme levels		Hepatitis, jaundice		
Skin and subcutaneous tissue disorders	Urticaria, itching, rash		Petechiae purpura, hair loss, erythema multiforme, photosensitivity	Steven-Johnson syndrome*‡, toxic epidermal necrolysis*‡	Subacute cutaneous lupus erythematosus*‡, drug reaction with eosinophilia and systemic symptom (DRESS)* ‡
Musculoskeletal and connective tissue disorders		Arthralgia, myalgia, fracture of the hip, wrist or spine ‡			
Renal and urinary disorders			Tubulointerstitial nephritis (with		

			possible progression to renal failure)		
Reproductive system and breast disorders			gynecomastia		
General disorders and administration site conditions	fatigue	oedema	Angioedema, fever, hyperhidrosis, anorexia, impotence		
Investigations				Increase in cholesterol and triglyceride levels	

* Postmarketing events

Hypocalcaemia and/or hypokalaemia may be related to the occurrence of hypomagnesaemia (see section 4.4)

‡See section 4.4

Reporting of suspected adverse reactions:

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Any suspected adverse events should be reported to the Ministry of Health according to the National Regulation by using an online form <https://sideeffects.health.gov.il>.

4.9 Overdose

The effects of overdose on lanton in humans are not known (although the acute toxicity is likely to be low) and, consequently, instruction for treatment cannot be given. However, daily doses of up to 180mg of lanton orally and up to 90mg of lanton intravenously have been administered in trials without significant undesirable effects.

Please refer to section 4.8 for possible symptoms of lanton overdose.

In the case of suspected overdose the patient should be monitored. Lanton is not significantly eliminated by haemodialysis. If necessary, gastric emptying, charcoal and symptomatic therapy is recommended.

5 PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Proton pump inhibitors, ATC code: A02BC03.

Lanton is a gastric proton pump inhibitor. It inhibits the final stage of gastric acid formation by inhibiting the activity of H⁺/K⁺ ATPase of the parietal cells in the stomach. The inhibition is dose-dependent and reversible, and the effect applies to both basal and stimulated secretion of gastric acid. Lanton is concentrated in the

parietal cells and becomes active in their acidic environment, whereupon it reacts with the sulphhydryl group of H⁺/K⁺ATPase causing inhibition of the enzyme activity.

Effect on gastric acid secretion:

Lanton is a specific inhibitor of the parietal cell proton pump. A single oral 30mg dose of lanton inhibits pentagastrin-stimulated gastric acid secretion by about 80%. After repeated daily administration for seven days, about 90% inhibition of gastric acid secretion is achieved. It has a corresponding effect on the basal secretion of gastric acid. A single oral dose of 30mg reduces basal secretion by about 70%, and the patients' symptoms are consequently relieved starting from the very first dose. After eight days of repeated administration the reduction is about 85%. A rapid relief of symptoms is obtained by 30mg daily, and most patients with duodenal ulcer recover within 2 weeks, patients with gastric ulcer and reflux oesophagitis within 4 weeks.

By reducing gastric acidity, lanton creates an environment in which appropriate antibiotics can be effective against *H. pylori*.

During treatment with antisecretory medicinal products, serum gastrin increases in response to the decreased acid secretion. Also CgA increases due to decreased gastric acidity. The increased CgA level may interfere with investigations for neuroendocrine tumours.

Available published evidence suggests that proton pump inhibitors should be discontinued between 5 days and 2 weeks prior to CgA measurements. This is to allow CgA levels that might be spuriously elevated following PPI treatment to return to reference range.

5.2 Pharmacokinetic properties

Lanton is a racemate of two active enantiomers that are biotransformed into the active form in the acidic environment of the parietal cells. As lanton is rapidly inactivated by gastric acid, it is administered orally in enteric-coated form(s) for systemic absorption.

Absorption and distribution

Lanton exhibits high (80-90%) bioavailability with a single dose. Peak plasma levels occur within 1.5 to 2.0 hours. Intake of food slows the absorption rate of lanton and reduces the bioavailability by about 50%.

The plasma protein binding is 97%.

Biotransformation and elimination

Lanton is extensively metabolised by the liver and the metabolites are excreted by both the renal and biliary route. The metabolism of lanton is mainly catalysed by the enzyme CYP2C19. The enzyme CYP3A4 also contributes to the metabolism. The plasma elimination half-life ranges from 1 to 2 hours following single or multiple doses in healthy subjects. There is no evidence of accumulation following multiple doses in healthy subjects. Sulphone, sulphide and 5-hydroxyl derivatives of lanton have been identified in plasma. These metabolites have very little or no antisecretory activity.

A study with ¹⁴C labelled lanton indicated that approximately one-third of the administered radiation was excreted in the urine and two-thirds was recovered in the faeces.

Pharmacokinetics in elderly patients

The clearance of lanton is decreased in the elderly, with elimination half-life increased approximately 50% to 100%. Peak plasma levels were not increased in the elderly.

Pharmacokinetics in paediatric patients

The evaluation of the pharmacokinetics in children aged 1 –17 years of age showed a similar exposure as compared to adults with doses of 15 mg for those below 30 kg of weight and 30 mg for those above. The investigation of a dose of 17 mg/m² body surface or 1 mg/kg body weight also resulted in comparable exposure of lanton in children aged 2-3 months up to one year of age compared to adults.

Higher exposure to lanton in comparison to adults has been seen in infants below the age of 2-3 months with doses of both 1.0 mg/kg and 0.5 mg/kg body weight given as a single dose.

Pharmacokinetics in hepatic insufficiency

The exposure of lanton is doubled in patients with mild hepatic impairment and much more increased in patients with moderate and severe hepatic impairment.

CYP2C19 poor metabolisers

CYP2C19 is subject to genetic polymorphism and 2-6 % of the population, called poor metabolisers (PMs), are homozygote for a mutant CYP2C19 allele and therefore lacks a functional CYP2C19 enzyme. The exposure of lanton is several-fold higher in PMs than in extensive metabolisers (EMs).

5.3 Preclinical safety data

Non-clinical data reveal no special hazards for humans based on conventional studies of safety pharmacology, repeated dose toxicity, toxicity to reproduction or genotoxicity.

In two rat carcinogenicity studies, lanton produced dose-related gastric ECL cell hyperplasia and ECL cell carcinoids associated with hypergastrinaemia due to inhibition of acid secretion. Intestinal metaplasia was also observed, as were Leydig cell hyperplasia and benign Leydig cell tumours. After 18 months of treatment retinal atrophy was observed. This was not seen in monkeys, dogs or mice.

In mouse carcinogenicity studies dose-related gastric ECL cell hyperplasia developed as well as liver tumours and adenoma of rete testis. The clinical relevance of these findings is unknown.

Juvenile animal studies:

In juvenile rats lanton was administered from postnatal Day 7 (age equivalent to neonatal humans) through postnatal Day 62 (age equivalent to approximately 14 years in humans).

Studies in juvenile rats (8-week study, 6-week toxicokinetic dose titration study, developmental sensitivity study) have shown an increased incidence of cardiac valve thickening. The findings reversed or trended towards reversibility after a 4-week drug free recovery period. Juvenile rats younger than postnatal Day 21 (age equivalent to approximately 2 years in humans) were more sensitive to the development of cardiac valve thickening. The safety margin to the expected human exposure is in the range of 3- to 6-fold the exposure in juvenile studies based on the AUC at the no-observed effect level (NOEL) (8-week study, 6-week toxicokinetic dose titration study) or lowest-observed-effect level (LOEL)(developmental sensitivity study).

These studies have also shown changes in male reproductive tissue (testis and epididymis).

Moreover, growth retardation has been recorded either in males or in female rats but this led to delayed femoral growth plate thickness only in males.
The relevance of these findings to paediatric patients is unknown.

6 PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Capsule contents:

Sucrose, corn starch, methacrylic acid – ethyl acrylate copolymer (Eudragit), hypromellose, mannitol, talc, macrogol 6000, titanium dioxide, polysorbate 80, N-methylglucamine, sodium lauryl sulfate.

Capsule Shells:

Lanton 15: Gelatin, titanium dioxide, quinoline yellow, water.

Lanton 30: Gelatin, titanium dioxide, water.

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

The expiry date of the product is indicated on the packaging materials.

6.4 Special precautions for storage

Store below 25°C.

6.5 Nature and contents of container

Blister packs of 28 capsules.

7 MANUFACTURER AND REGISTRATION HOLDER

Manufacturer: Laboratories Liconza S.A., Spain.

Registration holder: Rafa Laboratories LTD., P.O. Box 405, Jerusalem 9100301, Israel.

8 MARKETING AUTHORISATION NUMBER(S)

Lanton 15: 131 08 30988

Lanton 30: 129 25 30761

Revised in February 2024.