

Consult your doctor immediately if you have any of the following symptoms, which may be a sign of liver problems:

- Pain or discomfort in the upper right stomach area
- Yellowing of your skin or the white part of your eyes
- Loss of appetite
- Nausea or vomiting
- Dark, amber-colored urine

Serious allergic reactions can happen to people who are treated with Trikafta. Call your doctor or go to the emergency room right away if you have any symptoms of an allergic reaction.

Symptoms of an allergic reaction may include:

- Rash or hives
- Tightness of the chest or throat or difficulty breathing
- Swelling of the face, lips and/or tongue, difficulty swallowing
- Light-headedness or dizziness

Side effects seen with Trikafta:

Very common side effects (occur in more than one in ten users)

- headache
- upper respiratory tract infection (common cold) including stuffy and runny nose
- stomach (abdominal) pain
- diarrhea
- rash
- increase in liver enzymes

Common side effects (occur in 1-10 in 100 users)

- nasal congestion
- increased creatine phosphokinase (sign of muscle breakdown) seen in blood tests
- stuffy or runny nose
- flu
- sinusitis
- increase in blood bilirubin
- flatulence
- abdominal distension
- conjunctivitis
- pharyngitis
- respiratory tract infection
- tonsillitis
- urinary tract infection
- increase in c-reactive protein
- low blood sugar (hypoglycemia)
- dizziness
- dysmenorrhea
- acne
- eczema
- pruritus

