

SUMMARY OF PRODUCT CHARACTERISTICS

1. NAME OF THE MEDICINAL PRODUCT

PROLOL 10

PROLOL 40

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Prolol 10:

Each tablet contains Propranolol hydrochloride 10mg.

Excipients with known effect:

Each tablet contains approximately 87mg lactose monohydrate.

Each tablet contains orange lake E-110.

Prolol 40:

Each tablet contains Propranolol hydrochloride 40mg.

Excipient with known effect:

Each tablet contains approximately 156mg lactose monohydrate.

For the full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Tablets.

Prolol 10: peach coloured, round tablets, scored on one side.

Prolol 40: pink, round tablets, scored on one side.

Prolol tablets can be divided into equal doses.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

- Management of angina pectoris.
- Control of essential and renal hypertension, essential tremor, most forms of cardiac dysrhythmias.
- As an adjunct in the management of tachycardias and arrhythmias due to thyrotoxicosis and thyrotoxic crises, hypertrophic obstructive cardiomyopathy, pheochromocytoma (with an alpha-blocker).
- Long-term prophylaxis following recovery from acute myocardial infarction (treatment to be initiated by a hospital physician).
- Migraine prophylaxis.

4.2 Posology and method of administration

Since the half-life may be increased in patients with significant hepatic or renal impairment, caution must be exercised when starting treatment and selecting the initial dose.

As with any other potent beta-blocking agent, the optimum dosage of propranolol has to be determined individually for each patient.

As a guide for the correct dosage, bradycardia (usually less than 45 beats/minute) is indicative that dosage should not be further increased.

When discontinuation of chronic administration of propranolol is planned in patients with ischemic heart disease, the dosage should be reduced gradually and the patients carefully monitored.

Angina Pectoris, Essential Tremor, Migraine Prophylaxis

Adults

The recommended initial dosage is 40 mg, 2-3 times daily. This may be increased by the same amount, at weekly intervals, according to patient's response. An adequate response in the management of migraine and essential tremor is usually seen in the range of 80-160 mg/day, and in the treatment of angina in the range of 120-240 mg/day.

Children

The dosage for migraine prophylaxis in children is as follows:

Under the age of 12 years: 20 mg 2 or 3 times daily.

Over the age of 12 years: The adult dosage.

Hypertension

The recommended initial dosage is 80 mg, twice a day. This may be increased at weekly intervals, according to patient response. The usual dosage range is 160-320 mg/day.

With concurrent diuretic or other antihypertensive drugs, a further reduction of blood pressure is obtained.

Arrhythmias, Hypertrophic Obstructive Cardiomyopathy, Thyrotoxicosis, Thyrotoxic Crises:

Adults

A dosage range of 10-40 mg, 3-4 times daily, usually achieves the desired response.

Children

For these indications as well as for pheochromocytoma, dosage should be individually determined, and the following is only a guide: A dosage of 0.25-0.5 mg per kg body weight 3-4 times daily, as required.

Pheochromocytoma

Use only with an alpha-receptor blocking drug. Preoperatively, a dosage of 60 mg/day, for 3 days, is recommended. In non-operable malignant cases, administer 30 mg/day.

Post-Myocardial Infarction

Treatment should start between days 5 and 21 after myocardial infarction, with an initial dose of 40 mg four times a day for 2 or 3 days. In order to improve compliance, the total daily dosage may thereafter be given as 80 mg twice a day.

Use in the Elderly

Evidence concerning the relation between blood level and age is conflicting. Propranolol should be used to treat the elderly with caution. It is suggested that treatment should

start with the lowest dose. With regard to the elderly, the optimum dose should be individually determined according to clinical response.

Method of administration

For oral use.

Food may slow the hepatic metabolism of propranolol therapy, enhancing its bioavailability. In order to standardize conditions, it is recommended that propranolol should be taken with meals.

4.3 Contraindications

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

Propranolol must not be used if there is a history of bronchial asthma or bronchospasm. The product label states the following warning: "Do not take Propranolol if you have a history of asthma or wheezing". A similar warning appears in the patient information leaflet.

Bronchospasm can usually be reversed by beta₂ agonist bronchodilators such as salbutamol. Large doses of the beta₂ agonist bronchodilator may be required to overcome the beta blockade produced by propranolol and the dose should be titrated according to the clinical response; both intravenous and inhalational administration should be considered. The use of intravenous aminophylline and/or the use of ipratropium (given by nebuliser) may also be considered. Glucagon (1 to 2 mg given intravenously) has also been reported to produce a bronchodilator effect in asthmatic patients. Oxygen or artificial ventilation may be required in severe cases.

Propranolol as with other beta-blockers must not be used in patients with any of the following conditions: bradycardia; cardiogenic shock; hypotension; metabolic acidosis; after prolonged fasting; severe peripheral arterial circulatory disturbances; second or third degree heart block; sick sinus syndrome; untreated phaeochromocytoma; uncontrolled heart failure or Prinzmetal's angina.

Propranolol must not be used in patients prone to hypoglycaemia, i.e., patients after prolonged fasting or patients with restricted counter-regulatory reserves. Patients with restricted counter regulatory reserves may have reduced autonomic and hormonal responses to hypoglycaemia which includes glycogenolysis, gluconeogenesis and /or impaired modulation of insulin secretion. Patients at risk for an inadequate response to hypoglycaemia includes individuals with malnutrition, prolonged fasting, starvation, chronic liver disease, diabetes and concomitant use of drugs which block the full response to catecholamines.

4.4 Special warnings and precautions for use

Propranolol as with other beta-blockers:

- although contraindicated in uncontrolled heart failure (see section 4.3), may be used in patients whose signs of heart failure have been controlled. Caution must be exercised in patients whose cardiac reserve is poor.
- should not be used in combination with calcium channel blockers with negative inotropic effects (e.g. verapamil, diltiazem), as it can lead to an exaggeration of these

effects particularly in patients with impaired ventricular function and/or SA or AV conduction abnormalities. This may result in severe hypotension, bradycardia and cardiac failure. Neither the beta-blocker nor the calcium channel blocker should be administered intravenously within 48 hours of discontinuing the other.

- although contraindicated in severe peripheral arterial circulatory disturbances (see section 4.3), may also aggravate less severe peripheral arterial circulatory disturbances.
- due to its negative effect on conduction time, caution must be exercised if it is given to patients with first degree heart block.
- may block/modify the signs and symptoms of the hypoglycaemia (especially tachycardia). Propranolol occasionally causes hypoglycaemia, even in non-diabetic patients, e.g. neonates, infants, children, elderly patients, patients on haemodialysis or patients suffering from chronic liver disease and patients suffering from overdose. Severe hypoglycaemia associated with propranolol has rarely presented with seizures and/or coma in isolated patients. Caution must be exercised in the concurrent use of propranolol and hypoglycaemic therapy in diabetic patients. propranolol may prolong the hypoglycaemic response to insulin (see section 4.3).
- may mask the signs of thyrotoxicosis.
- should not be used in untreated phaeochromocytoma. However, in patients with phaeochromocytoma, an alpha-blocker may be given concomitantly.
- will reduce heart rate as a result of its pharmacological action. In the rare instances when a treated patient develops symptoms which may be attributable to a slow heart rate, the dose may be reduced.
- may cause a more severe reaction to a variety of allergens when given to patients with a history of anaphylactic reaction to such allergens. Such patients may be unresponsive to the usual doses of adrenaline used to treat the allergic reactions.

Abrupt withdrawal of beta-blockers is to be avoided. The dosage should be withdrawn gradually over a period of 7 to 14 days. Patients should be followed during withdrawal especially those with ischaemic heart disease.

When a patient is scheduled for surgery and a decision is made to discontinue beta-blocker therapy, this should be done at least 24 hours prior to the procedure. The risk/benefit of stopping beta blockade should be made for each patient.

Since the half-life may be increased in patients with significant hepatic or renal impairment, caution must be exercised when starting treatment and selecting the initial dose.

Propranolol must be used with caution in patients with decompensated cirrhosis (see section 4.2).

In patients with portal hypertension, liver function may deteriorate and hepatic encephalopathy may develop. There have been reports suggesting that treatment with propranolol may increase the risk of developing hepatic encephalopathy (see section 4.2).

Interference with laboratory tests: propranolol has been reported to interfere with the estimation of serum bilirubin by the diazo method and with the determination of catecholamines by methods using fluorescence.

Prolol tablets contain lactose. Patients with rare hereditary problems of galactose intolerance, total lactase deficiency or glucose-galactose malabsorption should not take this medicine.

Prolol 10 contains orange lake E-110, which may cause allergic reactions.

4.5 Interaction with other medicinal products and other forms of interaction

Propranolol modifies the tachycardia of hypoglycaemia. Caution must be exercised in the concurrent use of propranolol and hypoglycaemic therapy in diabetic patients. Propranolol may prolong the hypoglycaemic response to insulin (see sections 4.3 and 4.4).

Simultaneous administration of rizatriptan and propranolol can cause an increased rizatriptan AUC and C_{max} by approximately 70-80%. The increased rizatriptan exposure is presumed to be caused by inhibition of first-pass metabolism of rizatriptan through inhibition of monoamine oxidase-A. If both drugs are to be used, a rizatriptan dose of 5 mg has been recommended.

Class I anti-arrhythmic drugs (e.g. disopyramide) and amiodarone may have potentiating effect on atrial-conduction time and induce negative inotropic effect.

Digitalis glycosides in association with beta-blockers may increase atrioventricular conduction time.

Combined use of beta-blockers and calcium channel blockers with negative inotropic effects (e.g. verapamil, diltiazem) can lead to an exaggeration of these effects particularly in patients with impaired ventricular function and/or SA or AV conduction abnormalities. This may result in severe hypotension, bradycardia and cardiac failure. Neither the beta-blocker nor the calcium channel blocker should be administered intravenously within 48 hours of discontinuing the other.

Concomitant therapy with dihydropyridine calcium channel blockers, e.g. nifedipine, may increase the risk of hypotension, and cardiac failure may occur in patients with latent cardiac insufficiency.

Concomitant use of sympathomimetic agents e.g. adrenaline, may counteract the effect of beta-blockers. Caution must be exercised in the parenteral administration of preparations containing adrenaline to patients taking beta-blockers as, in rare cases, vasoconstriction, hypertension and bradycardia may result.

Administration of propranolol during infusion of lidocaine may increase the plasma concentration of lidocaine by approximately 30%. Patients already receiving propranolol tend to have higher lidocaine levels than controls. The combination should be avoided.

Concomitant use of cimetidine or hydralazine will increase plasma levels of propranolol, and concomitant use of alcohol may increase the plasma levels of propranolol.

Beta-blockers may exacerbate the rebound hypertension which can follow the withdrawal of clonidine. If the two drugs are co-administered, the beta-blocker should be withdrawn several days before discontinuing clonidine. If replacing clonidine with beta-blocker therapy, the introduction of beta-blockers should be delayed for several days after clonidine administration has stopped.

Caution must be exercised if ergotamine, dihydroergotamine or related compounds are given in combination with propranolol, since vasospastic reactions have been reported in a few patients.

Concomitant use of prostaglandin synthetase inhibiting drugs e.g., ibuprofen and indometacin, may decrease the hypotensive effects of propranolol.

Concomitant administration of propranolol and chlorpromazine may result in an increase in plasma levels of both drugs. This may lead to an enhanced antipsychotic effect for chlorpromazine and an increased antihypertensive effect for propranolol.

Caution must be exercised when using anaesthetic agents with propranolol. The anaesthetist should be informed and the choice of anaesthetic should be an agent with as little negative inotropic activity as possible. Use of beta-blockers with anaesthetic drugs may result in attenuation of the reflex tachycardia and increase the risk of hypotension. Anaesthetic agents causing myocardial depression are best avoided.

Pharmacokinetic studies have shown that the following agents may interact with propranolol due to effects on enzyme systems in the liver which metabolise propranolol and these agents: quinidine, propafenone, rifampicin, theophylline, warfarin, thioridazine and dihydropyridine calcium channel blockers such as nifedipine, nisoldipine, nicardipine, isradipine, and lacidipine. Owing to the fact that blood concentrations of either agent may be affected, dosage adjustments may be needed according to clinical judgement (see also the interaction above concerning the concomitant therapy with dihydropyridine calcium channel blockers).

4.6 Pregnancy and lactation

Pregnancy

As with all drugs, propranolol should not be given during pregnancy unless its use is essential. There is no evidence of teratogenicity with propranolol. However beta-blockers reduce placental perfusion, which may result in intra-uterine foetal death, immature and premature deliveries. In addition, adverse effects (especially hypoglycaemia and bradycardia in the neonate and bradycardia in the foetus) may occur. There is an increased risk of cardiac and pulmonary complications in the neonate in the post-natal period.

Breast-feeding

Most beta-blockers, particularly lipophilic compounds, will pass into breast milk although to a variable extent. Breast-feeding is therefore not recommended following administration of these compounds.

4.7 Effects on ability to drive and use machines

Propranolol has no or negligible influence on the ability to drive and use machines. However it should be taken into account that occasionally dizziness or fatigue may occur.

4.8 Undesirable effects

Propranolol is usually well tolerated. In clinical studies the undesired events reported are usually attributable to the pharmacological actions of propranolol.

The following undesired events, listed by body system, have been reported.

The following definitions of frequencies are used:

Very common ($\geq 1/10$); common ($\geq 1/100$ to $< 1/10$); uncommon ($\geq 1/1,000$ to $< 1/100$); rare ($\geq 1/10,000$ to $< 1/1,000$); very rare ($< 1/10,000$); not known (cannot be estimated from the available data).

System Organ class	Frequency	Undesirable Effect
Blood and lymphatic system disorders	Rare	Thrombocytopaenia
Endocrine disorders	Not known	Hypoglycaemia in neonates, infants, children, elderly patients, patients on haemodialysis, patients on concomitant antidiabetic therapy, patients with prolonged fasting and patients with chronic liver disease has been reported, seizure linked to hypoglycaemia
Nervous system disorders	Common	Sleep disturbances, nightmares
	Rare	Hallucinations, psychoses, mood changes, confusion, memory loss, paraesthesia
	Very rare	Isolated reports of myasthenia gravis like syndrome or exacerbation of myasthenia gravis have been reported
Eye disorders	Rare	Dry eyes, visual disturbances
Cardiovascular disorders	Common	Bradycardia, cold extremities, Raynaud's phenomenon
	Rare	Heart failure deterioration, precipitation of heart block, postural hypotension which may be associated with syncope, exacerbation of intermittent claudication
Respiratory, thoracic and mediastinal disorders	Rare	Bronchospasm may occur in patients with bronchial asthma or a history of asthmatic complaints, sometimes with fatal outcome
Gastrointestinal disorders	Uncommon	Gastrointestinal disturbance, such as nausea, vomiting, diarrhoea
Skin and subcutaneous tissue disorders	Rare	Purpura, alopecia, psoriasiform skin reactions, exacerbation of psoriasis, skin rashes
General disorders and administration site conditions	Common	Fatigue and/or lassitude (often transient)
	Rare	Dizziness

Investigations	Very rare	An increase in ANA (Antinuclear Antibodies) has been observed, however the clinical relevance of this is not clear
----------------	-----------	--

Discontinuance of the drug should be considered if, according to clinical judgement, the well-being of the patient is adversely affected by any of the above reactions. Cessation of therapy with a beta-blocker should be gradual. In the rare event of intolerance, manifested as bradycardia and hypotension, the drug should be withdrawn and, if necessary, treatment for overdose instituted.

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Any suspected adverse events should be reported to the Ministry of Health according to the National Regulation by using an online form:

<https://sideeffects.health.gov.il>

4.9 Overdose

Propranolol is known to cause severe toxicity when used in overdose. Patients should be informed of the signs of overdose and advised to seek urgent medical assistance if an overdose of propranolol has been taken.

Clinical features:

Cardiac

Bradycardia, hypotension, pulmonary oedema, syncope and cardiogenic shock may develop. QRS complex prolongation, ventricular tachycardia, first to third degree AV block, ventricular fibrillation or asystole may also occur. Development of cardiovascular complications is more likely if other cardioactive drugs, especially calcium channel blockers, digoxin, cyclic antidepressants or neuroleptics have also been ingested. Older patients and those with underlying ischaemic heart disease are at risk of developing severe cardiovascular compromise.

CNS

Drowsiness, confusion, seizures, hallucinations, dilated pupils and in severe cases coma may occur. Neurological signs such as coma or absence of pupil reactivity are unreliable prognostic indicators during resuscitation.

Other features

Bronchospasm, hyperkalaemia and occasionally CNS-mediated respiratory depression may occur.

Management

In cases of overdose or extreme falls in heart rate or blood pressure, treatment with propranolol must be stopped. Management should include general symptomatic and supportive measures including a clear airway and monitoring of vital signs until stable. In symptomatic patients, or patients with an abnormal ECG, early discussion with critical care should be considered.

Consult national clinical guidance for further information on the management of overdose.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Beta blocking agents, non-selective, ATC code: C07AA05

Propranolol is a competitive antagonist at both the beta₁- and beta₂- adrenoceptors. It has no agonist activity at the beta-adrenoceptor, but has membrane stabilising activity at concentrations exceeding 1-3 mg/litre, though such concentrations are rarely achieved during oral therapy. Competitive beta-blockade has been demonstrated in man by a parallel shift to the right in the dose-heart rate response curve to beta-agonists such as isoprenaline.

Propranolol as with other beta-blockers, has negative inotropic effects, and is therefore contraindicated in uncontrolled heart failure.

Propranolol is a racemic mixture and the active form is the S (-) isomer of propranolol. With the exception of inhibition of the conversion of thyroxine to triiodothyronine, it is unlikely that any additional ancillary properties possessed by R (+) propranolol, in comparison with the racemic mixture, will give rise to different therapeutic effects.

Propranolol is effective and well tolerated in most ethnic populations, although the response may be less in black patients.

5.2 Pharmacokinetic properties

Following intravenous administration, the plasma half-life of propranolol is about 2 hours and the ratio of metabolites to parent drug in the blood is lower than after oral administration. In particular 4-hydroxypropranolol is not present after intravenous administration. Propranolol is completely absorbed after oral administration and peak plasma concentrations occur 1 to 2 hours after dosing in fasting patients. The liver removes up to 90% of an oral dose with an elimination half-life of 3 to 6 hours. Propranolol is widely and rapidly distributed throughout the body with highest levels occurring in the lungs, liver, kidney, brain and heart. Propranolol is highly protein bound (80 to 95%).

5.3 Preclinical safety data

Propranolol is a drug on which extensive clinical experience has been obtained. All relevant information for the prescriber is provided elsewhere in the Summary of Product Characteristics.

6 PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Prolol 10

Lactose monohydrate, cellulose microcrystalline, sodium starch glycolate, magnesium stearate, carmellose sodium, silica colloidal anhydrous, orange lake E-110

Prolol 40

Lactose monohydrate, sodium starch glycolate, cellulose microcrystalline, magnesium stearate, carmellose sodium, silica colloidal anhydrous, red lake E-127, blue lake E-132

6.2 Incompatibilities

None known.

6.3 Shelf life

The expiry date of the product is indicated on the packaging materials.

6.4 Special precautions for storage

Store below 25°C.

6.5 Nature and contents of container

Blister

Pack sizes: 10, 28, 30, 48, 50, 56 or 1000 tablets.

Not all pack sizes may be marketed.

6.6 Special precautions for disposal and other handling

Not applicable.

7 MARKETING AUTHORISATION HOLDER

Dexcel Ltd. 1 Dexcel Street, Or Akiva 3060000, Israel

8 MARKETING AUTHORISATION NUMBER

Prolol 10: 049-80-23645-00

Prolol 40: 048-37-23646-00

Revised in July 2024 according to MOH guidelines