



הבדון:

Augmentin 250 mg/5 ml Suspension

Powder for oral suspension

Augmentin 400 mg/5 ml Suspension

Powder for oral suspension

Augmentin ES

Powder for oral suspension

Augmentin 250 mg Tablets

Film-coated tablets

Augmentin 500 mg Tablets

Film-coated tablets

Augmentin 875 mg Tablets

Film-coated tablets

אוגמנטין 250 מ"ג/5 מיל' טרחיף

אבוקה להכנת טרחיף לשתייה

אוגמנטין 400 מ"ג/5 מיל' טרחיף

אבוקה להכנת טרחיף לשתייה

אוגמנטין ES

אבוקה להכנת טרחיף לשתייה

אוגמנטין 250 מ"ג טבליות

טבליות מצופות

אוגמנטין 500 מ"ג טבליות

טבליות מצופות

אוגמנטין 875 מ"ג טבליות

טבליות מצופות

רופא/ה נכבד/ה,

רוקחת/ת נכבד/ה,

חברת גלקסוסמייטקלין ישראל בע"מ (GSK) מבקשת להודיע על עדכון עלוני לרופא ועלוני לצרכן של התכשירים שבנדון.
בהודעה זו מצוינים העדכונים המהותיים בלבד.

מרכיבים פעילים וחזקם:

	Augmentin ES	Augmentin 250 mg/5 ml Suspension	Augmentin 400 mg/5 ml Suspension	Augmentin 250 mg Tablets	Augmentin 500 mg Tablets	Augmentin 875 mg Tablets
Amoxicillin (as trihydrate)	600 mg/ 5 ml	250 mg/ 5 ml	400 mg/ 5 ml	250 mg/ tablet	500 mg/ tablet	875 mg/ tablet
Clavulanic acid (as potassium salt)	42.9 mg/ 5 ml	62.5 mg/ 5 ml	57 mg/ 5 ml	125 mg/ tablet	125 mg/ tablet	125 mg/ tablet

התוויה הרשומה לתכשיר בישראל:

Augmentin 250 mg/5 ml Suspension	Augmentin 400 mg/5 ml Suspension	Augmentin 250 mg Tablets	Augmentin 500 mg Tablets	Augmentin 875 mg Tablets
Augmentin is indicated for the treatment of the following infections in adults and children: <ul style="list-style-type: none"> • Acute bacterial sinusitis (adequately diagnosed) • Acute otitis media • Acute exacerbations of chronic bronchitis (adequately diagnosed) • Community acquired pneumonia • Cystitis • Pyelonephritis • Skin and soft tissue infections in particular cellulitis, animal bites, severe dental abscess with spreading cellulitis • Bone and joint infections, in particular osteomyelitis. 	Augmentin is indicated for the treatment of the following infections in adults and children: <ul style="list-style-type: none"> • Acute bacterial sinusitis (adequately diagnosed) • Acute otitis media • Acute exacerbations of chronic bronchitis (adequately diagnosed) • Community acquired pneumonia • Cystitis • Pyelonephritis • Skin and soft tissue infections in particular cellulitis, animal bites, severe dental abscess with spreading cellulitis. • Bone and joint infections, in particular osteomyelitis. 	Augmentin is indicated for the treatment of the following infections in adults and children: <ul style="list-style-type: none"> • Acute bacterial sinusitis (adequately diagnosed) • Acute otitis media • Acute exacerbations of chronic bronchitis (adequately diagnosed) • Community acquired pneumonia • Cystitis • Pyelonephritis • Skin and soft tissue infections in particular cellulitis, animal bites, severe dental abscess with spreading cellulitis. • Bone and joint infections, in particular osteomyelitis. 	Augmentin is indicated for the treatment of the following infections in adults and children: <ul style="list-style-type: none"> • Acute bacterial sinusitis (adequately diagnosed) • Acute otitis media • Acute exacerbations of chronic bronchitis (adequately diagnosed) • Community acquired pneumonia • Cystitis • Pyelonephritis • Skin and soft tissue infections in particular cellulitis, animal bites, severe dental abscess with spreading cellulitis. • Bone and joint infections, in particular osteomyelitis. 	Augmentin is indicated for the treatment of the following infections in adults and children: <ul style="list-style-type: none"> • Acute bacterial sinusitis (adequately diagnosed) • Acute otitis media • Acute exacerbations of chronic bronchitis (adequately diagnosed) • Community acquired pneumonia • Cystitis • Pyelonephritis • Skin and soft tissue infections in particular cellulitis, animal bites, severe dental abscess with spreading cellulitis. • Bone and joint infections, in particular osteomyelitis.

4.8 Undesirable effects

[...]

Undesirable effects	Frequency
[...]	
Skin and subcutaneous tissue disorders ⁶	
[...]	
Symmetrical drug-related intertriginous and flexural exanthema (SDRIFE) (baboon syndrome)	Not known
[...]	

5.1 Pharmacodynamic properties

[...]

Susceptibility testing breakpoints

MIC (minimum inhibitory concentration) interpretive criteria for susceptibility testing have been established by the European Committee on Antimicrobial Susceptibility Testing (EUCAST) for amoxicillin/clavulanic acid and are listed here: https://www.ema.europa.eu/documents/other/minimum-inhibitory-concentration-mic-breakpoints_en.xlsx

Breakpoints

MIC breakpoints for amoxicillin/clavulanic acid are those of the European Committee on Antimicrobial Susceptibility Testing (EUCAST). (...)

Organism	Susceptibility Breakpoints ($\mu\text{g/ml}$)	
	Susceptible	Resistant
<i>Haemophilus influenzae</i>	$\leq 0.001^+$	$>2^+$
<i>Moraxella catarrhalis</i>	$\leq 1^+$	$>1^+$
<i>Staphylococcus spp.</i>	Note ^{2a,3a,3b,4}	Note ^{2a,3a,3b,4}
<i>Enterococcus spp.</i> ⁷	$\leq 4^{1,5}$	$>8^{1,5}$
Streptococcus groups A, B, C, G ^{2b,8} (indications other than meningitis)	Note ^{2b}	Note ^{2b}
<i>Streptococcus pneumoniae</i> ⁸	$\leq 0.5^{1,6}$	$>1^{1,6}$
Enterobacterales in uncomplicated UTIs	$\leq 32^+$	$>32^+$
Gram-negative Anaerobes	$\leq 4^+$	$>8^+$
Gram-positive Anaerobes (except <i>Clostridioides difficile</i>)	$\leq 4^+$	$>8^+$
Non-species related breakpoints	$\leq 2^+$	$>8^+$
Viridans group streptococci ⁸	Note ^{2a,9}	Note ^{2a,9}
<i>Pasteurella multocida</i>	$\leq 1^+$	$>1^+$
<i>Burkholderia pseudomallei</i>	$\leq 0.001^+$	$>8^+$
[...]		

[...]

Revised in August July 2023-2024, according to MOH guidelines.

עדכוניים מהותיים שנעשו בעליון לצריך:

4. תופעות לוואי

[...]

תופעת לוואי שכיחותן אינה ידועה (לא ניתן לדעת את שכיחות מתוך הנתונים הקיימים)

[...]

תגובה ערירות חמורות:

[...]

- פריה אדומה המופיעה בדרך כלל משני צידי הישבן, הירכיהם הפנימיות העליונות, בית השחי, ה佐ואר (אקסנטומה סימטרית

משולבת בכלפי העור הנגרמת על ידי תרופות [SDRIFE]).

[...]

• נערך באוגוסט-בולי 2024 בהתאם להנחיות משרד הבריאות.

[...]

למיידע נוסף יש לעיין בעליון לרופא ובעליון לצריך מעודכנים.
עלון לרופא ועלון לצריך נשלחו לפרסום במאג'ר התרופות שבאתר משרד הבריאות:
<https://data.health.gov.il/drugs/index.html#/byDrug>

וניתן לקבלם מודפסים על-ידי פניה לחברת גלקסוסמיתקליין רח' בזל 25 פתח תקווה טלפון: 03-9297100.