

## PHYSICIAN'S PRESCRIBING INFORMATION

### 1. NAME OF THE MEDICINAL PRODUCT

Onpattro 2 mg/mL  
concentrate for solution for infusion.

### 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each mL contains patisiran sodium equivalent to 2 mg patisiran.

Each vial contains patisiran sodium equivalent to 10 mg patisiran formulated as lipid nanoparticles.

#### Excipients with known effect

Each mL of concentrate contains 3.99 mg sodium.

For the full list of excipients, see section 6.1.

### 3. PHARMACEUTICAL FORM

Concentrate for solution for infusion (sterile concentrate).

White to off-white, opalescent, homogeneous solution (pH: 6.6-7.5).

### 4. CLINICAL PARTICULARS

#### 4.1 Therapeutic indications

Onpattro is indicated for the treatment of hereditary transthyretin-mediated amyloidosis (hATTR amyloidosis) in adult patients with stage 1 or stage 2 polyneuropathy.

#### 4.2 Posology and method of administration

Therapy should be initiated under the supervision of a physician knowledgeable in the management of amyloidosis.

#### Posology

The recommended dose of Onpattro is 300 micrograms per kg body weight administered via intravenous (IV) infusion once every 3 weeks.

Dosing is based on actual body weight. For patients weighing  $\geq 100$  kg, the maximum recommended dose is 30 mg.

Vitamin A supplementation at approximately 2 500 IU vitamin A per day is advised for patients treated with Onpattro (see section 4.4).

### Required premedication

All patients should receive premedication prior to Onpattro administration to reduce the risk of infusion-related reactions (IRRs) (see section 4.4). Each of the following medicinal products should be given on the day of Onpattro infusion at least 60 minutes prior to the start of infusion:

- Intravenous corticosteroid (dexamethasone 10 mg, or equivalent)
- Oral paracetamol (500 mg)
- Intravenous H1 blocker (diphenhydramine 50 mg, or equivalent)
- Intravenous H2 blocker (famotidine 20 mg, or equivalent)

For premedications not available or not tolerated intravenously, equivalents may be administered orally.

If clinically indicated, the corticosteroid may be tapered in decrements no greater than 2.5 mg to a minimum dose of 5 mg of dexamethasone (IV), or equivalent. The patient should receive at least 3 consecutive IV infusions of Onpattro without experiencing IRRs before each reduction in corticosteroid premedication.

Additional or higher doses of one or more of the premedications may be administered to reduce the risk of IRRs, if needed (see sections 4.4 and 4.8).

### Missed dose

If a dose is missed, Onpattro should be administered as soon as possible.

- If Onpattro is administered within 3 days of the missed dose, dosing should be continued according to the patient's original schedule.
- If Onpattro is administered more than 3 days after the missed dose, dosing should be continued every 3 weeks thereafter.

### Special populations

#### *Elderly patients*

No dose adjustment is required in patients  $\geq 65$  years of age (see section 5.2).

#### *Hepatic impairment*

No dose adjustment is necessary in patients with mild hepatic impairment (bilirubin  $\leq 1$  x upper limit of normal [ULN] and aspartate aminotransferase [AST]  $> 1$  x ULN, or bilirubin  $> 1.0$  to  $1.5$  x ULN and any AST). Onpattro has not been studied in patients with moderate or severe hepatic impairment and should not be used in these patients unless the anticipated clinical benefit outweighs the potential risk (see section 5.2).

#### *Renal impairment*

No dose adjustment is necessary in patients with mild or moderate renal impairment (estimated glomerular filtration rate [eGFR]  $\geq 30$  to  $< 90$  mL/min/1.73m<sup>2</sup>). Onpattro has not been studied in patients with severe renal impairment or end-stage renal disease and should not be used in these patients unless the anticipated clinical benefit outweighs the potential risk (see section 5.2).

#### *Paediatric population*

The safety and efficacy of Onpattro in children or adolescents  $< 18$  years of age have not been established. No data are available.

Onpattro is not indicated for children and adolescents under 18 years old

### Method of administration

Onpattro is for intravenous use.

- Onpattro must be diluted prior to intravenous infusion (see section 6.6).

- A dedicated line with an infusion set containing a 1.2 micron polyethersulfone (PES) in-line infusion filter must be used. The infusion sets and lines must be free of di(2-ethylhexyl)phthalate (DEHP).
- The diluted solution of Onpattro should be infused intravenously over approximately 80 minutes at an initial infusion rate of approximately 1 mL/min for the first 15 minutes, followed by an increase to approximately 3 mL/min for the remainder of the infusion. The duration of infusion may be extended in the event of an IRR (see section 4.4).
- Onpattro must be administered through a free-flowing venous access line. The infusion site should be monitored for possible infiltration during administration. Suspected extravasation should be managed according to local standard practice for non-vesicants.
- The patient should be observed during the infusion and, if clinically indicated, following the infusion (see section 4.4).
- After completion of the infusion, the intravenous administration set should be flushed with sodium chloride 9 mg/mL (0.9%) solution to ensure that all medicinal product has been administered.

Infusion of Onpattro at home may be considered for patients who have tolerated at least 3 infusions well in the clinic. The decision for a patient to receive home infusions should be made after evaluation and recommendation by the treating physician. Home infusions should be performed by a healthcare professional.

### **4.3 Contraindications**

Severe hypersensitivity (e.g., anaphylaxis) to the active substance or any of the excipients listed in section 6.1.

### **4.4 Special warnings and precautions for use**

#### Infusion-related reactions

IRRs have been observed in patients treated with Onpattro. In patients experiencing an IRR, the majority experienced the first IRR within the first 2 infusions (see section 4.8). Across clinical studies, the most common symptoms (reported in  $\geq 2\%$  of patients) of IRRs were flushing, back pain, nausea, abdominal pain, dyspnoea, and headache. IRRs may also include hypotension and syncope.

To reduce the risk of IRRs, patients should receive premedications on the day of Onpattro infusion, at least 60 minutes prior to the start of infusion (see section 4.2). If an IRR occurs, slowing or interrupting the infusion and institution of medical management (e.g., corticosteroids or other symptomatic treatment) should be considered, as clinically indicated. If the infusion is interrupted, resumption of the infusion at a slower infusion rate may be considered after symptoms have resolved. The infusion should be discontinued in the case of a serious or life-threatening IRR.

Some patients who experience IRRs may benefit from a slower infusion rate or additional or higher doses of one or more of the premedications with subsequent infusions to reduce the risk of IRRs.

#### Vitamin A deficiency

By reducing serum TTR protein, Onpattro treatment leads to a decrease in serum vitamin A (retinol) levels (see section 5.1). Serum vitamin A levels below the lower limit of normal should be corrected and any ocular symptoms or signs due to vitamin A deficiency should be evaluated prior to initiation of treatment.

Patients receiving Onpattro should take oral supplementation of approximately 2 500 IU vitamin A per day to reduce the potential risk of ocular toxicity due to vitamin A deficiency. Referral for ophthalmological assessment is recommended if patients develop ocular symptoms suggestive of

vitamin A deficiency, including reduced night vision or night blindness, persistent dry eyes, eye inflammation, corneal inflammation or ulceration, corneal thickening or corneal perforation.

Serum vitamin A levels should not be used to guide vitamin A supplementation during treatment with Onpattro (see section 4.5).

During the first 60 days of pregnancy, both too high or too low vitamin A levels may be associated with an increased risk of foetal malformation. Therefore, pregnancy should be excluded before initiating Onpattro and women of childbearing potential should practise effective contraception. If a woman intends to become pregnant, Onpattro and vitamin A supplementation should be discontinued, and serum vitamin A levels should be monitored and have returned to normal before conception is attempted.

In the event of an unplanned pregnancy, Onpattro should be discontinued (see section 4.6). Vitamin A supplementation should be discontinued during the first trimester, unless the pregnant woman has clinical signs of vitamin A deficiency. If such signs are present, vitamin A supplementation should not exceed 2 500 IU per day. Thereafter, vitamin A supplementation of 2 500 IU per day should be resumed in the second and third trimesters if serum vitamin A levels have not returned to normal, because of the increased risk of vitamin A deficiency in the third trimester.

#### Excipients with known effect

This medicinal product contains 3.99 mg sodium per mL, equivalent to 0.2% of the WHO recommended maximum daily intake of 2 g sodium for an adult.

#### **4.5 Interaction with other medicinal products and other forms of interaction**

No formal clinical drug interaction studies have been performed. At doses higher than the clinically relevant dose, induction and time-dependent inhibition of CYP2B6 was observed in vitro. The net effect on CYP2B6 substrates (e.g., bupropion and efavirenz) in vivo is unknown.

Onpattro is not expected to cause interactions or be affected by inhibitors or inducers of cytochrome P450 enzymes.

#### Vitamin A testing

Serum TTR is a carrier of retinol binding protein, which facilitates transport of vitamin A in the blood. Treatment with Onpattro reduces serum TTR levels, which results in reduced levels of retinol binding protein and vitamin A in the serum. However, transport and tissue uptake of vitamin A can occur through alternative mechanisms in the absence of retinol binding protein. As a result, during treatment with Onpattro, laboratory tests for serum vitamin A do not reflect the total amount of vitamin A in the body and should not be used to guide vitamin A supplementation (see sections 4.4 and 5.1).

#### **4.6 Fertility, pregnancy and lactation**

##### Women of childbearing potential

Treatment with Onpattro reduces serum levels of vitamin A. Both too high or too low vitamin A levels may be associated with an increased risk of foetal malformation. Therefore, pregnancy should be excluded before initiation of treatment and women of childbearing potential should use effective contraception. If a woman intends to become pregnant, Onpattro and vitamin A supplementation should be discontinued and serum vitamin A levels should be monitored and have returned to normal before conception is attempted.

## Pregnancy

There are no data on the use of Onpattro in pregnant women. Animal studies are insufficient with respect to reproductive toxicity (see section 5.3). Due to the potential teratogenic risk arising from unbalanced vitamin A levels, Onpattro should not be used during pregnancy, unless the clinical condition of the woman requires treatment. As a precautionary measure, vitamin A and thyroid stimulating hormone (TSH) levels should be obtained early in pregnancy (see section 5.3). Close monitoring of the foetus should be carried out in the event of an unplanned pregnancy, especially during the first trimester (see section 4.4). Women of childbearing potential have to use effective contraception during treatment with Onpattro.

## Breast-feeding

It is unknown whether Onpattro is excreted in human milk. A risk to the breastfed newborn/infant cannot be excluded. Available toxicological data in animals have shown excretion of small amounts of the lipid components DLin-MC3-DMA and PEG<sub>2000</sub>-C-DMG in milk (see section 5.3).

A decision must be made whether to discontinue breast-feeding or to discontinue/abstain from Onpattro, taking into account the benefit of breast-feeding for the child and the benefit of therapy for the woman.

## Fertility

There are no data on the effects of Onpattro on human fertility. No impact on male or female fertility was detected in animal studies (see section 5.3).

### **4.7 Effects on ability to drive and use machines**

On the basis of the pharmacodynamic and pharmacokinetic profiles, Onpattro is considered to have no or negligible influence on the ability to drive or use machines.

### **4.8 Undesirable effects**

#### Summary of the safety profile

The most frequently occurring adverse reactions reported in Onpattro-treated patients were peripheral oedema (29.7%) and infusion-related reactions (18.9%). One patient (0.7%) discontinued treatment during clinical studies due to an infusion-related reaction.

#### Tabulated list of adverse reactions

The adverse reactions are presented as MedDRA preferred terms under the MedDRA System Organ Class (SOC) by frequency. Within each frequency grouping, adverse reactions are presented in order of decreasing seriousness. The frequency of the adverse reactions is expressed according to the following categories:

- Very common ( $\geq 1/10$ )
- Common ( $\geq 1/100$  to  $< 1/10$ )
- Uncommon ( $\geq 1/1\ 000$  to  $< 1/100$ )

**Table 1: Adverse reactions reported for Onpattro 300 micrograms per kg**

System Organ Class	Adverse Reaction	Frequency
Infections and infestations	Bronchitis	Common
	Sinusitis	Common
	Rhinitis	Common
Immune system disorders	Infusion-related reaction	Very common
Ear and labyrinth disorders	Vertigo	Common
Respiratory, thoracic and mediastinal disorders	Dyspnoea	Common
Gastrointestinal disorders	Dyspepsia	Common
Skin and subcutaneous tissue disorders	Erythema	Common
Musculoskeletal and connective tissue disorders	Arthralgia	Common
	Muscle spasms	Common
General disorders and administration site conditions	Peripheral oedema	Very common
	Extravasation	Uncommon

### Description of selected adverse reactions

#### Infusion-related reactions

Symptoms of IRRs include, but are not limited to: arthralgia or pain (including back, neck, or musculoskeletal pain), flushing (including erythema of face or skin warm), nausea, abdominal pain, dyspnoea or cough, dysphonia, chest discomfort or chest pain, headache, rash, pruritus, chills, dizziness, fatigue, increased heart rate or palpitations, hypotension which may include syncope, hypertension, facial oedema.

In clinical studies, all patients received premedication with a corticosteroid, paracetamol, and H1 and H2 blockers to reduce the risk of IRRs. In the double-blind placebo-controlled study, 18.9% of Onpattro-treated patients experienced IRRs, compared to 9.1% of placebo-treated patients. In Onpattro-treated patients, all IRRs were either mild (95.2%) or moderate (4.8%) in severity. Among Onpattro-treated patients who experienced an IRR, 78.6% experienced the first IRR within the first 2 infusions. The frequency of IRRs decreased over time. Few IRRs led to infusion interruption. IRRs resulted in permanent discontinuation of Onpattro in < 1% of patients in clinical studies. For clinical management of IRRs, see section 4.4.

#### Peripheral oedema

In the placebo-controlled study, peripheral oedema was reported in 29.7% of Onpattro-treated patients and 22.1% of placebo-treated patients. All events were mild or moderate in severity and did not lead to treatment discontinuation. In Onpattro-treated patients, the events decreased in frequency over time.

#### Extravasation

Extravasation was observed in < 0.5% of infusions in clinical studies. Signs and symptoms included phlebitis or thrombophlebitis, infusion or injection site swelling, dermatitis (subcutaneous inflammation), cellulitis, erythema or injection site redness, burning sensation, or injection site pain.

### Other special population(s)

#### Liver transplant recipients

In an open-label study in 23 hATTR amyloidosis patients with polyneuropathy progression post liver transplant, the safety profile of patisiran was consistent with previous clinical studies (see section 5.1).

## Immunogenicity

Anti-drug antibodies to Onpattro were evaluated by measuring antibodies specific to PEG<sub>2000</sub>-C-DMG, a lipid component exposed on the surface of Onpattro. In the placebo-controlled and open-label clinical studies, 7 of 194 (3.6%) patients with hATTR amyloidosis developed anti-drug antibodies during treatment with Onpattro. One additional patient had pre-existing anti-drug antibodies. Anti-drug antibody titres were low and transient with no evidence of an effect on clinical efficacy, the safety profile, or the pharmacokinetic or pharmacodynamic profiles of Onpattro.

## Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product.

Any suspected adverse events should be reported to the Ministry of Health according to the National Regulation by using an online form <https://sideeffects.health.gov.il/>

## **4.9 Overdose**

In case of overdose, it is recommended that the patient be monitored for any signs or symptoms of adverse reactions and given symptomatic treatment, as appropriate.

## **5. PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: Other nervous System Drugs; ATC code: N07XX12.

#### Mechanism of action

Onpattro contains patisiran, a double-stranded small interfering ribonucleic acid (siRNA) that specifically targets a genetically conserved sequence in the 3' untranslated region of all variant and wild-type TTR mRNA. Patisiran is formulated as lipid nanoparticles to deliver the siRNA to hepatocytes, the primary source of TTR protein in the circulation. Through a natural process called RNA interference (RNAi), patisiran causes the catalytic degradation of TTR mRNA in the liver, resulting in a reduction of serum TTR protein.

#### Pharmacodynamic effects

Mean serum TTR was reduced by approximately 80% within 10 to 14 days after a single dose with 300 micrograms per kg Onpattro. With repeat dosing every 3 weeks, mean reductions of serum TTR after 9 and 18 months of treatment were 83% and 84%, respectively. Serum TTR reduction was maintained with continued dosing.

Serum TTR is a carrier of retinol binding protein, which facilitates transport of vitamin A in the blood. Mean reductions in serum retinol binding protein of 45% and serum vitamin A of 62% were observed over 18 months (see sections 4.4 and 4.5).

#### Clinical efficacy

The efficacy of Onpattro was studied in a randomised, double-blind, placebo-controlled study in 225 hATTR amyloidosis patients with a TTR mutation and symptomatic polyneuropathy. Patients were randomised 2:1 to receive 300 micrograms per kg Onpattro or placebo via intravenous infusion once every 3 weeks for 18 months. All patients received premedication with a corticosteroid, paracetamol, and H1 and H2 blockers.

In the study, 148 patients received Onpattro and 77 patients received placebo. The median patient age at baseline was 62 (range, 24 to 83) years and 74% of patients were male, 26% were female. Thirty-nine (39) different TTR mutations were represented; the most common ( $\geq 5\%$ ) were V30M (43%), A97S (9%), T60A (7%), E89Q (6%), and S50R (5%). Approximately 10% of patients had the V30M mutation and early onset of symptoms ( $< 50$  years of age). At baseline, 46% of patients had stage 1 disease (unimpaired ambulation; mostly mild sensory, motor and autonomic neuropathy in the lower limbs), and 53% had stage 2 disease (assistance with ambulation required; mostly moderate impairment progression to the lower limbs, upper limbs, and trunk). Approximately half (53%) of patients had prior treatment with tafamidis meglumine or diflunisal. Forty-nine percent (49%) and 50% of patients had a New York Heart Association (NYHA) Class of I or II, respectively. Approximately half of patients (56%) met pre-defined criteria for cardiac involvement (defined as baseline left ventricular [LV] wall thickness  $\geq 13$  mm with no history of hypertension or aortic valve disease). Patient demographics and baseline characteristics were balanced between treatment groups, except that a higher proportion of patients in the Onpattro group had a non-V30M mutation (62% vs. 48%). Ninety-three percent (93%) of Onpattro-treated and 62% of placebo-treated patients completed 18 months of the assigned treatment.

The primary efficacy endpoint was the change from baseline to 18 months in modified Neuropathy Impairment Score +7 (mNIS+7). This endpoint is a composite measure of motor, sensory, and autonomic polyneuropathy including assessments of motor strength and reflexes, quantitative sensory testing, nerve conduction studies, and postural blood pressure, with the score ranging from 0 to 304 points, where an increasing score indicates worsening impairment.

A statistically significant benefit in mNIS+7 with Onpattro relative to placebo was observed at 18 months (Table 2). Benefits relative to placebo were also observed across all mNIS+7 components. Changes were also seen at 9 months, the first post-baseline assessment in the study, where treatment with Onpattro led to a 16.0-point treatment difference, with a mean change from baseline of -2.0 points, compared to an increase of 14.0 points with placebo. In a threshold analysis of mNIS+7 (change from baseline of  $< 0$  points), 56.1% of Onpattro-treated patients versus 3.9% of placebo-treated patients experienced improvement in mNIS+7 ( $p < 0.001$ ).

Patients treated with Onpattro experienced statistically significant benefits in all secondary endpoints compared to patients who received placebo (all  $p < 0.001$ ) (Table 2).

The key secondary endpoint was the change from baseline to 18 months in Norfolk Quality of Life-Diabetic Neuropathy (QoL-DN) total score. The Norfolk QoL-DN questionnaire (patient-reported) includes domains relating to small fibre, large fibre, and autonomic nerve function, symptoms, and activities of daily living, with the total score ranging from -4 to 136, where an increasing score indicates worsening quality of life. At 18 months, a benefit with Onpattro to placebo was observed across all domains of Norfolk QoL-DN, and 51.4% of Onpattro-treated patients experienced an improvement in quality of life (Norfolk QoL-DN change from baseline of  $< 0$  points) compared to 10.4% of placebo-treated patients. Improvement was observed at 9 months, the first post-baseline assessment in the study.



**Table 2: Clinical efficacy results from the placebo-controlled study**

Endpoint <sup>a</sup>	Baseline, Mean (SD)		Change from baseline at 18 months, LS mean (SEM)		(Onpattro – Placebo) treatment difference, LS mean (95% CI)	p-value
	Onpattro N=148	Placebo N=77	Onpattro	Placebo		
<b>Primary</b>						
mNIS+7 <sup>b</sup>	80.9 (41.5)	74.6 (37.0)	-6.0 (1.7)	28.0 (2.6)	-34.0 (-39.9, -28.1)	p < 0.001
<b>Secondary</b>						
Norfolk QoL-DN <sup>b</sup>	59.6 (28.2)	55.5 (24.3)	-6.7 (1.8)	14.4 (2.7)	-21.1 (-27.2, -15.0)	p < 0.001
NIS-W <sup>b</sup>	32.7 (25.2)	29.0 (23.0)	0.05 (1.3)	17.9 (2.0)	-17.9 (-22.3, -13.4)	p < 0.001
R-ODS <sup>c</sup>	29.7 (11.5)	29.8 (10.8)	0.0 (0.6)	-8.9 (0.9)	9.0 (7.0, 10.9)	p < 0.001
10-metre walk test (m/sec) <sup>c</sup>	0.80 (0.40)	0.79 (0.32)	0.08 (0.02)	-0.24 (0.04)	0.31 (0.23, 0.39)	p < 0.001
mBMI <sup>d</sup>	970 (210)	990 (214)	-3.7 (9.6)	-119 (14.5)	116 (82, 149)	p < 0.001
COMPASS 31 <sup>b</sup>	30.6 (17.6)	30.3 (16.4)	-5.3 (1.3)	2.2 (1.9)	-7.5 (-11.9, -3.2)	p < 0.001

SD, standard deviation; LS mean, least squares mean; SEM, standard error of the mean; CI, confidence interval, NIS-W, NIS-weakness (motor strength); R-ODS, Rasch-Built Overall Disability (patient reported ability to perform activities of daily living); 10-metre walk test (gait speed); mBMI, modified body mass index (nutritional status); COMPASS 31, Composite Autonomic Symptom Score 31 (patient reported symptom score)

<sup>a</sup>All endpoints analysed using the mixed-effect model repeated measures (MMRM) method.

<sup>b</sup>A lower number indicates less impairment/fewer symptoms.

<sup>c</sup>A higher number indicates less disability/less impairment.

<sup>d</sup>mBMI: body mass index (BMI; kg/m<sup>2</sup>) multiplied by serum albumin (g/L); a higher number indicates better nutritional status; nutritional status favoured Onpattro as early as 3 months.

Patients receiving Onpattro experienced similar benefits relative to placebo in mNIS+7 and Norfolk QoL-DN score across all subgroups including age, sex, race, region, NIS score, V30M mutation status, prior tafamidis meglumine or diflunisal use, disease stage, and patients with pre-defined cardiac involvement. Patients experienced benefit across all TTR mutations and the full range of disease severity studied.

In patients with pre-defined cardiac involvement, centrally-assessed echocardiograms showed decreases in LV wall thickness (LS mean difference: -0.9 mm [95% CI -1.7, -0.2]) and longitudinal strain (LS mean difference: -1.37% [95% CI -2.48, -0.27]) with Onpattro treatment relative to placebo. N-terminal pro-B type natriuretic peptide (NT-proBNP) was 727 ng/L and 711 ng/L at baseline (geometric mean) in Onpattro-treated and placebo-treated patients, respectively. At 18 months, the adjusted geometric mean ratio to baseline was 0.89 with Onpattro and 1.97 with placebo (ratio, 0.45; p < 0.001), representing a 55% difference in favour of Onpattro.

#### Liver transplant recipients

In an open-label study, 23 patients with hATTR amyloidosis and polyneuropathy progression after receiving a liver transplant were treated with patisiran at a dose of 300 micrograms per kg via IV infusion once every 3 weeks. Median time from transplant to first patisiran dose was 9.4 years and median duration of patisiran treatment was 13.1 months. All patients received concomitant immunosuppressants. The study demonstrated a statistically significant median reduction in serum

TTR levels from baseline of 91% ( $p < 0.001$ ). Patients also showed stable or improved efficacy endpoints at Month 12 compared to baseline. This was consistent with the findings in the placebo-controlled patisiran study.

## 5.2 Pharmacokinetic properties

The pharmacokinetic properties of Onpattro were characterised by measuring the plasma concentrations of patisiran and the lipid components DLin-MC3-DMA and PEG<sub>2000</sub>-C-DMG.

### Absorption

Greater than 95% of patisiran in the circulation is associated with lipid nanoparticles. At the dose regimen of 300 micrograms per kg every 3 weeks, steady state was reached by 24 weeks of treatment. The estimated patisiran mean  $\pm$  SD steady-state peak concentration ( $C_{max}$ ), trough concentration ( $C_{trough}$ ), and area under the curve ( $AUC_{\tau}$ ) were  $7.15 \pm 2.14$   $\mu\text{g/mL}$ ,  $0.021 \pm 0.044$   $\mu\text{g/mL}$ , and  $184 \pm 159$   $\mu\text{g}\cdot\text{h/mL}$ , respectively. The accumulation of  $AUC_{\tau}$  was 3.2-fold at steady-state compared to the first dose.

The estimated DLin-MC3-DMA mean  $\pm$  SD steady-state  $C_{max}$ ,  $C_{trough}$  and  $AUC_{\tau}$  were  $40.2 \pm 11.5$   $\mu\text{g/mL}$ ,  $1.75 \pm 0.698$   $\mu\text{g/mL}$ , and  $1403 \pm 105$   $\mu\text{g}\cdot\text{h/mL}$ , respectively. The accumulation of  $AUC_{\tau}$  was 1.76-fold at steady-state compared to the first dose.

The estimated PEG<sub>2000</sub>-C-DMG mean  $\pm$  SD steady-state  $C_{max}$ ,  $C_{trough}$  and  $AUC_{\tau}$  were  $4.22 \pm 1.22$   $\mu\text{g/mL}$ ,  $0.0236 \pm 0.0093$   $\mu\text{g/mL}$ , and  $145 \pm 64.7$   $\mu\text{g}\cdot\text{h/mL}$ , respectively. There was no accumulation of  $AUC_{\tau}$  at steady-state compared to the first dose.

### Distribution

Plasma protein binding of Onpattro is low, with  $\leq 2.1\%$  binding observed in vitro with human serum albumin and human  $\alpha 1$ -acid glycoprotein. At the dose regimen of 300 micrograms per kg every 3 weeks, the mean  $\pm$  SD steady-state volume of distribution ( $V_{ss}$ ) of patisiran, DLin-MC3-DMA and PEG<sub>2000</sub>-C-DMG was  $0.26 \pm 0.20$  L/kg,  $0.47 \pm 0.24$  L/kg and  $0.13 \pm 0.05$  L/kg, respectively.

### Biotransformation

Patisiran is metabolized by nucleases to nucleotides of various lengths. DLin-MC3-DMA is primarily metabolised to 4-dimethylaminobutyric acid (DMBA) by hydrolysis. There is little to no metabolism of PEG<sub>2000</sub>-C-DMG.

### Elimination

At the dose regimen of 300 micrograms per kg every 3 weeks, mean  $\pm$  SD steady state plasma clearance ( $CL_{ss}$ ) of patisiran was  $3.0 \pm 2.5$  mL/h/kg. The mean  $\pm$  SD terminal elimination half-life ( $t_{1/2\beta}$ ) of patisiran was  $3.2 \pm 1.8$  days. Less than 1% of patisiran in the administered dose was recovered intact in urine.

The estimated DLin-MC3-DMA mean  $\pm$  SD steady-state  $CL_{ss}$  was  $2.1 \pm 0.8$  mL/h/kg. Approximately 5.5% of DLin-MC3-DMA was recovered after 96 hours as its metabolite (DMBA) in urine.

The estimated PEG<sub>2000</sub>-C-DMG mean  $\pm$  SD steady-state  $CL_{ss}$  was  $2.1 \pm 0.6$  mL/h/kg. In rats and monkeys, PEG<sub>2000</sub>-C-DMG is eliminated unchanged in the bile. PEG<sub>2000</sub>-C-DMG excretion in humans was not measured.

### Linearity/non-linearity

Exposure to patisiran and the lipid components (DLin-MC3-DMA and PEG<sub>2000</sub>-C-DMG) increased proportionally with increase in dose over the range evaluated in clinical studies (10 to 500 micrograms

per kg). Patisiran and the lipid components exhibit linear and time-independent pharmacokinetics with chronic dosing at the dose regimen of 300 micrograms per kg every 3 weeks.

#### Pharmacokinetic/pharmacodynamic relationship(s)

Increasing the dose of patisiran resulted in greater TTR reduction, with maximal reductions plateauing at patisiran exposures obtained with 300 micrograms per kg every 3 weeks dosing.

#### Interactions

The components of Onpattro are not inhibitors or inducers of cytochrome P450 enzymes or transporters, except for CYP2B6 (see section 4.5). Patisiran is not a substrate of cytochrome P450 enzymes.

#### Special populations

##### Gender and race

Clinical studies did not identify significant differences in steady state pharmacokinetic parameters or TTR reduction according to gender or race (non-Caucasian vs. Caucasian).

##### Weight

No data are available for patients weighing  $\geq 110$  kg.

##### Elderly patients

In the placebo-controlled study, 62 (41.9%) patients treated with Onpattro were  $\geq 65$  years of age and 9 (6.1%) patients were  $\geq 75$  years of age. There were no significant differences in steady state pharmacokinetic parameters or TTR reduction between patients  $< 65$  years of age and  $\geq 65$  years of age.

##### Hepatic impairment

Population pharmacokinetic and pharmacodynamic analyses indicated no impact of mild hepatic impairment (bilirubin  $\leq 1$  x ULN and AST  $> 1$  x ULN, or bilirubin  $> 1.0$  to  $1.5$  x ULN and any AST) on patisiran exposure or TTR reduction compared to patients with normal hepatic function. Onpattro has not been studied in patients with moderate or severe hepatic impairment.

##### Liver transplant

In a clinical study in hATTR amyloidosis patients who had undergone prior liver transplant, steady state pharmacokinetic parameters and TTR reduction were comparable to those observed in patients without a liver transplant.

##### Renal impairment

Population pharmacokinetic and pharmacodynamic analyses indicated no impact of mild or moderate renal impairment (eGFR  $\geq 30$  to  $< 90$  mL/min/1.73m<sup>2</sup>) on patisiran exposure or TTR reduction compared to subjects with normal renal function. Onpattro has not been studied in patients with severe renal impairment or end-stage renal disease.

### **5.3 Preclinical safety data**

#### General toxicology

Liver and spleen were the primary target organs of toxicity in both rats and monkeys. Intravenous administration of Onpattro led to increases in serum liver markers (alanine aminotransferase [ALT], AST, alkaline phosphatase [ALP], and/or total bilirubin) and histopathology findings in the liver (hepatocellular/single cell necrosis, inflammation, pigment deposition, and/or monocytic infiltration) at doses  $> 100$  micrograms per kg every 4 weeks and  $> 1.0$  mg/kg every 3 weeks in rats and monkeys, respectively. In spleen, lymphoid atrophy/necrosis and histiocytosis in the white pulp was observed in rats and hypocellularity of the red pulp was observed in monkeys.

In general, all findings observed at the end of dosing in the rat and monkey toxicity studies had either a full recovery or were observed with reduced severity at the end of the 60-90 day recovery period, indicating at least partial reversibility.

#### Genotoxicity/Carcinogenicity

Onpattro did not exhibit a genotoxic potential in vitro and in vivo and was not carcinogenic in transgenic rasH2 mice.

#### Reproductive toxicity

In rats, while there were parental decreases in serum TTR ( $\geq 90\%$ ), thyroxine ( $\geq 66\%$ ) and vitamin A ( $\geq 75\%$ ) levels using a rat specific surrogate to patisiran, no effects were found on male or female fertility, embryo-foetal development, or pre-/post-natal development.

In rabbits, Onpattro generated spontaneous abortions, reduced embryo-foetal survival, and reduced foetal body weights at maternally toxic doses  $\geq 1$  mg/kg (human equivalent dose [HED] 3.2 times the recommended human dose [RHD]). As patisiran is not pharmacologically active in rabbits, these effects are not due to reductions in TTR, thyroxine or vitamin A.

Intravenous administration of Onpattro had no effect on male reproductive assessments in sexually mature cynomolgus monkeys.

In lactating rats, patisiran was not present in milk, although small amounts of the lipid components DLin-MC3-DMA and PEG<sub>2000</sub>-C-DMG were present in milk (up to 7% of concomitant maternal plasma concentrations). There were no adverse effects on the pups.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

DLin-MC3-DMA, cholesterol, DSPC, PEG<sub>2000</sub>-C-DMG

Phosphate buffered saline:  
sodium chloride  
sodium phosphate, dibasic, heptahydrate  
potassium phosphate, monobasic, anhydrous  
water for injection

### **6.2 Incompatibilities**

This medicinal product must not be mixed with other medicinal products except those mentioned in section 6.6.

### **6.3 Shelf life**

The expiry date of the product is indicated on the packaging materials.

#### After dilution

Chemical and physical in-use stability has been demonstrated for 16 hours at room temperature (up to 30°C). From a microbiological point of view, it is recommended that the product should be used immediately. If not used immediately, in-use storage times and conditions prior to use are the

responsibility of the user and should not be longer than 16 hours at either 2°C to 8°C or room temperature (up to 30°C), including infusion time.

#### **6.4 Special precautions for storage**

Store in a refrigerator (2°C to 8°C).

Do not freeze.

If refrigeration is not available, Onpattro can be stored at room temperature up to 25°C for up to 14 days.

For storage conditions after dilution of the medicinal product, see section 6.3.

#### **6.5 Nature and contents of container**

Type I glass vial with a chlorobutyl stopper and an aluminium flip-off cap. Pack size of 1 vial containing 5 mL concentrate.

#### **6.6 Special precautions for disposal and other handling**

This medicinal product is for single-use only.

Onpattro must be diluted with sodium chloride 9 mg/mL (0.9%) solution prior to intravenous infusion. The diluted solution for infusion should be prepared by a healthcare professional using aseptic technique as follows:

- Remove Onpattro from the refrigerator. Do not shake or vortex.
- Discard vial if it has been frozen.
- Inspect visually for particulate matter and discoloration. Do not use if discoloration or foreign particles are present. Onpattro is a white to off-white, opalescent, homogeneous solution. A white to off-white coating may be observed on the inner surface of the vial, typically at the liquid-headspace interface. Product quality is not impacted by presence of the white to off-white coating.
- Calculate the required volume of Onpattro based on the recommended weight-based dosage (see section 4.2).
- Withdraw the entire contents of one or more vials into a single sterile syringe.
- Filter Onpattro through a sterile 0.45 micron polyethersulfone (PES) syringe filter into a sterile container.
- Withdraw the required volume of filtered Onpattro from the sterile container using a sterile syringe.
- Dilute the required volume of filtered Onpattro into an infusion bag containing sodium chloride 9 mg/mL (0.9%) solution for a total volume of 200 mL. Use infusion bags that are free of di(2-ethylhexyl)phthalate (DEHP).
- Gently invert the bag to mix the solution. Do not shake. Do not mix or dilute with other medicinal products.
- Discard any unused portion of Onpattro. Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

### **7. MANUFACTURER**

Alnylam Netherlands B.V.  
Antonio Vivaldistraat 150,1083 HP  
Amsterdam, 1077 ZX  
Netherlands

## **8. LICENSE HOLDER**

Medison Pharma Ltd., 10 Hashiloach, St., P.O.B 7090, Petach Tikva

Registration Number :165-33-36108

Revised in September 2024 according to MOHs guidelines

Onpattro- SPC-0924-V1