

**PATIENT LEAFLET IN ACCORDANCE  
WITH THE PHARMACISTS' REGULATIONS  
(PREPARATIONS) – 1986**

This medicine can be sold  
without a physician's prescription

## Agisten® Aloevera Cream

**Active ingredient:**  
Clotrimazole 1% w/w

Inactive ingredients and allergens in the product – see section 2 "Important information about some of the ingredients of this medicine" and section 6 "Additional information".

**Read this entire leaflet carefully before using this medicine.** This leaflet contains concise information about the product. If you have any further questions, ask the physician or pharmacist. Use this product according to the instructions in the dosage section in this leaflet.

Consult the pharmacist if you need additional information. Contact a physician if the symptoms of your illness get worse or if they do not improve after 4 weeks.

### 1. WHAT IS THE MEDICINE INTENDED FOR?

Agisten Aloevera Cream is used to treat skin and mucous tissue infections caused by various types of fungi. It is also intended for the treatment of Candida in the genitals.

**Therapeutic group:** Antifungal agents for local use of the imidazole group.

### 2. BEFORE USING THE MEDICINE

**Do not use the medicine:**

- If you are hypersensitive (allergic) to the active ingredient or to any of the other ingredients that this medicine contains, including cetostearyl alcohol (see section 2 "Important information about some of the ingredients of this medicine" and section 6 "Additional information").
- To treat fungus on your nails or on the scalp.

**Special warnings regarding the use of the product:**

Do not smoke or get close to an open fire source due to risk of severe burns. Fabric (clothing, linens, dressings and the like) that was in contact with this medicine can easily catch fire and is a serious fire hazard. Washing clothing and bedding may reduce product build-up, but not totally remove it. As with other creams, Agisten Aloevera Cream may reduce the effectiveness of contraceptives made of rubber, such as a condom or diaphragm. If you are using Agisten Aloevera Cream on the vulva or penis, alternative contraceptives should be used for at least 5 days after the use of this product. If your condition has not improved within 4 weeks, contact a physician.

### Drug Interactions:

**If you are taking, or have recently taken other medicines including non-prescription medicines and nutritional supplements, tell the physician or pharmacist.**

**Pregnancy, breastfeeding and fertility:**

### Pregnancy and breastfeeding:

If you are pregnant, breastfeeding or trying to become pregnant, consult the physician or pharmacist before using the product and follow his instructions. This product can be used during pregnancy, but only under the supervision of a physician.

This product can be used while breastfeeding, but the nipple area must be washed before breastfeeding if the medicine was applied in the nipple area.

### Fertility:

There is no data indicating fertility problems resulting from use of the product.

### Driving and using machines:

Agisten Aloevera Cream does not affect the ability to drive or use machines or has only a negligible effect.

**Important information about some of the ingredients of this medicine:**

Agisten Aloevera Cream contains cetostearyl

alcohol that can cause local reaction of the skin, such as contact dermatitis.

Agisten Aloevera Cream contains 10 mg benzyl alcohol per 1 gram of cream that can cause an allergic reaction and mild localized irritation.

### 3. HOW TO USE THE MEDICINE?

Check with the doctor or pharmacist if you are uncertain about the dosage or about how you should use this medicine.

If this product has been prescribed for you by your physician, follow any instructions your physician may have given you. If you purchased this product without a prescription, it is to be used as described in this section.

**The usual dosage is generally:** 2-3 treatments a day.

#### How to use:

- Instructions for opening – to open the tube, turn the cap over and pierce the tube seal by pressing the cap.
- For treatment of foot fungus: Before applying the cream, rinse the feet and dry well, especially between the toes.
- Apply a thin and even layer on the affected areas 2 to 3 times a day and massage the cream into the skin.
- Typically, a small amount of ½ cm will be enough to treat a palm-sized area.
- The duration of treatment of fungal skin infections depends on the type of infection. Usually, the duration of treatment is at least 2 weeks, although it may take up to 4 weeks of treatment.
- If you have foot fungus, it is advisable to also use an antifungal powder. Consult a physician or pharmacist.
- Symptoms of skin infection such as itching and pain are expected to improve within a few days of treatment, but symptoms such as redness and peeling of the skin may disappear after a longer period of time.
- If your condition has not improved within 4 weeks, contact your physician.

**Do not exceed the recommended dose.**

For external use only.

Do not put the medicine in your mouth or swallow it.

- If you accidentally swallowed the product, you should contact your physician immediately or go to the emergency room at the nearest hospital.
- Avoid contact of the product with the eyes or mouth. In case of contact with the eyes or mouth, rinse well with water and consult a physician.
- **If you have accidentally taken a higher dosage**, if you have taken an overdose or if a child or someone else has swallowed the medicine by mistake, proceed immediately to a physician or to the emergency room of a hospital and take the medicine package with you.
- **If you forget to apply the product on time**, apply immediately as soon as you have remembered and continue the rest of the treatment as usual.
- **How can you contribute to the success of the treatment?**
  - Although the affected area is itchy, avoid scratching the area. Scratching may cause damage to the skin surface and thus spread the infection.
  - Keep infected areas clean.
  - Dry the area thoroughly but avoid excessive rubbing.
  - Do not share towels, bath mats etc. with other people as they may get the infection.
  - Always wash your hands after treating the infection to prevent its spread.
- If you suffer from foot fungus:
  - Be sure to dry the skin between the toes thoroughly.
  - Wash your socks and stockings/tights thoroughly in hot water to remove any shed skin or fungal spores.
  - Change your footwear daily if possible.

**Do not take medicines in the dark! Check**

**the label and the dose each time you take a medicine. Wear glasses if you need them. If you have any further questions about the use of the medicine, ask your physician or pharmacist.**

### 4. SIDE EFFECTS

Like all medicines, the use of Agisten Aloevera Cream may cause side effects in some users. Do not be alarmed by reading the list of side effects. You may not suffer from any of them.

As with all medicines, some patients may be hypersensitive (allergic) to this cream. If you are allergic, a reaction will develop soon after using this product. **Discontinue use and contact a physician or proceed to a hospital emergency room immediately if an allergic reaction develops.**

Signs of allergic reaction may include:

- Rash
- Swallowing or breathing problems
- Swelling of the lips, face, throat or tongue
- Weakness, dizziness or fainting
- Nausea

**After using the medicine you may experience:** itching, rash, blisters, burning, discomfort, swelling, irritation, pins and needles, redness or peeling of the skin.

**If a side effect appears, if one of the side effects gets worse, or if you suffer from a side effect which is not mentioned in this leaflet, consult the physician.**

Side effects can be reported to the Ministry of Health by clicking on the link "Reporting Side Effects from Drug Treatment" that can be found on the homepage of the Ministry of Health website ([www.health.gov.il](http://www.health.gov.il)) directing to the online form of adverse events reporting or by clicking on the following link: <https://sideeffects.health.gov.il>

Additionally, side effects can be reported to Padagis via the following address: [Padagis.co.il](mailto:Padagis.co.il)

### 5. HOW TO STORE THE MEDICINE?

- Avoid poisoning! This medicine and any other medicine should be kept in a safe place out of the reach and sight of children and/or infants to avoid poisoning. Do not induce vomiting without an explicit instruction from the physician.
- Do not use the medicine after the expiry date (exp. date) that appears on the package. The expiry date refers to the last day of that month.
- **Storage conditions:**
  - Store in a cool place, below 25°C.
  - After first opening, the medicine can be used for 6 months and not later than the expiry date that appears on the package.
  - Caution! Inflammable material. Do not light a cigarette or expose to fire until the medicine has fully dried.

### 6. ADDITIONAL INFORMATION

**In addition to the active ingredient, Agisten Aloevera Cream also contains:**

- Octyldodecanol, cetostearyl alcohol, cetyl esters wax, sorbitan monostearate, polysorbate 60, benzyl alcohol, aloe vera powder decolorized, purified water.
- What does the medicine look like and what is the contents of the package: Agisten Aloevera Cream is a white cream packaged in an aluminum tube that contains 20 grams.
- **Manufacturer and registration holder:** Padagis Israel Pharmaceuticals Ltd., 1 Rakefet St., Shoham.
- Revised in April 2024 according to MOH guidelines.
- Registration number of the medicine in the National Drug Registry of the Ministry of Health: 043-17-22461

Agisten® is a trademark of Padagis Israel Pharmaceuticals Ltd.

30200761

04/2024  
SH107561 PL 11