

Veterinary medicine package leaflet

This medicine is dispensed with a veterinarian's prescription only
For use in animals only

Dolpac 10 veterinary - oral tablets for dogs

Dolpac 25 veterinary - oral tablets for dogs

Composition:

Active ingredients:

Dolpac 10 veterinary:

200.28 mg of oxantel (as embonate)
49.94 mg of pyrantel (as embonate)
50.00 mg of praziquantel

Dolpac 25 veterinary:

500.70 mg of oxantel (as embonate)
124.85 mg of pyrantel (as embonate)
125.00 mg of praziquantel

Inactive ingredients:

dextrates, povidone K30, lauryl sodium sulphate, bacon flavor, crospovidone, sodium stearyl fumarate.

Indication:

Treatment of parasitic infestation with adult stages of roundworms and tapeworms in dogs

Nematodes and cestodes:

Nematodes: *Toxocara canis*, *Toxascaris leonina*, *Ancylostoma caninum*, *Uncinaria stenocephala*, *Trichuris vulpis*.

Cestodes: *Dipylidium caninum*, *Taenia* spp.

Side effects:

Vomiting and diarrhea may appear in the dog during the course of treatment. Anorexia is a common side effect of praziquantel; however, it was not observed in studies of Dolpac. If you notice any serious side effect not mentioned in this leaflet, tell your veterinarian.

You can report side effects to the Ministry of Health by following the link 'Reporting Side Effects of Drug Treatment' on the Ministry of Health home page (www.health.gov.il) which links to an online form for reporting side effects. You can also use this link:
<https://sideeffects.health.gov.il>

Dosage and directions for use:

Dosage table:

Dolpac 10	Dog weight	Number of tablets
The recommended dose is 20 mg oxantel / 5 mg pyrantel / 5 mg praziquantel per kg bodyweight, which is one tablet	3.1-5 kg	½
	5.1-10 kg	1
	10.1-20 kg	2
	20.1-30 kg	3

per 10 kg bodyweight. For oral administration.		
Dolpac 25 The recommended dose is 20 mg oxantel / 5 mg pyrantel / 5 mg praziquantel per kg bodyweight, which is one tablet per 25 kg bodyweight. For oral administration.	Dog weight	Number of tablets
	10.1-12.5 kg 12.6-25 kg 25.1-50 kg 50.1-75 kg	½ 1 2 3

Administer the required number of tablets, according to the dog's bodyweight, orally, in a single administration. Preferably, dogs should not eat before treatment. Food may be given one hour or more after treatment. The tablets can be split in half. Dogs kept together should be treated at the same time.

Special warnings:

Parasite resistance may develop following frequent, repeated use of this type of medicine.

Fleas serve as intermediate hosts for one of the common tapeworms – *Dipylidium caninum*. Tapeworm infestation may reoccur if the flea problem is left untreated.

Special precautions for use in animals:

Roundworm and hookworm infections:

In some of the treated dogs, the treatment may not fully eradicate *Ancylostoma caninum* and *Toxocara canis*, resulting in a continued risk of the dogs shedding parasite eggs. Follow-up on treatment results and examination of the feces are advisable and if parasite eggs are seen in the feces, the dog must be treated with a nematocidal product.

This product is not recommended for use in pups younger than 2 months old or weighing less than 1 kg.

In debilitated or heavily infested dogs, the product should be used only according to a benefit/risk assessment by the veterinarian.

Do not use in animals with known sensitivity to any of the ingredients of the product.

Gestation and lactation:

The safety of using this product during gestation and lactation has not been established. So its use is not recommended in gestating or lactating bitches.

Drug interactions: Do not use simultaneously with levamisole, piperazine or choline esterase inhibitors.

Overdose:

No serious side effects were observed in healthy dogs given 5 times the recommended dose for 6 consecutive weeks.

Special precautions for the person administering the product:

Some ingredients of this product may cause allergic reactions or skin irritation.

Avoid contact with the skin.

Avoid contact with this product if you have a known sensitivity to any of its ingredients.

Wash hands after administration.

If accidentally ingested, seek medical advice and show the package leaflet to the doctor.

Storage: at room temperature, below 25°C.

Keep out of children's reach. Do not use the medicine after the expiry date (exp. date) stated on the carton.

Dispose of any remaining veterinary medicine or waste obtained from using a veterinary medicine in the same manner as toxic waste; do not discard down the drain.

Registration number of the medicine in the Ministry of Health National Drug Registry:

Dolpac 25 veterinary: 142 77 92433 00

Dolpac 10 veterinary: 142 76 92430 00

Manufacturer: Vetoquinol, Magny-Vernois 70200 Lure, France

Registration holder: Eliezer Linevitz Ltd., 6 Edom St., POB 7006, Kanot Industrial Park.

Revised in July 2021 according to MOH guidelines.