

**PATIENT PACKAGE INSERT  
IN ACCORDANCE WITH THE  
PHARMACISTS' REGULATIONS  
(PREPARATIONS) – 1986**

The medicine is dispensed without a doctor's prescription

## **SALATAC GEL**

### **Composition:**

Salicylic acid 12%, Lactic acid 4%

**Inactive ingredients and allergens in the preparation:** see section 6.

**Read the leaflet carefully in its entirety before using the medicine.** This leaflet contains concise information about the medicine. If you have further questions, refer to the doctor or pharmacist.

Use according to the instructions in the dosage section of this leaflet. Consult a pharmacist if you need more information. Consult a doctor if the symptoms of the illness worsen or do not improve after 12 weeks.

### **1. WHAT IS THE MEDICINE INTENDED FOR?**

The medicine is intended to remove viral warts, corns and hardenings of the skin in the form of calluses.

The active ingredients soften and dissolve the cornified skin of the wart.

**Therapeutic group:** Combination of keratolytic active ingredients.

### **2. BEFORE USING THE MEDICINE**

#### **Do not use the medicine if:**

- You are sensitive (allergic) to the active ingredients (salicylic acid, lactic acid) or any of the other ingredients contained in the medicine (see the list of inactive ingredients in section 6).
- Do not use on or near your face, armpit, breasts, bottom and genital area.
- Do not use on birthmarks, moles, hairy warts, or on any other spot.
- Do not use the medicine if you are diabetic or suffer from poor blood circulation to your hands and feet.

### **Special warning regarding use of the medicine:**

- If you are unsure whether you have a wart that is suitable for treatment with Salatac Gel, ask your doctor or pharmacist before starting treatment.
- Apply the gel carefully only to the wart, by squeezing the tube gently:
  - o Avoid contact of the gel with the healthy skin surrounding the wart, especially on young children's delicate skin, because this may cause irritation. If the gel accidentally touches healthy skin, clean with a tissue immediately and if necessary, rinse the area.
  - o Do not use an excessive amount of gel.
- Avoid contact with the eyes, nose, mouth, injured skin, cuts and scratches. If the gel accidentally enters the eyes or mouth, damage may occur. Rinse eyes or mouth with plenty of water. If rinsing one eye, be sure to avoid letting the rinsing water contact the other eye. Then seek urgent medical attention.
- Avoid spillage of the substance as this may cause damage to you, your clothing and your possessions:
  - o Do not fold the tube because this may damage it and make it leak.
- The medicine is volatile and highly flammable. Do not use the medicine near flames or ignition sources (e.g., burning cigarettes or anything else that might ignite it).
- Avoid inhaling the preparation. If you inhale a lot of the substance, it may make you feel light-headed, inhale a lot of fresh air.

### **Drug interactions:**

**If you are taking, or have recently taken, other medicines, including non-prescription medicines and nutritional supplements, tell the doctor or pharmacist.** This preparation is not known to affect, or to be affected by any other medicines.

### **Pregnancy, breast-feeding and fertility:**

This medicine can be used during pregnancy and while breast-feeding. The ingredients of the medicine have been in widespread use in this and similar preparations for many years, without reports of problems. However, safety trials have not been conducted.

### **Driving and using machinery:**

Using this preparation is not known to affect your ability to drive or use machinery.

### **3. HOW SHOULD YOU USE THE MEDICINE?**

Check with the doctor or pharmacist if you are uncertain regarding the dosage and treatment regimen of the preparation.

#### **Directions for use:**

How should you treat warts and hardened skin in the form of a callus?

1. Remove any adhesive plaster.
2. Wash the affected areas, then soak in warm water for 2 to 3 minutes.
3. Dry the area thoroughly with your own towel.
4. Apply 1 to 2 drops of the gel directly on the wart. Wait until a film forms over the wart. The film will appear within a short time after the substance dries. Take care to avoid contact of the gel with the skin surrounding the wart. It is not necessary to cover it with an adhesive plaster.

5. The following evening, carefully remove the film formed from the previous evening's treatment. Repeat the treatment.

6. Once a week, before the treatment, gently rub the treated surface with the emery board provided with the package, or with a pumice stone, and then carry on with the treatment.

**Remember:** Be sure to keep using Salatac Gel until the complete disappearance of the warts.

### **What is the frequency of treatment with Salatac Gel?**

You should use Salatac Gel regularly, once a day, generally in the evening at bedtime, until healing is complete. Removal of viral warts can take up to 12 weeks.

**Note:** Before opening the tube, hold it upright with the opening facing upwards for 30 seconds. Unscrew the cap, turn it upside down and use it to puncture the opening of the tube. Take the additional long-nozzled cap out of the box and tightly close the tube with it.

### **Do not exceed the recommended dosage.**

Attention:

- Do not swallow!
- This medicine is intended for external use only.

### **If you forget to take the medicine:**

If you forgot to take the medicine at the designated time, do not take a double dose. Take the next dose at the regular time.

**Do not take medicines in the dark! Check the label and the dose each time you take medicine. Wear glasses if you need them.**

**If you have further questions regarding use of the medicine, consult a doctor or pharmacist.**

### **4. SIDE EFFECTS**

As with any medicine, use of Salatac Gel may cause side effects in some users. Do not be alarmed when reading the list of side effects. You may not suffer from any of them.

While using Salatac Gel you may feel a slight tingling sensation and/or some mild tenderness at the treated area. This sensation is usually temporary, and in rare cases may appear as a temporary blemish on the skin.

If the gel was mistakenly allowed to spread onto and remain in contact with areas of normal skin (see "Special warnings regarding use of the medicine" in section 2), it may cause irritation.

Spillage of the medicine on your clothes, fabrics, jewelry, metal and polished surfaces, may damage them permanently. Discontinue use and tell your doctor or pharmacist:

- o If you experience intolerable discomfort or if irritation persists.

**If a side effect occurs, if one of the side effects worsens or if you suffer from a side effect not mentioned in the leaflet, consult with the doctor.**

#### **Reporting side effects**

Side effects can be reported to the Ministry of Health by clicking on the link "Report Side Effects of Drug Treatment" found on the Ministry of Health homepage ([www.health.gov.il](http://www.health.gov.il)) that directs you to the online form for reporting side effects, or by entering the link:

<https://sideeffects.health.gov.il/>

### **5. HOW SHOULD THE MEDICINE BE STORED?**

**Avoid poisoning!** This medicine, and any other medicine, must be stored in a safe place out of the reach and sight of children and/or infants, to avoid poisoning. Do not induce vomiting unless explicitly instructed to do so by a doctor. Do not use the medicine after the expiry date (exp. date) appearing on the package and on the label. The expiry date refers to the last day of that month.

Do not discard medicines into the sewer or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help protect the environment.

#### **Storage conditions:**

Do not store above 25°C.

Caution! Flammable material, keep away from fire! See "Special warnings regarding use of the medicine" in section 2.

Close the tube tightly with the cap, at the end of use.

### **6. FURTHER INFORMATION**

**In addition to the active ingredients the medicine also contains:**

Ethyl acetate, Ethanol, Pyroxylin, Camphor.

**What does the medicine look like and what are the contents of the package:**

Clear, colorless gel.

The medicine is provided in a tube containing 8 grams of gel. The tube has a special applicator nozzle. Each pack also includes an emery board.

**License holder:** Trupharm Marketing 1985 Ltd., Poleg Industrial Park, P.O.B. 8105, South Netanya Industrial Zone.

**Manufacturer:** Dermal Laboratories Ltd., Gosmore, Herts, UK.

**Registration number of the medicine in the National Drug Registry of the Ministry of Health:** 053-62-26348

Revised in March 2021 according to MOH guidelines.



TRUPHARM