

**PATIENT LEAFLET IN ACCORDANCE WITH
THE PHARMACISTS' REGULATIONS
(PREPARATIONS) – 1986**

This medicine is dispensed without a doctor's prescription

Spirit Salicyl 2% Floris Liquid

Composition:

Salicylic acid 2% w/v

Inactive ingredients and allergens in the medicine – see section 2 "Important information about some of the ingredients of the medicine" and section 6 "Additional information".

Read the entire leaflet carefully before using the medicine. This leaflet contains concise information about the medicine. If you have any other questions, refer to the doctor or the pharmacist.

This medicine is dispensed without requiring a doctor's prescription and is usually not intended for children and infants under two years of age.

Use the medicine according to the instructions in section 3 "How should you use the medicine?" in this leaflet. Consult the pharmacist if you need more information. Refer to the doctor if symptoms of the ailment worsen or do not improve.

1. What is the medicine intended for?

The medicine is intended for disinfection and peeling of the skin.

Therapeutic class:

Keratolytic preparation from the salicylate group.

2. Before using the medicine

Do not use the preparation if:

- You are sensitive (allergic) to the active ingredient salicylic acid or to any of the other ingredients this medicine contains (see section 6 – "Additional information").

Special warnings regarding the use of the medicine

- Do not light a cigarette or go near an open flame – there is a risk of severe burns. Fabric (clothes, bed linen, dressings etc.) that comes in contact with the medicine will burn more easily and is a serious fire hazard. Washing clothes and bed linen can decrease the accumulation of the medicine, but not remove it completely.
- Avoid applying the preparation on damaged or inflamed skin.
- Avoid applying additional preparations on the same area of skin concomitantly with the preparation.

Drug interactions:

If you are taking, or have recently taken, other medicines including non-prescription medicines and nutritional supplements, tell the doctor or the pharmacist.

Pregnancy, breastfeeding and fertility:

If you are pregnant, might be pregnant or are breastfeeding, consult the doctor or pharmacist before using the medicine. Do not use this medicine during pregnancy or breastfeeding, unless the doctor instructs you to do so.

Use in children

The parents should report to the doctor about any side effect and any other medicine given to the child.

Important information about some of the ingredients of the medicine

The medicine contains alcohol which may cause a burning sensation on damaged skin.

This medicine contains 70.08 ml of alcohol in each 100 ml, a concentration equivalent to 0.7008 ml/ml.

3. How should you use the medicine?

Check with the doctor or pharmacist if you are uncertain about the dosage and how to use the medicine.

The accepted dosage when no other instruction

from the doctor is given is:

Wash and dry the skin, apply on the skin 1-3 times a day using a cotton wool ball soaked in the substance.

Do not exceed the recommended dose.

Method of administration:

Do not swallow, this medicine is intended for external use only.

If you accidentally took a higher dosage

If you took an overdose or if a child swallowed this medicine by mistake, go to the doctor or to a hospital emergency room immediately and take the package of the medicine with you.

Do not take medicines in the dark! Check the label and the dose every time you take a medicine. Wear glasses if you need them.

If you have any other questions regarding the use of the medicine, consult the doctor or the pharmacist.

4. Side effects

As with any medicine, using Spirit Salicyl 2% Floris may cause side effects in some users. Do not be alarmed when reading the list of side effects. You may not experience any of them.

Stop using this medicine and refer to a doctor if the following side effects appear:

Allergic reactions, dry skin and skin irritation.

Application over large skin areas or on the skin of a baby may cause dizziness, deafness and hearing problems.

If a side effect occurs, if one of the side effects worsens, or if you suffer from a side effect not mentioned in this leaflet, consult your doctor.

Reporting side effects

Side effects may be reported to the Ministry of Health by clicking on the link "Report side effects due to medicinal treatment" found on the Ministry of Health website homepage (www.health.gov.il), which will direct you to the online form for reporting side effects, or by clicking on the following link:

<https://sideeffects.health.gov.il>

5. How to store the medicine?

- Avoid poisoning! This medicine and any other medicine must be kept in a closed place out of the reach and sight of children and/or infants to avoid poisoning. Do not induce vomiting without an explicit instruction from the doctor.
- Do not use the medicine after the expiry date (exp. Date) appearing on the package. The expiry date refers to the last day of that month.

Storage conditions:

- Store in a cool place at a temperature not exceeding 25°C.
- Caution! Flammable substance, keep away from fire.
- After opening, the product can be used until it expires.

6. Additional information:

In addition to the active ingredient, the medicine also contains:

Ethanol 96% (Alcohol 96%), Purified water.

What does the medicine look like and what are the contents of the package?

A brown bottle that contains a clear alcoholic solution.

Package content: 100 ml

Manufacturer and license holder:

Ben Shimon Floris Ltd., Industrial Park Misgav, D.N. Misgav 2017400, Israel.

This leaflet was revised in 01/2025 in accordance with the Ministry of Health guidelines.

Registration number of the medicine in the national drug registry of the Ministry of Health: 060-92-27486-00