

Tamiflu®



Oseltamivir

CAPSULES

1. NAME OF THE MEDICINAL PRODUCT

Tamiflu 30 mg
Tamiflu 45 mg
Tamiflu 75 mg

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Tamiflu 30 mg hard capsules

Each hard capsule contains oseltamivir (as phosphate) equivalent to 30 mg of oseltamivir.
For the full list of excipients, see section 6.1.

Tamiflu 45 mg hard capsules

Each hard capsule contains oseltamivir (as phosphate) equivalent to 45 mg of oseltamivir.
For the full list of excipients, see section 6.1.

Tamiflu 75 mg hard capsules

Each hard capsule contains oseltamivir (as phosphate) equivalent to 75 mg of oseltamivir.
For the full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Tamiflu 30 mg hard capsules:

The hard capsule consists of a light yellow opaque body bearing the imprint "ROCHE" and a light yellow opaque cap bearing the imprint "30 mg". Imprints are blue.

Tamiflu 45 mg hard capsules:

The hard capsule consists of a grey opaque body bearing the imprint "ROCHE" and a grey opaque cap bearing the imprint "45 mg". Imprints are blue.

Tamiflu 75 mg hard capsules:

The hard capsule consists of a grey opaque body bearing the imprint "ROCHE" and a light yellow opaque cap bearing the imprint "75 mg". Imprints are blue.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

Tamiflu capsules **75 mg** is indicated for adults patients and patients who weight > 40 kg body weight.

Tamiflu capsules **30 mg and 45 mg** are indicated for adults and children 1 year of age or older.

Treatment of influenza

The treatment of uncomplicated acute illness due to influenza infection in adults and children 1 year of age or older who have been symptomatic for no more than 2 days.

Prophylaxis of influenza

- Post-exposure prevention in adults and children 1 year of age or older following contact with a clinically diagnosed influenza case when influenza virus is circulating in the community.
- The appropriate use of Tamiflu for prevention of influenza should be determined on a case by case basis by the circumstances and the population requiring protection. In exceptional situations (e.g. in case of a mismatch between the circulating and vaccine virus strains, and a pandemic situation) seasonal prevention could be considered in adults and children one year of age or older.

Tamiflu is not a substitute for influenza vaccination.

The use of antivirals for the treatment and prevention of influenza should be determined on the basis of official recommendations. Decisions regarding the use of oseltamivir for treatment and prophylaxis should take into consideration what is known about the characteristics of the circulating influenza viruses, available information on influenza drug susceptibility patterns for each season and the impact of the disease in different geographical areas and patient populations (see section 5.1).

4.2 Posology and method of administration

Posology

75 mg doses can be administered as either

- one 75 mg capsule or
- one 30 mg capsule plus one 45 mg capsule

Adults, and adolescents 13 years and over

Treatment: The recommended oral dose is 75 mg oseltamivir twice daily for 5 days for adolescents (13 to 17 years of age) and adults.

Body Weight	Recommended dose for 5 days
> 40 kg	75 mg twice daily

Treatment should be initiated as soon as possible within the first two days of onset of symptoms of influenza.

Post-exposure prevention: The recommended dose for prevention of influenza following close contact with an infected individual is 75 mg oseltamivir once daily for 10 days for adolescents (13 to 17 years of age) and adults.

Body Weight	Recommended dose for 10 days
> 40 kg	75 mg once daily

Therapy should begin as soon as possible within two days of exposure to an infected individual.

Prevention during an influenza epidemic in the community: The recommended dose for prevention of influenza during a community outbreak is 75 mg oseltamivir once daily for up to 6 weeks. Safety has been demonstrated for up to 12 weeks in immunocompromised patients. The duration of protection lasts for as long as dosing is continued.

Paediatric population

Children 1 to 12 years of age

Tamiflu 30 mg, 45 mg and 75 mg capsules are available for children 1 year of age or older

Treatment: The following weight-adjusted dosing regimens are recommended for treatment of children 1 year of age or older:

Body Weight	Recommended dose for 5 days
15 kg or less	30 mg twice daily
> 15 kg to 23 kg	45 mg twice daily
> 23 kg to 40 kg	60 mg twice daily
> 40 kg	75 mg twice daily

Treatment should be initiated as soon as possible within the first two days of onset of symptoms of influenza.

Post-exposure prevention: The recommended post-exposure prevention dose of Tamiflu is:

Body Weight	Recommended dose for 10 days
15 kg or less	30 mg once daily
> 15 kg to 23 kg	45 mg once daily
> 23 kg to 40 kg	60 mg once daily
> 40 kg	75 mg once daily

Prevention during an influenza epidemic in the community: For prophylaxis in pediatric patients during a community outbreak of oseltamivir susceptible influenza, dosing may be continued for up to 6 weeks.

For instructions on preparing the extemporaneous formulation, see section 6.6.

Special populations

Hepatic impairment

No dose adjustment is required either for treatment or for prevention in patients with mild or moderate hepatic impairment (Child-Pugh score <9). The safety and pharmacokinetics in patients with severe hepatic impairment have not been evaluated.

Renal impairment

Treatment of influenza: Dose adjustment is recommended for adults and adolescents (13 to 17 years of age) with moderate or severe renal impairment. Recommended doses are detailed in the table below.

Creatinine clearance	Recommended dose for treatment
Mild Creatinine Clearance >60-90 mL/min	75 mg twice daily for 5 days
Moderate Creatinine Clearance >30-60 mL/min	30 mg twice daily for 5 days
Severe Creatinine Clearance >10-30 mL/min	30 mg once daily for 5 days
ESRD patient not undergoing dialysis	Not recommended
ESRD Patients on Hemodialysis Creatinine Clearance ≤10 mL/min	30 mg after each haemodialysis session Treatment duration not to exceed 5 days*
ESRD Patients on Continuous Ambulatory Peritoneal Dialysis** Creatinine Clearance ≤10 mL/min	30 mg single dose administered immediately after a dialysis exchange

*Assuming three hemodialysis sessions are performed in the 5-day period. Treatment can be initiated immediately if influenza symptoms develop during the 48 hours between hemodialysis sessions; however, the post-hemodialysis dose should still be administered independently of time of administration of the initial dose.

** Data derived from studies in continuous ambulatory peritoneal dialysis (CAPD) patients.

Prevention of influenza: Dose adjustment is recommended for adults and adolescents (13 to 17 years of age) with moderate or severe renal impairment as detailed in the table below. The duration of prophylaxis is the same as recommended for patients with normal renal function.

Creatinine clearance	Recommended dose for prevention
Mild Creatinine Clearance >60-90 mL/min	75 mg once daily
Moderate Creatinine Clearance >30-60 mL/min	30 mg once daily
Severe Creatinine Clearance >10-30 mL/min	30 mg every second day
ESRD patient not undergoing dialysis	Not recommended

ESRD Patients on Hemodialysis Creatinine Clearance ≤ 10 mL/min	30 mg after every second haemodialysis session*
ESRD Patients on Continuous Ambulatory Peritoneal Dialysis** Creatinine Clearance ≤ 10 mL/min	30 mg once weekly immediately after dialysis exchange

*An initial dose can be administered prior to the start of dialysis.

** Data derived from studies in continuous ambulatory peritoneal dialysis (CAPD) patients.

Elderly

No dose adjustment is required for geriatric patients.

Immunocompromised patients

Treatment: In immunocompromised patients the excretion of influenza virus is longer than in non-immunocompromised patients and it might be reduced by a 10-days treatment.

Seasonal prophylaxis: Safety has been demonstrated for up to 12 weeks in immunocompromised patients.

Method of administration

Oral use.

4.3 Contraindications

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

4.4 Special warnings and precautions for use

Oseltamivir is effective only against illness caused by influenza viruses. There is no evidence for efficacy of oseltamivir in any illness caused by agents other than influenza viruses (see section 5.1).

Tamiflu is not a substitute for influenza vaccination. Use of Tamiflu must not affect the evaluation of individuals for annual influenza vaccination. The protection against influenza lasts only as long as Tamiflu is administered. Tamiflu should be used for the treatment and prevention of influenza only when reliable epidemiological data indicate that influenza virus is circulating in the community. Susceptibility of circulating influenza virus strains to oseltamivir has been shown to be highly variable (see section 5.1). Therefore, prescribers should take into account the most recent information available on oseltamivir susceptibility patterns of the currently circulating viruses when deciding whether to use Tamiflu.

Severe concomitant condition

No information is available regarding the safety and efficacy of oseltamivir in patients with any medical condition sufficiently severe or unstable to be considered at imminent risk of requiring hospitalisation.

Immunocompromised patients

The efficacy of oseltamivir in either treatment or prophylaxis of influenza in immunocompromised patients has not been firmly established.

Cardiac / respiratory disease

Efficacy of oseltamivir in the treatment of subjects with chronic cardiac disease and/or respiratory disease has not been established. No difference in the incidence of complications was observed between the treatment and placebo groups in this population (see section 5.1).

Paediatric population

Tamiflu is not indicated in infants less than 1 year of age.

Severe renal impairment

Dose adjustment is recommended for both treatment and prevention in adolescents (13 to 17 years of age) and adults with severe renal impairment. There is insufficient clinical data available in children (1 year of age or older) with renal impairment to be able to make any dosing recommendation (see sections 4.2 and 5.2).

Neuropsychiatric events

Neuropsychiatric events have been reported during administration of Tamiflu in patients with influenza, especially in children and adolescents. These events are also experienced by patients with influenza without oseltamivir administration. Patients should be closely monitored for behavioural changes, and the benefits and risks of continuing treatment should be carefully evaluated for each patient (see section 4.8).

4.5 Interaction with other medicinal products and other forms of interaction

Pharmacokinetic properties of oseltamivir, such as low protein binding and metabolism independent of the CYP450 and glucuronidase systems (see section 5.2), suggest that clinically significant drug interactions via these mechanisms are unlikely.

Probenecid

No dose adjustment is required when co-administering with probenecid in patients with normal renal function. Co-administration of probenecid, a potent inhibitor of the anionic pathway of renal tubular secretion, results in an approximate 2-fold increase in exposure to the active metabolite of oseltamivir.

Amoxicillin

Oseltamivir has no kinetic interaction with amoxicillin, which is eliminated via the same pathway, suggesting that oseltamivir interaction with this pathway is weak.

Renal elimination

Clinically important drug interactions involving competition for renal tubular secretion are unlikely, due to the known safety margin for most of these substances, the elimination characteristics of the active metabolite (glomerular filtration and anionic tubular secretion) and the excretion capacity of these pathways. However, care should be taken when prescribing oseltamivir in subjects when taking co-excreted agents with a narrow therapeutic margin (e.g. chlorpropamide, methotrexate, phenylbutazone).

Additional information

No pharmacokinetic interactions between oseltamivir or its major metabolite have been observed when co-administering oseltamivir with paracetamol, acetylsalicylic acid, cimetidine, antacids (magnesium and aluminium hydroxides and calcium carbonates), rimantadine or warfarin (in subjects stable on warfarin and without influenza).

4.6 Fertility, pregnancy and lactation

Pregnancy

Influenza is associated with adverse pregnancy and foetal outcomes, with a risk of major congenital malformations, including congenital heart defects. A large amount of data on oseltamivir exposure of pregnant women from post-marketing reports and observational studies (more than 1000 exposed outcomes during the first trimester) indicate no malformative nor fetoneonatal toxicity by oseltamivir.

However, in one observational study, while the overall malformation risk was not increased, the results for major congenital heart defects diagnosed within 12 months of birth were not conclusive. In this study, the rate of major congenital heart defects following oseltamivir exposure during the first trimester was 1.76% (7 infants out of 397 pregnancies) compared to 1.01% in unexposed pregnancies from the general population (Odds Ratio 1.75, 95% Confidence Interval 0.51 to 5.98). The clinical significance of this finding is not clear, as the study had limited power. Additionally, this study was too small to reliably assess individual types of major malformations; moreover women exposed to oseltamivir and women unexposed could not be made fully comparable, in particular whether or not they had influenza.

Animal studies do not indicate reproductive toxicity (see section 5.3).

The use of Tamiflu may be considered during pregnancy if necessary and after considering the available safety and benefit information (for data on benefit in pregnant women please refer to section 5.1 “treatment of influenza in pregnant women”), and the pathogenicity of the circulating influenza virus strain.

Breastfeeding

In lactating rats, oseltamivir and the active metabolite are excreted in milk. Very limited information is available on children breast-fed by mothers taking oseltamivir and on excretion of oseltamivir in breast milk. Limited data

demonstrated that oseltamivir and the active metabolite were detected in breast milk, however the levels were low, which would result in a subtherapeutic dose to the infant. Considering this information, the pathogenicity of the circulating influenza virus strain and the underlying condition of the breastfeeding woman, administration of oseltamivir may be considered, where there are clear potential benefits to breastfeeding mothers.

Fertility

Based on preclinical data, there is no evidence that Tamiflu has an effect on male or female fertility (see section 5.3).

4.7 Effects on ability to drive and use machines

Tamiflu has no influence on the ability to drive and use machines.

4.8 Undesirable effects

Summary of the safety profile

The overall safety profile of Tamiflu is based on data from 6049 adult/adolescent and 1473 paediatric patients treated with Tamiflu or placebo for influenza, and on data from 3990 adult/adolescent and 253 paediatric patients receiving Tamiflu or placebo/no treatment for the prophylaxis of influenza in clinical trials. In addition, 199 immunocompromised adult patients received Tamiflu for the treatment of influenza and 475 immunocompromised patients (including 18 children, of these 10 Tamiflu and 8 placebo) received Tamiflu or placebo for the prophylaxis of influenza.

In adults/adolescents, the most commonly reported adverse reactions (ARs) were nausea and vomiting in the treatment studies, and nausea in the prevention studies. The majority of these ARs were reported on a single occasion on either the first or second treatment day and resolved spontaneously within 1-2 days. In children, the most commonly reported adverse reaction was vomiting. In the majority of patients, these ARs did not lead to discontinuation of Tamiflu.

The following serious adverse reactions have been rarely reported since oseltamivir has been marketed: Anaphylactic and anaphylactoid reactions, hepatic disorders (fulminant hepatitis, hepatic function disorder and jaundice), angioneurotic oedema, Stevens-Johnson syndrome and toxic epidermal necrolysis, gastrointestinal bleeding and neuropsychiatric disorders.

(Regarding neuropsychiatric disorders, see section 4.4.)

Tabulated list of adverse reactions

The ARs listed in the tables below fall into the following categories: Very common ($\geq 1/10$), common ($\geq 1/100$ to $< 1/10$), uncommon ($\geq 1/1,000$ to $< 1/100$), rare ($\geq 1/10,000$ to $< 1/1,000$), and very rare ($< 1/10,000$). ARs are added to the appropriate category in the tables according to the pooled analysis from clinical studies.

Treatment and prevention of influenza in adults and adolescents:

In adult/adolescent treatment and prevention studies, ARs that occurred the most frequently at the recommended dose (75 mg bid for 5 days for treatment and 75 mg od for up to 6 weeks for prophylaxis) are shown in Table 1.

The safety profile reported in subjects who received the recommended dose of Tamiflu for prophylaxis (75 mg once daily for up to 6 weeks) was qualitatively similar to that seen in the treatment studies, despite a longer duration of dosing in the prophylaxis studies.

Table 1 Adverse reactions in studies investigating Tamiflu for treatment and prevention of influenza in adults and adolescents or through post-marketing surveillance

System Organ Class (SOC)	Adverse reactions according to frequency			
	Very common	Common	Uncommon	Rare
Infections and infestations		Bronchitis, Herpes simplex, Nasopharyngitis, Upper respiratory tract infections, Sinusitis		

System Organ Class (SOC)	Adverse reactions according to frequency			
	Very common	Common	Uncommon	Rare
Blood and lymphatic system disorders				Thrombocytopenia
Immune system disorders			Hypersensitivity reaction	Anaphylactic reactions, Anaphylactoid reactions
Psychiatric disorders				Agitation, Abnormal behaviour, Anxiety, Confusion, Delusions, Delirium, Hallucination, Nightmares, Self-injury
Nervous system disorders	Headache	Insomnia	Altered level of consciousness, Convulsion	
Eye disorders				Visual disturbance
Cardiac disorders			Cardiac arrhythmia	
Respiratory, thoracic and mediastinal disorders		Cough, Sore throat, Rhinorrhea		
Gastrointestinal disorders	Nausea	Vomiting Abdominal pain (incl. upper abdominal pain), Dyspepsia		Gastrointestinal bleedings, Haemorrhagic colitis
Hepatobiliary disorders			Elevated liver enzymes	Fulminant hepatitis, Hepatic failure, Hepatitis
Skin and subcutaneous tissue disorders			Eczema, Dermatitis, Rash, Urticaria	Angioneurotic oedema, Erythema multiforme, Stevens-Johnson syndrome, Toxic epidermal necrolysis
General disorders and administration site conditions		Pain Dizziness (incl. vertigo), Fatigue, Pyrexia, Pain in limb		

Treatment and prevention of influenza in children 1 year of age:

A total of 1473 children (including otherwise healthy children aged 1-12 years old and asthmatic children aged 6-12 years old) participated in clinical studies of oseltamivir given for the treatment of influenza. Of those, 851 children received treatment with oseltamivir suspension. A total of 158 children received the recommended dose of Tamiflu once daily in a post-exposure prophylaxis study in households (n = 99), a 6-week paediatric seasonal prophylaxis study (n = 49) and a 12-week paediatric seasonal prophylaxis study in immunocompromised subjects (n = 10).

Table 2 shows the most frequently reported ARs from paediatric clinical trials.

Table 2 Adverse reactions in studies investigating Tamiflu for treatment and prevention of influenza in children (age/weight-based dosing [30 mg to 75 mg o.d.]

System Organ Class (SOC)	Adverse reactions according to frequency			
	Very common	Common	Uncommon	Rare
Infections and infestations		Otitis media		
Nervous system disorders		Headache		
Eye disorders:		Conjunctivitis (including red eyes, eye discharge and eye pain)		
Ear and labyrinth disorders:		Ear ache	Tympanic membrane disorder	
Respiratory, thoracic and mediastinal disorders	Cough, Nasal congestion	Rhinorrhoea		
Gastrointestinal disorders	Vomiting	Abdominal pain (incl. upper abdominal pain), Dyspepsia, Nausea		
Skin and subcutaneous tissue disorders			Dermatitis (including allergic and atopic dermatitis)	

Description of selected adverse reactions

Psychiatric disorders and nervous system disorders

Influenza can be associated with a variety of neurologic and behavioural symptoms which can include events such as hallucinations, delirium, and abnormal behaviour, in some cases resulting in fatal outcomes. These events may occur in the setting of encephalitis or encephalopathy but can occur without obvious severe disease.

In patients with influenza who were receiving Tamiflu, there have been postmarketing reports of convulsions and delirium (including symptoms such as altered level of consciousness, confusion, abnormal behaviour, delusions, hallucinations, agitation, anxiety, nightmares), in a very few cases resulting in self-injury or fatal outcomes. These events were reported primarily among paediatric and adolescent patients and often had an abrupt onset and rapid resolution. The contribution of Tamiflu to those events is unknown. Such neuropsychiatric events have also been reported in patients with influenza who were not taking Tamiflu.

Hepato-biliary disorders

Hepato-biliary system disorders, including hepatitis and elevated liver enzymes in patients with influenza-like illness. These cases include fatal fulminant hepatitis/hepatic failure.

Other special populations

Older people and patients with chronic cardiac and/or respiratory disease

The population included in the influenza treatment studies is comprised of otherwise healthy adults/adolescents and patients “at risk” (patients at higher risk of developing complications associated with influenza, e.g. older people and patients with chronic cardiac or respiratory disease). In general, the safety profile in the patients “at risk” was qualitatively similar to that in otherwise healthy adults/adolescents.

Immunocompromised patients

In a double blind study for the treatment of influenza, a total of 199 adult immunocompromised patients (evaluable for safety) were randomized to receive Tamiflu for 10 days: 98 patients received the standard dose (75 mg twice daily) and 101 patients received the double dose (150 mg twice daily). The safety profile of Tamiflu observed in this study was consistent with that observed in previous clinical trials where Tamiflu was administered for treatment of influenza in non-immunocompromised patients (otherwise healthy patients or “at risk” patients [i.e., those with respiratory and/or cardiac co-morbidities]). The percentage of patients reporting adverse events was lower in the standard dose group compared to the double dose group (49.0% vs 59.4 %, respectively) (See section 5.1).

In a 12-week prophylaxis study in 475 immunocompromised patients, including 18 children 1 to 12 years of age and older, the safety profile in the 238 patients who received oseltamivir was consistent with that previously observed in Tamiflu prophylaxis clinical studies.

Children with pre-existing bronchial asthma

In general, the adverse reaction profile in children with pre-existing bronchial asthma was qualitatively similar to that of otherwise healthy children.

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product.

Any suspected adverse events should be reported to the Ministry of Health according to the National Regulation by using an online form /<https://sideeffects.health.gov.il>

4.9 Overdose

Reports of overdoses with Tamiflu have been received from clinical trials and during post-marketing experience. In the majority of cases reporting overdose, no adverse events were reported.

Adverse events reported following overdose were similar in nature and distribution to those observed with therapeutic doses of Tamiflu, described in section 4.8 Undesirable effects.

No specific antidote is known.

Paediatric population

Overdose has been reported more frequently for children than adults and adolescents. Caution should be exercised when preparing Tamiflu oral suspension and when administering Tamiflu products to children.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Antivirals for systemic use, neuraminidase inhibitors ATC code: J05AH02

Oseltamivir phosphate is a pro-drug of the active metabolite (oseltamivir carboxylate). The active metabolite is a selective inhibitor of influenza virus neuraminidase enzymes, which are glycoproteins found on the virion surface. Viral neuraminidase enzyme activity is important both for viral entry into uninfected cells and for the release of recently formed virus particles from infected cells, and for the further spread of infectious virus in the body.

Oseltamivir carboxylate inhibits influenza A and B neuraminidases *in vitro*. Oseltamivir phosphate inhibits influenza virus infection and replication *in vitro*. Oseltamivir given orally inhibits influenza A and B virus replication and pathogenicity *in vivo* in animal models of influenza infection at antiviral exposures similar to that achieved in man with 75 mg twice daily.

Antiviral activity of oseltamivir was supported for influenza A and B by experimental challenge studies in healthy volunteers.

Neuraminidase enzyme IC50 values for oseltamivir for clinically isolated influenza A ranged from 0.1 nM to 1.3 nM, and for influenza B was 2.6 nM. Higher IC50 values for influenza B, up to a median of 8.5 nM, have been observed in published studies.

Clinical studies

Treatment of influenza infection

The indication is based on clinical studies of naturally occurring influenza in which the predominant infection was influenza A.

Oseltamivir is effective only against illnesses caused by influenza virus. Statistical analyses are therefore presented only for influenza-infected subjects. In the pooled treatment study population, which included both influenza-positive and -negative subjects (ITT), primary efficacy was reduced proportionally to the number of influenza-negative individuals. In the overall treatment population, influenza infection was confirmed in 67 % (range 46 % to 74 %) of the recruited patients. Of the older subjects, 64 % were influenza-positive and of those with chronic cardiac and/or respiratory disease 62 % were influenza-positive. In all phase III treatment studies, patients were recruited only during the period in which influenza was circulating in the local community.

Adults and adolescents 13 years of age and older: Patients were eligible if they reported within 36 hours of onset of symptoms, had fever ≥ 37.8 °C, accompanied by at least one respiratory symptom (cough, nasal symptoms or sore throat) and at least one systemic symptom (myalgia, chills/sweats, malaise, fatigue or headache). In a pooled analysis of all influenza-positive adults and adolescents (N = 2,413) enrolled into treatment studies, oseltamivir 75 mg twice daily for 5 days reduced the median duration of influenza illness by approximately one day from 5.2 days (95 % CI 4.9 – 5.5 days) in the placebo group to 4.2 days (95 % CI 4.0 – 4.4 days; $p \leq 0.0001$).

The proportion of subjects who developed specified lower respiratory tract complications (mainly bronchitis) treated with antibiotics was reduced from 12.7 % (135/1,063) in the placebo group to 8.6 % (116/1,350) in the oseltamivir treated population ($p = 0.0012$).

Treatment of influenza in high risk populations: The median duration of influenza illness in older subjects (≥ 65 years) and in subjects with chronic cardiac and/or respiratory disease receiving oseltamivir 75 mg twice daily for 5 days was not reduced significantly. The total duration of fever was reduced by one day in the groups treated with oseltamivir. In influenza-positive older people, oseltamivir significantly reduced the incidence of specified lower respiratory tract complications (mainly bronchitis) treated with antibiotics from 19 % (52/268) in the placebo group to 12 % (29/250) in the oseltamivir treated population ($p = 0.0156$).

In influenza-positive patients with chronic cardiac and/or respiratory disease, the combined incidence of lower respiratory tract complications (mainly bronchitis) treated with antibiotics was 17 % (22/133) in the placebo group and 14 % (16/118) in the oseltamivir treated population ($p = 0.5976$).

Treatment of influenza in pregnant women: No controlled clinical studies have been conducted on the use of oseltamivir in pregnant women, however, there is evidence from post-marketing and retrospective observational studies showing benefit of the current dosing regimen in this patient population in terms of lower morbidity/mortality. Results from pharmacokinetic analyses indicate a lower exposure to the active metabolite, however dose adjustments are not recommended for pregnant women in the treatment or prophylaxis of influenza (see section 5.2, Pharmacokinetics, Special Population).

Treatment of influenza in children: In a study of otherwise healthy children (65 % influenza-positive) aged 1 to 12 years (mean age 5.3 years) who had fever (≥ 37.8 °C) plus either cough or coryza, 67 % of influenza-positive patients were infected with influenza A and 33 % with influenza B. Oseltamivir treatment, started within 48 hours of onset of symptoms, significantly reduced the time to freedom from illness (defined as the simultaneous return to normal health and activity and alleviation of fever, cough and coryza) by 1.5 days (95 % CI 0.6 – 2.2 days; $p < 0.0001$) compared to placebo. Oseltamivir reduced the incidence of acute otitis media from 26.5 % (53/200) in the placebo group to 16 % (29/183) in the oseltamivir treated children ($p = 0.013$).

A second study was completed in 334 asthmatic children aged 6 to 12 years old of which 53.6 % were influenza-positive. In the oseltamivir treated group, the median duration of illness was not reduced significantly.

By day 6 (the last day of treatment) FEV₁ had increased by 10.8 % in the oseltamivir treated group compared to 4.7 % on placebo (p = 0.0148) in this population.

Treatment of influenza B infection: Overall, 15 % of the influenza-positive population were infected by influenza B, proportions ranging from 1 to 33 % in individual studies. The median duration of illness in influenza B infected subjects did not differ significantly between the treatment groups in individual studies. Data from 504 influenza B infected subjects were pooled across all studies for analysis. Oseltamivir reduced the time to alleviation of all symptoms by 0.7 days (95 % CI 0.1 – 1.6 days; p = 0.022) and the duration of fever (≥ 37.8 °C), cough and coryza by one day (95 % CI 0.4 – 1.7 days; p < 0.001) compared to placebo.

Treatment of influenza in immunocompromised adults: A randomized, double blind study, to evaluate safety and characterize the effects of oseltamivir on the development of resistant influenza virus (primary analysis) in influenza-infected adult immunocompromised patients, included 151 patients evaluable for efficacy of oseltamivir (secondary analysis, not powered). The study included solid organ transplant [SOT] patients, haematopoietic stem cell transplant [HSCT] patients, HIV positive patients with a CD4+ cell count <500 cells/mm³, patients on systemic immunosuppressive therapy, and those with haematological malignancy. These patients were randomized to be treated, within 96 hours of symptoms onset, with standard dose (73 patients) or double dose (78 patients) of oseltamivir, for a duration of 10 days.

The median time to resolution of symptoms (TTRS) was similar between the standard dose group (103 hours [90% CI 75.4-110.0]) and double dose group (104 hours [90% CI 65.8-131.0]). The proportion of patients with secondary infections in the standard dose group and double dose group was comparable (8.2% vs 5.1%).

Prevention of influenza

The efficacy of oseltamivir in preventing naturally occurring influenza illness has been demonstrated in a post-exposure prevention study in households and two seasonal prevention studies. The primary efficacy parameter for all of these studies was the incidence of laboratory-confirmed influenza. The virulence of influenza epidemics is not predictable and varies within a region and from season to season, therefore the number needed to treat (NNT) in order to prevent one case of influenza illness varies.

Post-exposure prevention: In a study in contacts (12.6 % vaccinated against influenza) of an index case of influenza, oseltamivir 75 mg once daily was started within 2 days of onset of symptoms in the index case and continued for seven days. Influenza was confirmed in 163 out of 377 index cases. Oseltamivir significantly reduced the incidence of clinical influenza illness occurring in the contacts of confirmed influenza cases from 24/200 (12 %) in the placebo group to 2/205 (1 %) in the oseltamivir group (92 % reduction [95 % CI 6 – 16; p \leq 0.0001]). The number needed to treat (NNT) in contacts of true influenza cases was 10 (95 % CI 9 – 12) and was 16 (95 % CI 15 – 19) in the whole population (ITT) regardless of infection status in the index case.

The efficacy of oseltamivir in preventing naturally occurring influenza illness has been demonstrated in a post-exposure prevention study in households that included adults, adolescents, and children aged 1 to 12 years, both as index cases and as family contacts. The primary efficacy parameter for this study was the incidence of laboratory-confirmed clinical influenza in the households. Oseltamivir prophylaxis lasted for 10 days. In the total population, there was a reduction in the incidence of laboratory-confirmed clinical influenza in households from 20 % (27/136) in the group not receiving prevention to 7 % (10/135) in the group receiving prevention (62.7 % reduction [95 % CI 26.0 – 81.2; p = 0.0042]). In households of influenza-infected index cases, there was a reduction in the incidence of influenza from 26 % (23/89) in the group not receiving prevention to 11 % (9/84) in the group receiving prevention (58.5 % reduction [95 % CI 15.6 – 79.6; p = 0.0114]). According to subgroup analysis in children at 1 to 12 years of age, the incidence of laboratory-confirmed clinical influenza among children was significantly reduced from 19 % (21/111) in the group not receiving prevention to 7 % (7/104) in the group receiving prevention (64.4 % reduction [95 % CI 15.8 – 85.0; p = 0.0188]). Among children who were not already shedding virus at baseline, the incidence of laboratory-confirmed clinical influenza was reduced from 21 % (15/70) in the group not receiving prevention to 4 % (2/47) in the group receiving prevention (80.1 % reduction [95 % CI 22.0 – 94.9; p = 0.0206]). The NNT for the total paediatric population was 9 (95 % CI 7 – 24) and 8 (95 % CI 6, upper limit not estimable) in the whole population (ITT) and in paediatric contacts of infected index cases (ITTII), respectively.

Prevention during an influenza epidemic in the community: In a pooled analysis of two other studies conducted in unvaccinated otherwise healthy adults, oseltamivir 75 mg once daily given for 6 weeks significantly reduced the incidence of clinical influenza illness from 25/519 (4.8 %) in the placebo group to 6/520 (1.2 %) in the oseltamivir group (76 % reduction [95 % CI 1.6 – 5.7; p = 0.0006]) during a community outbreak of influenza. The NNT in this study was 28 (95 % CI 24 – 50).

A study in older people in nursing homes, where 80 % of participants received vaccine in the season of the study, oseltamivir 75 mg once daily given for 6 weeks significantly reduced the incidence of clinical influenza illness from 12/272 (4.4 %) in the placebo group to 1/276 (0.4 %) in the oseltamivir group (92 % reduction [95 % CI 1.5 – 6.6; p = 0.0015]). The NNT in this study was 25 (95 % CI 23 – 62).

Prophylaxis of influenza in immunocompromised patients: A double-blind, placebo-controlled, randomised study was conducted for seasonal prophylaxis of influenza in 475 immunocompromised patients (388 patients with solid organ transplantation [195 placebo; 193 oseltamivir], 87 patients with haemopoietic stem cell transplantation [43 placebo; 44 oseltamivir], no patient with other immunosuppressant conditions), including 18 children 1 to 12 years of age. The primary endpoint in this study was the incidence of laboratory-confirmed clinical influenza as determined by viral culture and/or a four-fold rise in HAI antibodies. The incidence of laboratory-confirmed clinical influenza was 2.9 % (7/238) in the placebo group and 2.1 % (5/237) in the oseltamivir group (95 % CI -2.3 % – 4.1 %; p = 0.772).

Specific studies have not been conducted to assess the reduction in the risk of complications.

Oseltamivir resistance

Clinical studies: The risk of emergence of influenza viruses with reduced susceptibility or frank resistance to oseltamivir has been examined during Roche-sponsored clinical studies. Developing oseltamivir-resistant virus during treatment was more frequent in children than adults, ranging from less than 1 % in adults to 18 % in infants aged below 1 year. Children who were found to carry oseltamivir-resistant virus in general shed the virus for a prolonged period compared with subjects with susceptible virus. However treatment-emergent resistance to oseltamivir did not affect treatment response and caused no prolongation of influenza symptoms.

An overall higher incidence of oseltamivir-resistance was observed in adult immunocompromised patients treated with standard dose or double dose of oseltamivir for a duration of 10 days [14.9% (10/67) in standard dose group and 2.8% (2/71) in double dose group], compared to data from studies with oseltamivir-treated otherwise healthy adult patients. The majority of patients that developed resistance were transplant recipients (8/10 patients in the standard dose group and 2/2 patients in the double dose group). Most of the patients with oseltamivir-resistant virus were infected with influenza type A and had prolonged viral shedding.

Incidence of Oseltamivir Resistance in Clinical Studies

Patient Population	Patients with Resistance Mutations (%)	
	Phenotyping*	Geno- and Phenotyping*
Adults and adolescents	0.88% (21/2377)	1.12% (27/2391)
Children (1-12 years)	3.89% (66/1698)	4.24% (72/1698)

* Full genotyping was not performed in all studies.

Prophylaxis of Influenza

There has been no evidence for emergence of drug resistance associated with the use of Tamiflu in clinical studies conducted to date in post-exposure (7 days), post-exposure within household groups (10 days) and seasonal (42 days) prevention of influenza in immunocompetent patients. There was no resistance observed during a 12-week prophylaxis study in immunocompromised patients.

Clinical and surveillance data: Natural mutations associated with reduced susceptibility to oseltamivir *in vitro* have been detected in influenza A and B viruses isolated from patients without exposure to oseltamivir. Resistant strains selected during oseltamivir treatment have been isolated from both immunocompetent and immunocompromised patients. Immunocompromised patients and young children are at a higher risk of developing oseltamivir-resistant virus during treatment.

Oseltamivir-resistant viruses isolated from oseltamivir-treated patients and oseltamivir-resistant laboratory strains of influenza viruses have been found to contain mutations in N1 and N2 neuraminidases. Resistance

mutations tend to be viral sub-type specific. Since 2007 naturally occurring resistance associated with the H275Y mutation in seasonal H1N1 strains has been sporadically detected. The susceptibility to oseltamivir and the prevalence of such viruses appear to vary seasonally and geographically. In 2008, H275Y was found in > 99 % of circulating H1N1 influenza isolates in Europe. The 2009 H1N1 influenza (“swine flu”) was almost uniformly susceptible to oseltamivir, with only sporadic reports of resistance in connection with both therapeutic and prophylactic regimens.

5.2 Pharmacokinetic properties

General Information

Absorption

Oseltamivir is readily absorbed from the gastrointestinal tract after oral administration of oseltamivir phosphate (pro-drug) and is extensively converted by predominantly hepatic esterases to the active metabolite (oseltamivir carboxylate). At least 75 % of an oral dose reaches the systemic circulation as the active metabolite. Exposure to the pro-drug is less than 5 % relative to the active metabolite. Plasma concentrations of both pro-drug and active metabolite are proportional to dose and are unaffected by co-administration with food.

Distribution

The mean volume of distribution at steady state of the oseltamivir carboxylate is approximately 23 litres in humans, a volume equivalent to extracellular body fluid. Since neuraminidase activity is extracellular, oseltamivir carboxylate distributes to all sites of influenza virus spread.

The binding of the oseltamivir carboxylate to human plasma protein is negligible (approximately 3 %).

Biotransformation

Oseltamivir is extensively converted to oseltamivir carboxylate by esterases located predominantly in the liver. *In vitro* studies demonstrated that neither oseltamivir nor the active metabolite is a substrate for, or an inhibitor of, the major cytochrome P450 isoforms. No phase 2 conjugates of either compound have been identified *in vivo*.

Elimination

Absorbed oseltamivir is primarily (> 90 %) eliminated by conversion to oseltamivir carboxylate. It is not further metabolised and is eliminated in the urine. Peak plasma concentrations of oseltamivir carboxylate decline with a half-life of 6 to 10 hours in most subjects. The active metabolite is eliminated entirely by renal excretion. Renal clearance (18.8 l/h) exceeds glomerular filtration rate (7.5 l/h) indicating that tubular secretion occurs in addition to glomerular filtration. Less than 20 % of an oral radiolabelled dose is eliminated in faeces.

Other special populations

Paediatric population

There are no data available for infants below 1 year of age for post exposure prevention of influenza.

Children 1 year of age or older: The pharmacokinetics of oseltamivir have been evaluated in single-dose pharmacokinetic studies in children and adolescents 1 to 16 years of age. Multiple-dose pharmacokinetics were studied in a small number of children enrolled in a clinical efficacy study. Younger children cleared both the pro-drug and its active metabolite faster than adults, resulting in a lower exposure for a given mg/kg dose. Doses of 2 mg/kg give oseltamivir carboxylate exposures comparable to those achieved in adults receiving a single 75 mg dose (approximately 1 mg/kg). The pharmacokinetics of oseltamivir in children and adolescents 12 years of age or older are similar to those in adults.

Elderly

Exposure to the active metabolite at steady state was 25 to 35 % higher in older people (age 65 to 78 years) compared to adults less than 65 years of age given comparable doses of oseltamivir. Half-lives observed in older people were similar to those seen in young adults. On the basis of drug exposure and tolerability, dosage

adjustments are not required for older people unless there is evidence of moderate or severe renal impairment (creatinine clearance below 60 ml /min) (see section 4.2).

Renal impairment

Administration of 100 mg oseltamivir phosphate twice daily for 5 days to patients with various degrees of renal impairment showed that exposure to oseltamivir carboxylate is inversely proportional to declining renal function. For dosing, see section 4.2.

Hepatic impairment

In vitro studies have concluded that exposure to oseltamivir is not expected to be increased significantly nor is exposure to the active metabolite expected to be significantly decreased in patients with hepatic impairment (see section 4.2).

Pregnant Women

A pooled population pharmacokinetic analysis indicates that the Tamiflu dosage regimen described in Section 4.2 Posology and method of administration results in lower exposure (30% on average across all trimesters) to the active metabolite in pregnant women compared to non-pregnant women. The lower predicted exposure however, remains above inhibitory concentrations (IC₉₅ values) and at a therapeutic level for a range of influenza virus strains. In addition, there is evidence from observational studies showing benefit of the current dosing regimen in this patient population. Therefore, dose adjustments are not recommended for pregnant women in the treatment or prophylaxis of influenza (see section 4.6 Fertility, pregnancy and lactation).

Immunocompromised Patients

Population pharmacokinetic analysis indicates that treatment of adult immunocompromised patients with oseltamivir (as described in Section 4.2. Posology and method of administration) results in an increased exposure (of up to 50%) to the active metabolite when compared to adult non-immunocompromised patients with comparable creatinine clearance. Due to the wide safety margin of the active metabolite, no dose adjustments are required in adults due to their immunocompromised status. However, for adult immunocompromised patients with renal impairment, doses should be adjusted as outlined in section 4.2. Posology and method of administration.

5.3 Preclinical safety data

Preclinical data reveal no special hazard for humans based on conventional studies of safety pharmacology, repeated-dose toxicity and genotoxicity. Results of the conventional rodent carcinogenicity studies showed a trend towards a dose-dependent increase in the incidence of some tumours that are typical for the rodent strains used. Considering the margins of exposure in relation to the expected exposure in the human use, these findings do not change the benefit-risk of Tamiflu in its adopted therapeutic indications.

Teratology studies have been conducted in rats and rabbits at doses of up to 1,500 mg/kg/day and 500 mg/kg/day, respectively. No effects on foetal development were observed. A rat fertility study up to a dose of 1,500 mg/kg/day demonstrated no adverse reactions on either sex. In pre- and post-natal rat studies, prolonged parturition was noted at 1,500 mg/kg/day: the safety margin between human exposure and the highest no-effect dose (500 mg/kg/day) in rats is 480-fold for oseltamivir and 44-fold for the active metabolite, respectively. Foetal exposure in the rats and rabbits was approximately 15 to 20 % of that of the mother.

In lactating rats, oseltamivir and the active metabolite are excreted in the milk. Limited data indicate that oseltamivir and the active metabolite are excreted in human milk. Extrapolation of the animal data provides estimates of 0.01 mg/day and 0.3 mg/day for the respective compounds.

A potential for skin sensitisation to oseltamivir was observed in a "maximisation" test in guinea pigs. Approximately 50 % of the animals treated with the unformulated active substance showed erythema after challenging the induced animals. Reversible irritancy of rabbits' eyes was detected.

Whereas very high oral single doses of oseltamivir phosphate salt, up to the highest dose tested (1,310 mg/kg), had no adverse reactions in adult rats, such doses resulted in toxicity in juvenile 7-day-old rat pups, including death. These reactions were seen at doses of 657 mg/kg and higher. At 500 mg/kg, no adverse reactions were seen, including upon chronic treatment (500 mg/kg/day administered from 7 to 21 days post partum).

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Tamiflu 30 mg hard capsules

Capsule core

Pregelatinised starch

Talc

Povidone K30

Croscarmellose sodium

Sodium stearyl fumarate

Capsule shell

Gelatin

Titanium dioxide (E171)

Yellow iron oxide (E172)

Red iron oxide (E172)

Printing ink

Tamiflu 45 mg hard capsules

Capsule core

Pregelatinised starch

Talc

Povidone K30

Croscarmellose sodium

Sodium stearyl fumarate

Capsule shell

Gelatin

Titanium dioxide (E171)

Black iron oxide (E172)

Printing ink

Tamiflu 75 mg hard capsules

Capsule core

Pregelatinised starch

Talc

Povidone K30

Croscarmellose sodium

Sodium stearyl fumarate

Capsule shell

Gelatin

Titanium dioxide (E171)

Yellow iron oxide (E172)

Red iron oxide (E172)

Black iron oxide (E172)

Printing ink

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

The expiry date of the product is indicated on the packaging materials.

Storage of the pharmacy compounded suspension

Shelf life of 10 days when stored below 25 °C.

6.4 Special precautions for storage

Do not store above 25 °C.

For storage conditions of the pharmacy compounded suspension, see section 6.3.

6.5 Nature and contents of container

Triplex blister pack (PVC/PE/PVDC, sealed with aluminium foil).

Pack-size 10 capsules.

6.6 Special precautions for disposal and other handling

Any unused product or waste material should be disposed of in accordance with local requirements.

Extemporaneous formulation

When Tamiflu powder for oral suspension is not available

Commercially manufactured Tamiflu for oral suspension (6 mg/ml) is the preferred product for paediatric and adult patients who have difficulties swallowing capsules or where lower doses are needed. In the event that commercially manufactured Tamiflu powder for oral suspension is not available, the pharmacist may compound a suspension (6 mg/ml) from Tamiflu capsules or patients can prepare the suspension from capsules at home.

The pharmacy preparation should be preferred to home preparation. Detailed information on the home preparation can be found in the package leaflet of Tamiflu capsules under “Making liquid Tamiflu at home”.

Syringes of appropriate volume and grading should be provided for administering the pharmacy compounded suspension as well as for the procedures involved in the home preparation. In both cases, the correct volumes should preferably be marked on the syringes.

Pharmacy compounding

Pharmacy compounded 6 mg/ml suspension prepared from capsules

Adults, adolescents and infants and children 1 year of age or older who are unable to swallow intact capsules

This procedure describes the preparation of a 6 mg/ml suspension that will provide one patient with enough medicine for a 5-day course of treatment or a 10-day course of prophylaxis.

The pharmacist may compound a 6 mg/ml suspension from Tamiflu 30 mg, 45 mg or 75 mg capsules using water containing 0.05 % w/v sodium benzoate added as a preservative.

First, calculate the total volume needed to be compounded and dispensed to provide a 5-day course of treatment or a 10-day course of prophylaxis for the patient. The total volume required is determined by the weight of the patient according to the recommendation in the table below. To allow for accurate volume withdrawal, the column indicating measurement loss is to be considered for compounding.

Volume of pharmacy compounded 6 mg/ml suspension prepared based upon the patient's weight for 5-day treatment or 10-day prophylaxis course

Body weight (kg)	Total volume to compound per patient weight (ml) Measurement loss not considered	Total volume to compound per patient weight (ml) Measurement loss considered
15 kg or less	50 ml	60 ml or 75 ml*
> 15 kg to 23 kg	75 ml	90 ml or 100 ml*
> 23 kg to 40 kg	100 ml	125 ml
> 40 kg	125 ml	137.5 ml (or 150 ml)*

* Depending on the capsule strength used.

Second, determine the number of capsules and the amount of vehicle (water containing 0.05 % w/v sodium benzoate added as a preservative) that is needed to prepare the total volume (calculated from the table above) of pharmacy compounded 6 mg/ml suspension as shown in the table below:

Number of capsules and amount of vehicle needed to prepare the total volume of a pharmacy compounded 6 mg/ml suspension (for 5 days of treatment or 10-days of prophylaxis)

Total volume of compounded suspension to be prepared	Required number of Tamiflu capsules (mg of oseltamivir)			Required volume of vehicle
	75 mg	45 mg	30 mg	
60 ml	Please use alternative capsule strength*	8 capsules (360 mg)	12 capsules (360 mg)	59.5 ml
75 ml	6 capsules (450 mg)	10 capsules (450 mg)	15 capsules (450 mg)	74 ml
90 ml	Please use alternative capsule strength*	12 capsules (540 mg)	18 capsules (540 mg)	89 ml
100 ml	8 capsules (600 mg)	Please use alternative capsule strength*	20 capsules (600 mg)	98.5 ml
125 ml	10 capsules (750 mg)	Please use alternative capsule strength*	25 capsules (750 mg)	123.5 ml
137.5 ml	11 capsules (825 mg)	Please use alternative capsule strength*	Please use alternative capsule strength*	136 ml

* There is no combination of this capsule strength that can be used to achieve the target concentration; therefore, please use an alternative capsule strength.

Third, follow the procedure below for compounding the 6 mg/ml suspension from Tamiflu capsules:

1. In a glass beaker of suitable size place the stated amount of water containing 0.05 % w/v sodium benzoate added as a preservative.
2. Open the stated amount of Tamiflu capsules and transfer the content of each capsule directly to the preserved water in the glass beaker.
3. With a suitable stirring device, stir for 2 minutes.

(Note: The drug substance, oseltamivir phosphate, readily dissolves in water. The suspension is caused by some of the excipients of Tamiflu capsules, which are insoluble.)

4. Transfer the suspension to an amber glass or amber polyethyleneterephthalate (PET) bottle. A funnel may be used to eliminate any spillage.
5. Close the bottle using a child-resistant cap.
6. Put an ancillary label on the bottle indicating “Shake Gently Before Use”.

(Note: This compounded suspension should be gently shaken prior to administration to minimise the tendency for air entrapment.)

7. Instruct the parent or caregiver that any remaining material following completion of therapy must be discarded. It is recommended that this information be provided by either affixing an ancillary label to the bottle or adding a statement to the pharmacy label instructions.
8. Place an appropriate expiration date label according to storage condition (see section 6.3).

Place a pharmacy label on the bottle that includes the patient’s name, dosing instructions, use by date, name of medicinal product and any other required information to be in compliance with local pharmacy regulations. Refer to the table below for the proper dosing instructions.

Dosing chart for pharmacy-compounded 6 mg/ml suspension prepared from Tamiflu capsules for patients 1 year of age or older

Body weight (kg)	Dose (mg)	Volume per dose 6 mg/ml	Treatment dose (for 5 days)	Prophylaxis dose (for 10 days)
15 kg or less	30 mg	5 ml	5 ml twice daily	5 ml once daily
> 15 kg to 23 kg	45 mg	7.5 ml	7.5 ml twice daily	7.5 ml once daily
> 23 kg to 40 kg	60 mg	10 ml	10 ml twice daily	10 ml once daily
> 40 kg	75 mg	12.5 ml	12.5 ml twice daily	12.5 ml once daily

Dispense the pharmacy compounded suspension with a graduated oral syringe for measuring small amounts of suspension. If possible, mark or highlight the graduation corresponding to the appropriate dose (according to the dosing table above) on the oral syringe for each patient.

The appropriate dose must be mixed by the caregiver with an equal quantity of sweet liquid food, such as sugar water, chocolate syrup, cherry syrup, dessert toppings (like caramel or fudge sauce) to mask the bitter taste.

Home preparation

When commercially manufactured Tamiflu oral suspension is not available, a pharmacy compounded suspension prepared from Tamiflu capsules must be used (see detailed instructions above). If the commercially manufactured Tamiflu oral suspension and the pharmacy compounded suspension is also not available, Tamiflu suspension may be prepared at home.

When appropriate capsule strengths are available for the dose needed, the dose is given by opening the capsule and mixing its contents with no more than one teaspoon of a suitable sweetened food product. The bitter taste can be masked by products such as sugar water, chocolate syrup, cherry syrup, dessert toppings (like caramel or fudge sauce). The mixture should be stirred and given entirely to the patient. The mixture must be swallowed immediately after its preparation.

When only 75 mg capsules are available, and doses of 30 mg or 45 mg are needed, the preparation of Tamiflu suspension involves additional steps. Detailed instructions can be found in the package leaflet of Tamiflu capsules under “Making liquid Tamiflu at home”.

7. MARKETING AUTHORISATION HOLDER

Roche Pharmaceuticals (Israel) Ltd., 6 Hacharash, P.O. B. 6391, Hod Hasharon 4524079.

8. LICENSE NUMBER(S)

Tamiflu Capsules 30 mg: 138 02 31733

Tamiflu Capsules 45 mg: 138 03 31734

Tamiflu Capsules 75 mg: 118 79 29952

9. MANUFACTURER

F. Hoffmann-La Roche Ltd., Grenzacherstrasse 124,CH-4058, Basel, Switzerland

Revised in February 2025