

# Ozurdex<sup>®</sup> Patient Guide

(dexamethasone intravitreal implant 0.7mg)

*This patient guide contains important safety information you should be aware of before and after treatment with Ozurdex<sup>®</sup>.*

## Why Should I Read this Patient Guide? 2

### Treatment 2

How Should I Prepare For My Treatment? 2

What Will My Treatment Involve? 2

What Will Happen After My Treatment? 3

Where To Find More Information 4

## Why Should I Read this Patient Guide?

Treatment with Ozurdex<sup>®</sup> may be associated with the following risks:

- Increased pressure in the eye, which can lead to glaucoma (an eye disease that causes vision loss).
- Inflammation inside the eye (usually caused by infection).

To prevent progression/minimize serious consequences, it is important for you to be aware of early signs and symptoms of these risks associated with Ozurdex<sup>®</sup>.

## How Should I Prepare For My Treatment?

Before your treatment, your doctor will give you antibiotic eye drops. You should use these for 3 days prior to your treatment. You should continue to use the eye drops for 3 days following your treatment.

Please do NOT wear any make-up on the day of the procedure.

## What Will My Treatment Involve?

To prepare you for your treatment, the doctor will:

- Clean your eye to minimize infection and may administer further antibiotic eye drops.
- Cover your face with a special drape.
- Numb your eye with anaesthetic so you feel no pain.
- Use a device to gently hold your eye open.

The doctor will then use a specially designed applicator to inject the medicine into your eye, through the white part of your eye. You may hear a clicking sound during this procedure as well as feeling a bit of pressure on the eye.

Your doctor will then perform routine eye examinations to confirm that the medicine has been successfully administered.

## What Will Happen After My Treatment?

After the injection:

- Your vision may be blurred for about a day.
- You may also see some floaters or spots, which is normal and should go away over time.
- If these symptoms do not go away, or worsen, please contact your doctor.
- If you normally drive or use machinery, you should not resume these activities until all blurring of your vision has stopped.

***You should contact your doctor or nurse immediately if you experience at any time any of these signs and symptoms following Ozurdex® treatment.***

- Worsening of vision after the injection.
- Pain or discomfort in or around the eye.
- An increase in floaters or spots in the vision.
- Blurred vision which lasts more than a day after the injection.
- Redness in the eye which continues to get worse.
- Discharge from the eye.



## Where You Can Find More Information About Your Medicine

As with all medicines, Ozurdex® can cause side effects, but not everybody gets them, such as:

- Ocular events of Cataract and retinal detachment.
- A detailed list of possible side effects can be found in the Patient Information Leaflet (PIL) that comes with Ozurdex®.

For more information about your medicine, please read the PIL. It can be accessed in a digital form at the MoH website

<https://israeldrugs.health.gov.il/#!/byDrug>

If you would like additional copies of the Patient Guide, please contact [PVIsrael@abbvie.com](mailto:PVIsrael@abbvie.com)

To report any side effects from the use of your medicine, please contact AbbVie at [PVIsrael@abbvie.com](mailto:PVIsrael@abbvie.com). You may also report any side effect to the MoH website [sideeffects.health.gov.il](https://sideeffects.health.gov.il)

This patient card and content were checked and approved by the Ministry of Health on 03 September 2024, V3.0