

# **PNEUMOVAX® 23**

## **Pneumococcal Vaccine Polyvalent Solution for S.C or I.M Injection**

Each 0.5-mL dose of vaccine contains 25 micrograms of each of the following pneumococcal polysaccharide types: 1, 2, 3, 4, 5, 6B, 7F, 8, 9N, 9V, 10A, 11A, 12F, 14, 15B, 17F, 18C, 19F, 19A, 20, 22F, 23F, and 33F.

### **1 THERAPEUTIC INDICATIONS**

PNEUMOVAX 23 is indicated for vaccination against pneumococcal disease caused by those pneumococcal types included in the vaccine.

Effectiveness of the vaccine in the prevention of pneumococcal pneumonia and pneumococcal bacteremia has been demonstrated in controlled trials in South Africa, France and in case-control studies.

PNEUMOVAX 23 will not prevent disease caused by capsular types of pneumococcus other than those contained in the vaccine.

Vaccination with PNEUMOVAX 23 is recommended for selected individuals as follows:

- routine vaccination for persons 50 years of age or older
- persons aged  $\geq 2$  years with certain chronic conditions or in special environments or social settings.

**Specific recommendations for the prevention of pneumococcal disease can be found at the following address: [https://www.health.gov.il/UnitsOffice/HD/PH/epidemiology/td/docs/tadrich\\_Chisunim.pdf](https://www.health.gov.il/UnitsOffice/HD/PH/epidemiology/td/docs/tadrich_Chisunim.pdf)**

### **2 DOSAGE AND ADMINISTRATION**

For intramuscular or subcutaneous injection only.

#### **2.1 Preparation**

- Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration. If either of these two conditions exists, the vaccine should not be administered.
- Do not mix PNEUMOVAX 23 with other vaccines in the same syringe.
- Use a separate sterile syringe and needle for each individual patient to prevent transmission of infectious agents from one person to another.

The package contains 2 needles. Attach the sterile needle to the prefilled syringe by twisting in a clockwise direction until the needle fits securely on the syringe.

#### **2.2 Administration**

Administer PNEUMOVAX 23 intramuscularly or subcutaneously into the deltoid muscle or lateral mid-thigh. Do not inject intravascularly or intradermally.

Administer the entire contents of the single-dose, prefilled syringe per standard protocol using a sterile needle. Discard syringe after use.

#### **2.3 Revaccination**

The Advisory Committee on Immunization Practices (ACIP) has recommendations for revaccination against pneumococcal disease for persons at high risk who were previously vaccinated with PNEUMOVAX 23. Routine revaccination of immunocompetent persons previously vaccinated with a 23-valent vaccine, is not recommended.

### **3 DOSAGE FORMS AND STRENGTHS**

PNEUMOVAX 23 is a clear, sterile solution supplied in a (0.5-mL dose) single-dose prefilled syringe [See *Description (9) and How Supplied/Storage and Handling (13).*]

### **4 CONTRAINDICATIONS**

#### **4.1 Hypersensitivity**

Do not administer PNEUMOVAX 23 to individuals with a history of anaphylactic/anaphylactoid or severe allergic reaction to any component of the vaccine [See *Description (9)*].

## **5 WARNINGS AND PRECAUTIONS**

### **5.1 Persons with Moderate or Severe Acute Illness**

Defer vaccination with PNEUMOVAX 23 in persons with moderate or severe acute illness.

### **5.2 Persons with Severely Compromised Cardiovascular or Pulmonary Function**

Caution and appropriate care should be exercised in administering PNEUMOVAX 23 to individuals with severely compromised cardiovascular and/or pulmonary function in whom a systemic reaction would pose a significant risk.

### **5.3 Use of Antibiotic Prophylaxis**

This vaccine does not replace the need for penicillin (or other antibiotic) prophylaxis against pneumococcal infection. In patients who require penicillin (or other antibiotic) prophylaxis against pneumococcal infection, such prophylaxis should not be discontinued after vaccination with PNEUMOVAX 23.

### **5.4 Persons with Altered Immunocompetence**

Persons who are immunocompromised, including persons receiving immunosuppressive therapy, may have a diminished immune response to PNEUMOVAX 23. [See *Use in Specific Populations* (8.6).]

### **5.5 Persons with Chronic Cerebrospinal Fluid Leakage**

PNEUMOVAX 23 may not be effective in preventing pneumococcal meningitis in patients who have chronic cerebrospinal fluid (CSF) leakage resulting from congenital lesions, skull fractures, or neurosurgical procedures.

### **5.6 Sodium**

This medicinal product contains less than 1 mmol (23 mg) sodium per dosage unit and is considered to be essentially sodium-free.

## **6 ADVERSE REACTIONS**

The most common adverse reactions, reported in >10% of subjects vaccinated with PNEUMOVAX 23 for the first time in a clinical trial, were: injection-site pain/soreness/tenderness (60.0%), injection-site swelling/induration (20.3%), headache (17.6%), injection-site erythema (16.4%), asthenia/fatigue (13.2%), and myalgia (11.9%). [See *Adverse Reactions* (6.1).]

### **6.1 Clinical Trials Experience**

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a vaccine cannot be directly compared to rates in the clinical trials of another vaccine and may not reflect the rates observed in practice.

#### **Primary Vaccination and Revaccination with PNEUMOVAX 23 in Adults 50 Years of Age or Older**

In a randomized, double-blind, placebo-controlled crossover clinical trial, subjects were enrolled in four different cohorts defined by age (50-64 years of age and ≥65 years of age) and vaccination status (no pneumococcal vaccination or receipt of a pneumococcal polysaccharide vaccine 3-5 years prior to the study). Subjects in each cohort were randomized to receive intramuscular injections of PNEUMOVAX 23 followed by placebo (saline containing 0.25% phenol), or placebo followed by PNEUMOVAX 23, at 30-day (±7 days) intervals. The safety of an initial vaccination (first dose) was compared to revaccination (second dose) with PNEUMOVAX 23 for 14 days following each vaccination.

All 1008 subjects (average age, 67 years; 49% male and 51% female; 91% Caucasian, 4.7% African-American, 3.5% Hispanic, and 0.8% Other) received placebo injections.

Initial vaccination was evaluated in a total of 444 subjects (average age 65 years; 32% male and 68% female; 93% Caucasian, 3.2% African-American, 3.4% Hispanic, and 1.1% Other).

Revaccination was evaluated in 564 subjects (average age 69 years; 53% male and 47% female; 90% Caucasian, 3.5% Hispanic, 6.0% African-American, and 0.5% Other).

#### **Serious Adverse Experiences**

In this study, 10 subjects had serious adverse experiences within 14 days of vaccination: 6 who received PNEUMOVAX 23 and 4 who received placebo. Serious adverse experiences within 14 days after PNEUMOVAX 23 included angina pectoris, heart failure, chest pain, ulcerative colitis, depression, and headache/tremor/stiffness/sweating. Serious adverse experiences within 14 days after placebo included

myocardial infarction complicated with heart failure, alcohol intoxication, angina pectoris, and edema/urinary retention/heart failure/diabetes.

Five subjects reported serious adverse experiences that occurred outside the 14-day follow-up window: 3 who received PNEUMOVAX 23 and 2 who received placebo. Serious adverse experiences after PNEUMOVAX 23 included cerebrovascular accident, lumbar radiculopathy, and pancreatitis/myocardial infarction resulting in death. Serious adverse experiences after placebo included heart failure and motor vehicle accident resulting in death.

### Solicited and Unsolicited Reactions

Table 1 presents the adverse event rates for all solicited and unsolicited reactions reported in  $\geq 1\%$  in any group in this study, without regard to causality.

The most common local adverse reactions reported at the injection site after initial vaccination with PNEUMOVAX 23 were pain/tenderness/soreness (60.0%), swelling/induration (20.3%), and erythema (16.4%). The most common systemic adverse experiences were headache (17.6%), asthenia/fatigue (13.2%), and myalgia (11.9%).

The most common local adverse reactions reported at the injection site after revaccination with PNEUMOVAX 23 were pain/soreness/tenderness (77.2%), swelling (39.8%), and erythema (34.5%). The most common systemic adverse reactions with revaccination were headache (18.1%), asthenia/fatigue (17.9%), and myalgia (17.3%). All of these adverse reactions were reported at a rate lower than 10% after receiving a placebo injection.

<b>Table 1: Incidence of Injection-Site and Systemic Complaints in Adults <math>\geq 50</math> Years of Age Receiving Their First (Initial) or Second (Revaccination) Dose of PNEUMOVAX 23 (Pneumococcal Polysaccharide Vaccine, 23 Valent) or Placebo Occurring at <math>\geq 1\%</math> in Any Group</b>			
	<b>PNEUMOVAX 23 Initial Vaccination N=444</b>	<b>PNEUMOVAX 23 Revaccination* N=564</b>	<b>Placebo Injection† N=1008</b>
<b>Number Followed for Safety</b>	438	548	984†
	AE Rate	AE Rate	AE Rate
<b>Injection-Site Complaints</b>			
<b>Solicited Events</b>			
Pain/Soreness/Tenderness	60.0%	77.2%	7.7%
Swelling/Induration	20.3%	39.8%	2.8%
Erythema	16.4%	34.5%	3.3%
<b>Unsolicited Events</b>			
Ecchymosis	0%	1.1%	0.3%
Pruritus	0.2%	1.6%	0.0%
<b>Systemic Complaints</b>			
<b>Solicited Events</b>			
Asthenia/Fatigue	13.2%	17.9%	6.7%
Chills	2.7%	7.8%	1.8%
Myalgia	11.9%	17.3%	3.3%
Headache	17.6%	18.1%	8.9%
<b>Unsolicited Events</b>			
Fever§	1.4%	2.0%	0.7%
Diarrhea	1.1%	0.7%	0.5%
Dyspepsia	1.1%	1.1%	0.9%
Nausea	1.8%	1.8%	0.9%
Back Pain	0.9%	0.9%	1.0%
Neck Pain	0.7%	1.5%	0.2%
Upper Respiratory Infection	1.8%	2.6%	1.8%
Pharyngitis	1.1%	0.4%	1.3%

\*Subjects receiving their second dose of pneumococcal polysaccharide vaccine as PNEUMOVAX 23 approximately 3-5 years after their first dose.

†Subjects receiving placebo injection from this study combined over periods.

‡The number of subjects receiving placebo followed for injection-site complaints. The corresponding number of subjects followed for systemic complaints was 981.

§Fever events include subjects who felt feverish in addition to subjects with elevated temperature.

In this clinical study an increased rate of local reactions was observed with revaccination at 3-5 years following initial vaccination.

For subjects aged 65 years or older, injection-site adverse reaction rate was higher following revaccination (79.3%) than following initial vaccination (52.9%). The proportion of subjects reporting injection site discomfort that interfered with or prevented usual activity or injection site induration  $\geq 4$  inches was higher following revaccination (30.6%) than following initial vaccination (10.4%). Injection site reactions typically resolved by 5 days following vaccination.

For subjects aged 50-64 years, the injection-site adverse reaction rate for revaccinees and initial vaccinees was similar (79.6% and 72.8% respectively).

The rate of systemic adverse reactions was similar among both initial vaccinees and revaccinees within each age group. The rate of vaccine-related systemic adverse reactions was higher following revaccination (33.1%) than following initial vaccination (21.7%) in subjects 65 years of age or older, and was similar following revaccination (37.5%) and initial vaccination (35.5%) in subjects 50-64 years of age. The most common systemic adverse reactions reported after PNEUMOVAX 23 were as follows: asthenia/fatigue, myalgia and headache.

Regardless of age, the observed increase in post vaccination use of analgesics ( $\leq 13\%$  in the revaccinees and  $\leq 4\%$  in the initial vaccinees) returned to baseline by day 5.

### **Sequential Administration of Prevnar 13 and PNEUMOVAX 23**

In a randomized, double-blind, placebo-controlled, multicenter study, healthy adults, 50 years of age and older, received Prevnar 13 followed by PNEUMOVAX 23 either 8 weeks later (Group 1) or 26 weeks later (Group 2). Placebo was administered instead of PNEUMOVAX 23 at 26 weeks (Group 1) or 8 weeks (Group 2). Solicited injection site adverse reactions were evaluated during Days 1 through 5 postvaccination. Solicited systemic adverse reactions and any other adverse reactions were evaluated during Days 1 through 14 postvaccination, and any serious adverse events (SAEs) were collected throughout the study period (through Week 30). [See *Clinical Studies* (11.2).]

Overall, subjects were a mean age of 64.2 years (range: 50 to 97 years). There were more females (n=219, 54.8%) than males (n=181, 45.3%). By race, 84.8% of subjects were White, 9.3% were Black or African-American, and 6.1% were other racial groups; the majority of subjects were not Hispanic or Latino (n=322, 80.5%).

#### **Serious Adverse Reactions**

There were 24 SAEs reported in 20 subjects (n=9 [4.5%] Group 1; n=11 [5.5%] Group 2). No SAEs were considered related to vaccination.

#### **Solicited Adverse Reactions**

Solicited injection site adverse reactions that occurred during Days 1 through 5 postvaccination with PNEUMOVAX 23, solicited systemic adverse reactions that occurred during Days 1 through 14, and fever that occurred during Days 1 through 5 postvaccination with PNEUMOVAX 23 are presented in Table 2. In this study, 81.4% of subjects in Group 1 and 64.0% of subjects in Group 2 reported at least 1 injection site adverse reaction from Days 1 through 5 postvaccination with PNEUMOVAX 23, and 64.9% of subjects in Group 1 and 54.9% of subjects in Group 2 reported at least 1 systemic adverse reaction from Days 1 through 14 postvaccination with PNEUMOVAX 23.

**Table 2: Rates (%) of Solicited Injection Site Reactions Occurring on Days 1 to 5 After PNEUMOVAX 23 and Solicited Systemic Adverse Reactions Occurring on Days 1 to 14 After PNEUMOVAX 23**

	Group 1* (Pprevnar 13 -> PNEUMOVAX 23 -> Placebo)		Group 2† (Pprevnar 13 -> Placebo -> PNEUMOVAX 23)	
	n	(%)	n	(%)
<b>Injection Site Adverse Reactions</b>				
Subjects in population with follow-up	188		164	
Any injection site reaction	153	(81.4)	105	(64.0)
Any Injection site pain‡	149	(79.3)	105	(64.0)
Mild	72	(38.3)	65	(39.6)
Moderate	65	(34.6)	36	(22.0)
Severe§	12	(6.4)	4	(2.4)
Any Injection site swelling	95	(50.5)	48	(29.3)
0 to <2.5 cm	28	(14.9)	19	(11.6)
≥2.5 to <5.1 cm	20	(10.6)	9	(5.5)
≥5.1 to <7.6 cm	20	(10.6)	10	(6.1)
≥7.6 to <10.2 cm	15	(8.0)	2	(1.2)
≥10.2 cm§	12	(6.4)	8	(4.9)
Any Injection site erythema	78	(41.5)	48	(29.3)
0 to <2.5 cm	26	(13.8)	20	(12.2)
≥2.5 to <5.1 cm	12	(6.4)	13	(7.9)
≥5.1 to <7.6 cm	12	(6.4)	6	(3.7)
≥7.6 to <10.2 cm	7	(3.7)	3	(1.8)
≥10.2 cm	19	(10.1)	6	(3.7)
Unknown [missing data]	2	(1.1)	0	(0.0)
<b>Systemic Adverse Reactions</b>				
Subjects in population with follow-up	188		164	
Any systemic adverse reaction	122	(64.9)	90	(54.9)
Myalgia	93	(49.5)	70	(42.7)
Fatigue	59	(31.4)	45	(27.4)
Headache	46	(24.5)	30	(18.3)
Arthralgia	37	(19.7)	25	(15.2)
Subjects with temperature data¶	185		161	
Temperature ≥ 100.4°F	1	(0.5)	0	(0.0)

Every subject is counted a single time for each applicable row and column.

A specific adverse reaction appears in this table only if its incidence in one or more of the columns meets the incidence criterion in the table title, after rounding.

\*Group 1: 8-week interval between Pprevnar 13 and PNEUMOVAX 23.

†Group 2: 26-week interval between Pprevnar 13 and PNEUMOVAX 23.

‡Pain was characterized as mild, moderate or severe. (Mild: awareness of sign or symptom, but easily tolerated. Moderate: discomfort enough to cause interference with usual activity. Severe: incapacitating with inability to work or do usual activity).

§One Group 1 subject with severe pain and swelling greater than 10.2 cm after receipt of PNEUMOVAX 23, went to the Emergency Room for medical attention.

¶Percentages are calculated based on number of subjects with temperature data. Oral temperature was solicited on Days 1 to 5 after PNEUMOVAX 23 vaccination.

### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product.

Any suspected adverse events should be reported to the Ministry of Health according to the National Regulation by using an online form:

<https://sideeffects.health.gov.il>

### 6.2 Post-Marketing Experience

The following list of adverse reactions includes those identified during post approval use of PNEUMOVAX 23. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or their causal relationship to product exposure.

#### *General disorders and administration site conditions*

- Cellulitis
- Malaise
- Fever (>38.9°C)

- Warmth at the injection site
- Decreased limb mobility
- Peripheral edema in the injected extremity
- Injection-site necrosis
- Digestive System*
  - Nausea
  - Vomiting
- Hematologic/ Lymphatic*
  - Lymphadenitis
  - Lymphadenopathy
  - Thrombocytopenia in patients with stabilized idiopathic thrombocytopenic purpura
  - Hemolytic anemia in patients who have had other hematologic disorders
  - Leukocytosis
- Hypersensitivity reactions including*
  - Anaphylactoid reactions
  - Serum Sickness
  - Angioneurotic edema
- Musculoskeletal System*
  - Arthralgia
  - Arthritis
- Nervous System*
  - Paresthesia
  - Radiculoneuropathy
  - Guillain-Barré syndrome
  - Febrile convulsion
- Skin*
  - Rash
  - Urticaria
  - Cellulitis-like reactions
  - Erythema multiforme
- Investigations*
  - Increased serum C-reactive protein

## **7 DRUG INTERACTIONS**

### **7.1 Concomitant Administration with Other Vaccines**

In a randomized clinical study, a reduced immune response to ZOSTAVAX® as measured by gpELISA was observed in individuals who received concurrent administration of PNEUMOVAX 23 and ZOSTAVAX compared with individuals who received these vaccines 4 weeks apart. Consider administration of the two vaccines separated by at least 4 weeks. [See *Clinical Studies (11.3).*]

Limited safety and immunogenicity data from clinical trials are available on the concurrent administration of PNEUMOVAX 23 and vaccines other than ZOSTAVAX.

## **8 USE IN SPECIFIC POPULATIONS**

### **8.1 Pregnancy**

#### Risk Summary

All pregnancies have a background risk of birth defect, loss, or other adverse outcomes.

Available human data from clinical trials of PNEUMOVAX 23 in pregnancy have not established the presence or absence of a vaccine-associated risk.

Developmental toxicity studies have not been conducted with PNEUMOVAX 23 in animals.

### **8.2 Lactation**

#### Risk Summary

It is not known whether PNEUMOVAX 23 is excreted in human milk. Data are not available to assess the effects of PNEUMOVAX 23 on the breastfed infant or on milk production/excretion.

The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for PNEUMOVAX 23 and any potential adverse effects on the breastfed child from PNEUMOVAX 23 or from the underlying maternal condition. For preventive vaccines, the underlying maternal condition is susceptibility to the disease prevented by the vaccine.

#### **8.4 Pediatric Use**

PNEUMOVAX 23 is not approved for use in children less than 2 years of age. Children in this age group do not develop an effective immune response to the capsular types contained in this polysaccharide vaccine. The ACIP has recommendations for use of PNEUMOVAX 23 in children 2 years of age or older, who have previously received pneumococcal vaccines, and who are at increased risk for pneumococcal disease.

#### **8.5 Geriatric Use**

In one clinical trial of PNEUMOVAX 23, conducted post-licensure, a total of 629 subjects who were aged  $\geq 65$  years and 201 subjects who were aged  $\geq 75$  years were enrolled.

In this trial, the safety of PNEUMOVAX 23 in adults 65 years of age and older (N=629) was compared to the safety of PNEUMOVAX 23 in adults 50 to 64 years of age (N=379). The subjects in this study had underlying chronic illness but were in stable condition; at least 1 medical condition at enrollment was reported by 86.3% of subjects who were 50 to 64 years old, and by 96.7% of subjects who were 65 to 91 years old. The rate of vaccine-related systemic adverse experiences was higher following revaccination (33.1%) than following primary vaccination (21.7%) in subjects  $\geq 65$  years of age, and was similar following revaccination (37.5%) and primary vaccination (35.5%) in subjects 50 to 64 years of age.

Since elderly individuals may not tolerate medical interventions as well as younger individuals, a higher frequency and/or a greater severity of reactions in some older individuals cannot be ruled out.

Post-marketing reports have been received in which some elderly individuals had severe adverse experiences and a complicated clinical course following vaccination. Some individuals with underlying medical conditions of varying severity experienced local reactions and fever associated with clinical deterioration requiring hospital care.

#### **8.6 Immunocompromised Individuals**

Persons who are immunocompromised, including persons receiving immunosuppressive therapy, may have a diminished immune response to PNEUMOVAX 23.

### **9 DESCRIPTION**

PNEUMOVAX 23 (Pneumococcal Vaccine Polyvalent) is a sterile, liquid vaccine consisting of a mixture of purified capsular polysaccharides from *Streptococcus pneumoniae* types (1, 2, 3, 4, 5, 6B, 7F, 8, 9N, 9V, 10A, 11A, 12F, 14, 15B, 17F, 18C, 19F, 19A, 20, 22F, 23F, and 33F).

PNEUMOVAX 23 is a clear, colorless solution. Each 0.5-mL dose of vaccine contains 25 micrograms of each polysaccharide type. Also contains 0.25% phenol as a preservative, sodium chloride and water for injections. The vaccine is used directly as supplied. No dilution or reconstitution is necessary.

Syringe plunger stopper and syringe tip cap are not made with natural rubber latex.

### **10 CLINICAL PHARMACOLOGY**

#### **10.1 Mechanism of Action**

PNEUMOVAX 23 induces type-specific antibodies that enhance opsonization, phagocytosis, and killing of pneumococci by leukocytes and other phagocytic cells. The levels of antibodies that correlate with protection against pneumococcal disease have not been clearly defined.

### **11 CLINICAL STUDIES**

#### **11.1 Effectiveness**

The protective efficacy of pneumococcal vaccines containing six (types 1, 2, 4, 8, 12F, and 25) or twelve (types 1, 2, 3, 4, 6A, 8, 9N, 12F, 25, 7F, 18C, and 46) capsular polysaccharides was investigated in two controlled studies in South Africa in male novice gold miners ranging in age from 16 to 58 years, in whom there was a high attack rate for pneumococcal pneumonia and bacteremia. In both studies, participants in the control groups received either meningococcal polysaccharide serogroup A vaccine or saline placebo. In both studies, attack rates for vaccine type pneumococcal pneumonia were observed for the period from 2 weeks through about 1 year after vaccination. Protective efficacy was 76% and 92%, respectively, for the 6- and 12-valent vaccines, for the capsular types represented.

Three similar studies in South African young adult male novice gold miners were carried out by Dr. R. Austrian and associates using similar pneumococcal vaccines prepared for the National Institute of Allergy and Infectious Diseases, with pneumococcal vaccines containing a 6-valent formulation (types 1, 3, 4, 7, 8, and 12) or a 13-valent formulation (types 1, 2, 3, 4, 6, 7, 8, 9, 12, 14, 18, 19, and 25) capsular

polysaccharides. The reduction in pneumococcal pneumonia caused by the capsular types contained in the vaccines was 79%. Reduction in type-specific pneumococcal bacteremia was 82%.

A prospective study in France found a pneumococcal vaccine containing fourteen (types 1, 2, 3, 4, 6A, 7F, 8, 9N, 12F, 14, 18C, 19F, 23F, and 25) capsular polysaccharides to be 77% (95%CI: 51% to 89%) effective in reducing the incidence of pneumonia among male and female nursing home residents with a mean age of 74 (standard deviation of 4 years).

In a study using a pneumococcal vaccine containing eight (types 1, 3, 6, 7, 14, 18, 19, and 23) capsular polysaccharides, vaccinated children and young adults aged 2 to 25 years who had sickle cell disease, congenital asplenia, or undergone a splenectomy experienced significantly less bacteremic pneumococcal disease than patients who were not vaccinated.

In the United States, one post-licensure randomized controlled trial, in the elderly or patients with chronic medical conditions who received a 14-valent pneumococcal polysaccharide vaccine (types 1, 2, 3, 4, 6A, 8, 9N, 12F, 14, 19F, 23F, 25, 7F, and 18C), did not support the efficacy of the vaccine for nonbacteremic pneumonia.

A retrospective cohort analysis study based on the U.S. Centers for Disease Control and Prevention (CDC) pneumococcal surveillance system, showed 57% (95%CI: 45% to 66%) overall protective effectiveness against invasive infections caused by serotypes included in PNEUMOVAX 23 in persons  $\geq 6$  years of age, 65 to 84% effectiveness among specific patient groups (e.g., persons with diabetes mellitus, coronary vascular disease, congestive heart failure, chronic pulmonary disease, and anatomic asplenia) and 75% (95%CI: 57% to 85%) effectiveness in immunocompetent persons aged  $\geq 65$  years of age. Vaccine effectiveness could not be confirmed for certain groups of immunocompromised patients.

## **11.2 Immunogenicity**

The levels of antibodies that correlate with protection against pneumococcal disease have not been clearly defined.

Antibody responses to most pneumococcal capsular types are generally low or inconsistent in children less than 2 years of age.

### **Sequential Administration of Prevnar 13 and PNEUMOVAX 23**

In a randomized, double-blind, placebo-controlled, multicenter study, healthy adults, 50 years of age and older, received Prevnar 13 followed by PNEUMOVAX 23 either 8 weeks later (Group 1) or 26 weeks later (Group 2). Four hundred subjects were randomized 1:1 into Group 1 or Group 2, all of whom were initially vaccinated with Prevnar 13; of these, 188 subjects received PNEUMOVAX 23 (Group 1) and 185 subjects received placebo (Group 2) at Week 8, and 172 subjects received placebo (Group 1) and 164 subjects received PNEUMOVAX 23 (Group 2) at Week 26.

Opsonophagocytic activity (OPA) titers were measured at prevaccination, at Week 12 and at Week 30 for the 12 shared serotypes contained in both PNEUMOVAX 23 and Prevnar 13 (1, 3, 4, 5, 6B, 7F, 9V, 14, 18C, 19A, 19F, and 23F), 2 of the 11 serotypes unique to PNEUMOVAX 23 (22F and 33F), and 1 serotype unique to Prevnar 13 (6A). OPA testing was performed on evaluable serum samples from all subjects at baseline (Day 1) and Week 12, and on sera from a random subset of subjects (approximately 50% of subjects) at Week 30. Estimated GMTs, GMT ratio, and 95% confidence intervals were obtained from a constrained Longitudinal Data Analysis model<sup>1</sup>.

For each of the shared serotypes, Week 12 OPA geometric mean titers (GMTs) in Group 1 were noninferior to those of Group 2, as the lower bounds of the 95% CIs for the OPA GMT ratios were  $>0.5$  for all 12 shared serotypes. For serotypes 22F and 33F, OPA GMTs in Group 1 at Week 12 were superior to those of Group 2 at Week 12, as the lower bounds of the 95% CIs for the OPA GMT ratios were  $>2.0$  for both serotypes.

The OPA GMTs to the 12 shared serotypes and 2 unique serotypes (22F and 33F) when measured 4 weeks after dosing with PNEUMOVAX 23 were generally similar between Group 1 (Week 12) and Group 2 (Week 30 subset).

### **11.3 Concomitant Administration with Other Vaccines**

In a double-blind, controlled clinical trial, 473 adults, 60 years of age or older, were randomized to receive ZOSTAVAX and PNEUMOVAX 23 concomitantly (N=237), or PNEUMOVAX 23 alone followed 4 weeks later by ZOSTAVAX alone (N=236). At four weeks postvaccination, the varicella-zoster virus (VZV) antibody levels following concomitant use were significantly lower than the VZV antibody levels following nonconcomitant administration (GMTs of 338 vs. 484 gpELISA units/mL, respectively; GMT ratio = 0.70 (95% CI: [0.61, 0.80])).

Limited safety and immunogenicity data from clinical trials are available on the concurrent administration of PNEUMOVAX 23 and vaccines other than ZOSTAVAX.

## 12 REFERENCES

1. Liang KY, Zeger S. Longitudinal data analysis of continuous and discrete responses for pre-post designs. *Sankhyā: The Indian Journal of Statistics (Series B)* 2000; 62: 134-148.

## 13 HOW SUPPLIED/STORAGE AND HANDLING

PNEUMOVAX 23 is supplied as follows:

A carton with 1 or 10 single-dose, pre-filled Luer-Lok™ syringe with 2 needles.

Not all pack sizes may be marketed.

### Storage and Handling

- Store at 2-8°C. Do not freeze.
- The expiry date of the product is indicated on the packaging materials.
- All vaccine must be discarded after the expiration date.
- Syringe plunger stopper and syringe tip cap are not made with natural rubber latex.

**Registration Number:** 136 52 23874

**License holder and importer:** Merck Sharp & Dohme (Israel-1996) Company Ltd., 34 Ha'charash St., Hod-Hasharon.

Revised in 03/2025.