

The format of this leaflet was set by the Ministry of Health who checked and approved its content.

**PATIENT PACKAGE INSERT IN ACCORDANCE WITH THE
PHARMACISTS' REGULATIONS (PREPARATIONS), 1986**

Prescription only medicine.

Read this package insert carefully in its entirety before using this medicine.

PHENIMIXIN, SOLUTION

Composition: Chloramphenicol 0.2%,
Polymixin B Sulphate 2500 u/ml.

Inactive Ingredients: PEG 300, Polyoxyl 40 Stearate, Boric Acid, Sodium Borate, Disodium Edetate, Benzalkonium Chloride, Purified Water.

Therapeutic group: antibiotics.

Therapeutic activity: For treatment of eye infections caused by organisms susceptible to the product.

When should the preparation not be used?

Do not use this medicine if you are pregnant or breastfeeding.

Do not use this medicine if you are sensitive to any of its ingredients.

Do not use this medicine without consulting a doctor before starting treatment if you are suffering, or have suffered in the past from impaired function of: the liver, the blood system (e.g. anemia, bone marrow disease).

Warnings: During treatment with this medicine, blood tests should be performed. If you are sensitive to any type of food or medicine, inform your doctor before commencing treatment with this medicine. The medicine contains the preservative Benzalkonium Chloride that can be absorbed by soft contact lenses. Do not use this medicine when soft contact lenses are inserted in your eyes. Remove contact lenses before using this medicine and reinsert into your eyes not less than 15 minutes after use of the medicine.

Drug Interactions: If you are taking another drug concomitantly or if you have just finished treatment with another medicine, inform the attending doctor, in order to prevent hazards or lack of efficacy arising from drug interactions. This is especially important for medicines belonging to the following groups: anticoagulants or other ophthalmic preparations especially those containing steroids.

Side Effects: In addition to the desired effect of the medicine, adverse reactions may occur during the course of taking this medicine.

Effects that require special attention: In the case of local sensitivity reactions like: irritation or burning in the eye, red or swollen eyes (which were not before using the medicine) or bone marrow depression, aplastic anemia with symptoms like fever, throat pain, paleness, abnormal bleeding, weakness (rare), stop treatment and refer to doctor. In the event that you experience side effects not mentioned in this leaflet, or if there is a change in your general health, consult your doctor immediately.

Adverse reactions and drug interactions in children and infants: parents must inform the attending doctor about any side effect as well as any additional medicine being taken by the child. See above for details of special side effects and drug interactions.

Dosage: Dosage is according to doctor's instructions only.

Do not exceed the recommended dosage. If there is no improvement in your condition within a few days, refer to your doctor. This medicine is to be taken at specific time intervals as determined by the attending doctor.

Attention: Do not swallow! This medicine is intended for external use only.

Directions for use: Wash your hands thoroughly. Tilt your head back or lie down on a bed and with the aid of a finger pull the lower eyelid slightly away from the eye. Instill the medicine into the gap, and keep your eyes closed for 1-2 minutes. Try not to blink. To prevent contamination of the eye drops, take care not to allow the tip of the bottle to come into contact with any surface including the finger and the eye itself, and close the bottle tightly after use. Do not use this medicine for more than one month after the bottle is first opened.

How can you contribute to the success of the treatment?

Complete the full course of treatment as instructed by the doctor. Even if there is an improvement in your health, do not discontinue use this medicine, before consulting your doctor.

Avoid poisoning! This medicine, and all other medicines, must be stored in a safe place out of the reach of children and/or infants, to avoid poisoning. If you have taken an overdose, or if a child has accidentally swallowed the medicine, proceed immediately to a hospital emergency room and bring the package of the medicine with you. Do not induce vomiting unless explicitly instructed to do so by a doctor! This medicine has been prescribed for the treatment of your ailment, in another patient, it may cause harm. Do not give this medicine to your relatives, neighbours or acquaintances. Do not take medicines in the dark! Check the label and the dose each time you take your medicine. Wear glasses if you need them.

Storage: Store In a refrigerator (2-8°C).

Even if kept in their original container and stored as recommended, medicines may be kept for a limited period only. Please note the expiry date of the medicine! In case of doubt, consult the pharmacist who dispensed the medicine to you. Do not store different medications in the same package.

Licence number: 1004628582

Manufacturer:

Vitamed Pharmaceutical Industries Ltd., P.O.B. 114, Binyamina 3055002, Israel.