

SUMMARY OF PRODUCT CHARACTERISTICS

1 NAME OF THE MEDICINAL PRODUCT

Undecyl Ointment

2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Each tube contains:

Zinc Undecylenate 21%

Undecylenic Acid 2%

Excipients with known effect:

Lanolin, Cetyl Alcohol and Stearyl Alcohol.

For the full list of excipients, see section 6.1

3 PHARMACEUTICAL FORM

A slightly yellow homogeneous ointment.

4 CLINICAL PARTICULARS

4.1 Therapeutic indications

Prophylaxis and treatment of mycotic skin infections particularly athlete's foot.

4.2 Posology and method of administration

Route of administration: Topical administration.

The medicine is not intended for children and infants under 2 years of age, unless recommended and supervised by a doctor.

Treatment

The usual recommended dosage for adults and children above 2 years of age:

Apply twice daily (morning and evening) on the foot and between the toes after washing and drying thoroughly.

Prevention

The usual recommended dosage for adults and children above 2 years of age:

Apply once daily on the foot and between the toes after washing and drying thoroughly.

Continue to use **Undecyl Ointment** for a week after the symptoms of the fungal infection have disappeared.

If the condition worsens, the patient should contact a doctor.

4.3 Contraindications

Hypersensitivity to the active ingredients (Zinc Undecylenate, Undecylenic Acid) or to any of the excipients listed in section 6.1.

4.4 Special warnings and precautions for use

- The medicine should not be swallowed! This medicine is intended for external use only.
- Contact with the eyes and mucous membranes should be avoided.
- Do not apply to broken skin.
- Treatment should be discontinued if irritation is severe.
- Keep all medicines out of the reach of children.

Excipients warnings:

Undecyl Ointment contains excipients with known effect: Lanolin, Cetyl Alcohol and Stearyl Alcohol, which may cause local skin reactions (e.g., contact dermatitis).

4.5 Interaction with other medicinal products and other forms of interaction

No clinically significant interactions known.

4.6 Pregnancy and lactation

The safety of the medicine during pregnancy and lactation has not been established, but use during these periods is not considered to constitute a hazard.

4.7 Effects on ability to drive and use machines

No or negligible influence.

4.8 Undesirable effects

Hypersensitivity reactions may occur occasionally. Irritation of the skin may rarely occur.

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/ risk balance of the medicinal product.

Any suspected adverse events should be reported to the Ministry of Health according to the National Regulation by using an online form <https://sideeffects.health.gov.il>

4.9 Overdose

Excessive application to the skin is unlikely to cause untoward effects. In the unlikely event of ingestion of the medicine symptoms of overdose may include nausea, vomiting, diarrhea and general gastrointestinal disturbances.

Treatment

Symptomatic.

5 PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

D01AE - Other antifungals for topical use.

Undecylenic Acid and Zinc Undecylenate have anti-fungal and anti-bacterial properties.

5.2 Pharmacokinetic properties

Not applicable.

5.3 Preclinical safety data

There are no preclinical data of relevance to the prescriber which are additional to that already included in other sections of the SPC.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Vaseline, Lanolin Anhydrous (Wool Fat), Purified Water, Mineral Oil Heavy, Cetyl Alcohol, Terpineol, Paraffin Hard (Non-Caking), Super Hartolan (Wool Alcohol), Stearyl Alcohol, Cholesterol.

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

The expiry date of the product is indicated on the packaging materials.

The medicine can be used for up to 28 days after the tube is first opened, and not later than the expiry date that appears on the package.

6.4 Special precautions for storage

Store in a dry place, below 25°C.
Close tightly, to prevent moisture penetration.

6.5 Nature and contents of container

Aluminium tube with HDPE cap.
Each pack contains 25 grams.

6.6 Special precautions for disposal

No special requirements for disposal.

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

7 MARKETING AUTHORISATION HOLDER AND MANUFACTURER

Rekah Pharmaceutical Industry Ltd., 30 Hamelacha St., Holon, 5881904, Israel.

8 MARKETING AUTHORISATION NUMBER(S)

137-46-21090-00

Revised in July 2025 according to MOH guidelines.