

**Patient package insert according to
Pharmacists' Regulations (Preparations)- 1986**
This medicine can be sold without a physician's prescription

UNDECYL POWDER

Each securitainer contains:
Zinc Undecylenate 20%
Undecylenic Acid 2%

Inactive ingredients and allergens in the medicine- see section 6 "Additional information" and in section 2 "Important information about some of the ingredients of this medicine".

Read the entire leaflet carefully before using this medicine. This leaflet contains concise information about the medicine. If you have any further questions, refer to the doctor or pharmacist.

Use this medicine according to the instructions in section 3 "How to use this medicine?" in this leaflet. Consult the pharmacist if you need additional information. Refer to a doctor if your condition worsens.

1. What is the medicine intended for?

For prophylaxis and treatment of mycotic skin infections, particularly athlete's foot.

Therapeutic group: Anti-fungal.

2. Before using this medicine

Do not use this medicine if:

- You are hypersensitive (allergic) to the active ingredients (Zinc Undecylenate, Undecylenic Acid) or to any of the other ingredients this medicine contains (see section 6 "Additional Information").

Special warnings regarding the use of this medicine

- Keep away from eyes and other sensitive areas (such as nose and mouth).
- Do not use this medicine on cracked skin.
- If the irritation is severe, discontinue use.

Children and adolescents:

This medicine is not intended for children and infants under two years of age, except on the recommendation and supervision of a doctor.

Drug interactions:

If you are taking or have recently taken other medicines, including non-prescription medicines and nutritional supplements, tell the doctor or pharmacist.

Pregnancy and breastfeeding:

If you are pregnant, may be pregnant, or are breastfeeding, consult the doctor or pharmacist before using this medicine.

Driving and using machines:

This medicine has no or negligible effect on your ability to drive or operate machinery.

3. How to use this medicine?

Check with the doctor or pharmacist if you are not sure about the dosage and manner of treatment with the medicine.

Treatment

The usual recommended dosage for adults and children above two years of age, unless otherwise instructed by the doctor:

Sprinkle twice daily (morning and evening), after washing and drying thoroughly, on the foot and between the toes. Additionally, it is recommended to sprinkle the powder once daily inside the socks, before wearing them.

Prevention

The usual recommended dosage for adults and children above two years of age, unless otherwise instructed by the doctor:

Sprinkle once daily, after washing and drying thoroughly, on the foot and between the toes. Additionally, it is recommended to sprinkle the powder once daily inside the socks, before wearing them.

Do not exceed the recommended dose.

Duration of treatment

Continue to use this medicine for a week after the symptoms of the fungal infection have disappeared.

If your condition worsens, refer to your doctor.

Method of administration-

Do not swallow! This medicine is intended for external use only.

If you have accidentally taken a higher dosage

If you have taken an overdose or if a child has accidentally swallowed the medicine, proceed immediately to a doctor or a hospital emergency room and bring the package of the medicine with you.

Do not take medicines in the dark! Check the label and the dose each time you take a medicine. Wear glasses if you need them.

If you have further questions on the use of this medicine, consult the doctor or pharmacist.

4. Side effects

As with any medicine, the use of **Undecyl Powder** may cause side effects in some users. Do not be alarmed when reading the list of side effects. You may not suffer from any of them.

Stop taking this medicine and refer to the doctor as soon as possible with the appearance of one or more of the following side effects:

- Skin irritation (redness and itching) or allergic reactions (rash, itching, wheezing).

If a side effect appears, if one of the side effects worsens or if you suffer from a side effect which is not mentioned in this leaflet, consult the doctor.

Report of side effects

Side effects can be reported to the Ministry of Health by clicking on the link "Report adverse effects and problems associated with medications and drugs" on the Ministry of Health home page (www.health.gov.il), which links to an online form for reporting side effects, or by following the link:

<https://sideeffects.health.gov.il>

5. How to store the medicine?

• Avoid poisoning! This medicine and any other medicine must be stored in a closed place out of the reach and sight of children and/ or infants to avoid poisoning. Do not induce vomiting unless explicitly instructed to do so by the doctor.

• Do not use the medicine after the expiry date (exp. Date) that appears on the package. The expiry date refers to the last day of that month.

Storage conditions:

- Store in a dry place, below 25°C.
- The medicine can be used for up to 60 days after the securitainer is first opened and not later than the expiry date, that appears on the package.
- Close tightly, to prevent moisture penetration.
- Do not throw away any medicine via wastewater or household waste. Ask the pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Additional information

In addition to the active ingredient, this medicine also contains:

Talc, Colloidal Silicon Dioxide (Aerosil 200), Terpineol.

What the medicine looks like and what the package contains:

Plastic securitainer containing white powder.

Approved package sizes:

30 grams, 80 grams.

Not all package sizes may be marketed.

Manufacturer and registration holder:

Rekah Pharmaceutical Industry Ltd., 30 Hamelacha St., Holon, 5881904, Israel.

Revised in July 2025 according to MOH guidelines.

Drug registration number at the national drug registry of the Ministry of Health:

029-47-25330-00