

## **STILNOX 10mg tablets**

### **Prescribing Information**

**WARNING: Zolpidem may be associated with potentially dangerous complex sleep-related behaviors which may include sleep walking, sleep driving and other bizarre behaviors. Zolpidem is not to be taken with alcohol. Caution is needed with other CNS depressant drugs. Limit use to four weeks maximum under close medical supervision.**

#### **1 NAME OF THE MEDICINAL PRODUCT**

Stilnox 10 mg Tablets

#### **2 QUALITATIVE AND QUANTITATIVE COMPOSITION**

Each tablet contains zolpidem tartrate 10 mg.

Excipients with known effect:

Each Stilnox tablet contains 90.4 mg lactose monohydrate

For the full list of excipients, *see section 6.1 List of excipients.*

#### **3 PHARMACEUTICAL FORM**

Film-coated tablets.

Stilnox tablets are white to off-white film coated oblong tablets

#### **4 CLINICAL PARTICULARS**

##### **4.1 THERAPEUTIC INDICATIONS**

Zolpidem is indicated for the short-term treatment of occasional or transient insomnia in adults in situations where the insomnia is debilitating or is causing severe distress for the patient.

##### **4.2 DOSE AND METHOD OF ADMINISTRATION**

As with all hypnotics long term use is not recommended and a course of treatment should not exceed 4 weeks.

Hypnotics should generally be limited to 7 to 10 days of use, and reevaluation of the patient is recommended if they are to be taken for more than 2 to 3 weeks.

### **Dosage in Adults**

Use the lowest effective dose for the patient. The recommended initial dose is 5 mg for women and either 5 or 10 mg for men, taken only once per night immediately before bedtime with at least 7-8 hours remaining before the planned time of awakening. If the 5 mg dose is not effective, the dose can be increased to 10 mg. In some patients, the higher morning blood levels following use of the 10 mg dose increase the risk of next day impairment of driving and other activities that require full alertness (*See section Special warnings and precautions for use*). The total dose of Stilnox should not exceed 10 mg once daily immediately before bedtime.

Stilnox should be taken as a single dose and should not be readministered during the same night.

The recommended initial doses for women and men are different because zolpidem clearance is lower in women.

Long-term use of Stilnox is not recommended. Treatment should be as short as possible. Extended treatment should not take place without re-evaluation of the patient's status because the risk of abuse and dependence increases with the duration of treatment (*See section 4.4 Abuse and Dependence*).

### **Special Populations**

Elderly or debilitated patients may be especially sensitive to the effects of zolpidem tartrate. Patients with hepatic insufficiency do not clear the drug as rapidly as normal subjects. The recommended dose of Stilnox in both of these patient populations is 5 mg (half tablet) once daily immediately before bedtime (*See section 4.4 Special warnings and precautions for use; 5.2 Special Populations*).

Stilnox should not be used in patient with severe hepatic insufficiency (*See section 4.3 Contraindications*).

### **Use with CNS Depressants**

Dosage adjustments may be necessary when Stilnox is combined with other CNS depressant drugs because of the potentially additive effects (*See section 4.4 Special warnings and precautions for use*).

### **Administration**

The effect of Stilnox may be slowed by ingestion with or immediately after a meal.

### **Dosage forms and strengths**

Stilnox is available in 10 mg strength tablets for oral administration.

It is possible to take half of tablet (5mg) if needed (*See section 4.2 Special population*).

## **4.3 CONTRAINDICATIONS**

- Sleep apnea.
- Known hypersensitivity to zolpidem or other ingredients in the tablet.
- Myasthenia gravis.
- Severe hepatic insufficiency.
- Acute and/or severe pulmonary insufficiency.
- Prior or concomitant intake with alcohol.
- Stilnox should not be prescribed for children.
- Patients who have previously experienced complex sleep behaviors after taking Stilnox.

#### **4.4 SPECIAL WARNINGS AND PRECAUTIONS FOR USE**

The cause of insomnia should be identified wherever possible and the underlying factors treated before a hypnotic is prescribed.

The failure of insomnia to remit after a 7-14 day course of treatment may indicate the presence of a primary psychiatric or physical disorder, and the patient should be carefully re-evaluated at regular intervals.

#### **Withdrawal, Rebound, Dependence and Tolerance**

##### **Tolerance**

Continuous long-term use of Stilnox is not recommended and should not exceed four weeks. Some loss of efficacy to the hypnotic effects of sedative/hypnotic agents may develop after repeated use for a few weeks.

##### **Dependence**

Use of zolpidem may lead to the development of abuse and/or physical and psychological dependence. The risk of dependence increases with dose and duration of treatment; it is also greater in patients with a history of psychiatric disorders and/or alcohol or drug abuse. Stilnox should be used with extreme caution in patients with current or a history of alcohol or drug abuse. These patients should be under careful surveillance when receiving hypnotics. Once physical dependence has developed, abrupt termination of treatment will be accompanied by withdrawal symptoms. These may consist of headaches or muscle pain, extreme anxiety and tension, restlessness, confusion and irritability. In severe cases the following symptoms may occur: derealisation, depersonalisation, hyperacusis, numbness and tingling of the extremities, hypersensitivity to light, noise and physical contact, hallucinations, delirium or epileptic seizures.

##### **Rebound insomnia**

A transient syndrome whereby the symptoms that led to treatment with sedative/hypnotic agents recur in an enhanced form may occur on withdrawal of hypnotic treatment. It may be accompanied by other reactions including mood changes, anxiety and restlessness.

It is important that the patient should be aware of the possibility of rebound phenomena, thereby minimising anxiety over such symptoms should they occur when the medicinal product is discontinued.

There are indications that, in the case of sedative/hypnotic agents with a short duration of action, withdrawal phenomena can manifest within the dosage interval, especially when the dosage is high.

When Stilnox is used in accordance with the recommendations for dosage, duration of treatment and warnings, the risk of withdrawal symptoms or rebound phenomena occurring is minimal.

### **Severe injuries**

Due to its pharmacological properties, zolpidem can cause drowsiness and a decreased level of consciousness, which may lead to falls and consequently to severe injuries.

### **Patients with Long QT syndrome**

An in vitro cardiac electrophysiological test showed that under experimental conditions, using very high concentration and pluripotent stem cells, zolpidem may reduce the hERG related potassium currents. As a precaution, the benefit/risk ratio of zolpidem treatment in patients with known congenital long QT syndrome should be carefully considered.

### **Chemical submission (Drug facilitated illicit use for criminal intent)**

The rapid onset of sedation, coupled with the amnesic features of Stilnox, particularly when combined with alcohol, administered without knowledge of the victim, has proven to induce incapacitation and thus facilitate criminal actions (which could be dangerous). Healthcare Providers should prescribe Stilnox according to their clinical evaluation and only in case of medical need as it may be used illicitly for chemical submission.

### **CNS effects**

As with all patients taking CNS-depressant medications, patients receiving Stilnox should be warned not to operate dangerous machinery or motor vehicles until it is known that they do not become drowsy or dizzy from Stilnox therapy. Patients should be advised that their tolerance for other CNS depressants will be diminished and that these medications should either be eliminated or given in reduced dosage in the presence of Stilnox. Prior or concomitant intake with alcohol is contraindicated (*See section 4.3 Contraindications*).

### **Respiratory function**

Both animal and human pharmacology studies performed with Stilnox have not observed any effect on the respiratory center. However, as other sedative/hypnotics have the capacity to depress respiratory drive, caution is advised when Stilnox is administered to patients with respiratory insufficiency (*See section 4.3 Contraindications*).

### **Use in hepatic impairment**

As clearance and metabolism of zolpidem is reduced in hepatic impairment, dosage should begin at 5 mg with particular caution being exercised in elderly patients. In adults (under 65 years) dosage may be increased to 10 mg only where the clinical response is inadequate and the drug is well tolerated. (*See Section 4.4 Special warnings and precautions for use 'Use in the elderly' and Section 4.2 Dose and method of administration*). Zolpidem must not be used in patients with severe hepatic impairment as it may contribute to encephalopathy.

### **Use in renal impairment**

Dosage reduction is not necessary in patients with renal impairment, however, as a general precaution, these patients should be monitored closely.

### **Use in the elderly or Debilitated Patient**

Elderly and debilitated patients may be particularly sensitive to the effects of Stilnox, therefore a 5 mg dose is recommended. This dose should not be exceeded in these patients. (*See section 4.2 Dose and method of administration*).

Such patients may be particularly susceptible to the sedative effects of the medication and associated giddiness, ataxias and confusion, which may increase the possibility of a fall.

### **Memory impairment**

Sedative/hypnotic agents may induce anterograde amnesia. The condition occurs most often several hours after ingesting the product and therefore to reduce the risk patients should ensure that they will be able to have an uninterrupted sleep of 7-8 hours.

### **Suicidality, Depression, Psychosis and Schizophrenia**

Several epidemiological studies show an increased incidence of suicide and suicide attempt in patients with or without depression, treated with benzodiazepines and other hypnotics, including zolpidem. Stilnox should be administered with caution in patients exhibiting symptoms of depression. Stilnox is not recommended as primary therapy in patients with psychotic illness, including depression and psychosis. In such conditions, psychiatric assessment and supervision are necessary as depression may increase in some patients and may contribute to deterioration in severely disturbed schizophrenics with confusion and withdrawal. Pre-existing depression may be unmasked during the use of Stilnox. Suicidal tendencies may be present or uncovered and protective measures may be required. Intentional overdosage is more common in this group of patients: therefore, the least amount of drug that is feasible should be prescribed for the patient at any one time.

### **Other Psychiatric and Paradoxical Reactions**

Other psychiatric and paradoxical reactions such as acute rage, restlessness, insomnia exacerbated, agitation, irritability, aggression, delusions, anger, nightmares, hallucinations, stimulation or excitement, abnormal behavior, delirium and other adverse behavioral effects are known to occur when using sedative/hypnotic agents like Stilnox. Should such reactions occur, Stilnox should be discontinued. These reactions are more likely to occur in the elderly.

### **Somnambulism and Associated Behaviors**

Complex sleep behaviors, including sleep-walking, sleep-driving, and engaging in other activities while not fully awake, may occur following the first or any subsequent use of Stilnox. Patients can be seriously injured or injure others during complex sleep behaviors. Such injuries may result in a fatal outcome. Other complex sleep behaviors (e.g., preparing and eating food, making phone calls, or having sex) have also been reported. Patients usually do not remember these events. Postmarketing reports have shown that complex sleep behaviors may occur with Stilnox alone at recommended doses, with or without the concomitant use of alcohol or other central nervous system (CNS) depressants (*See section*

*4.5 Interaction with other medicinal products and other forms of interactions*). Discontinue Stilnox immediately if a patient experiences a complex sleep behavior (*See section 4.3 Contraindications*). The use of alcohol and other CNS depressants with zolpidem appears to increase the risk of such behaviors, as does the use of Stilnox at doses exceeding the maximum recommended dose (*See section 4.5 Interactions with other medicines and other forms of interactions and section 4.8 Adverse Effects (Undesirable effects)*). These events can occur in sedative-hypnotic naive as well as sedative-hypnotic experienced patients.

### **Psychomotor Impairment**

Zolpidem has CNS-depressant effects. The risk of psychomotor impairment, including impaired driving ability, is increased if zolpidem is taken within less than 7-8 hours before performing activities that require mental alertness, a dose higher than the recommended dose is taken, or zolpidem is co-administered with other CNS depressants, alcohol, or with other drugs that increase the blood levels of zolpidem.

### **Interactions with Alcohol**

Prior or concomitant intake with alcohol is contraindicated (*See section 4.3 Contraindications*). Patients should be advised that their tolerance for alcohol and other CNS depressants might be reduced and have an additive effect on psychomotor performance (*See section 4.4 Special warnings and precautions for use 'Somnambulism and Associated Behaviors'* above).

### **Risks from Concomitant use with Opioids**

Concomitant use of sedative-hypnotic drugs, including zolpidem, with opioids may result in sedation, respiratory depression, coma and death. Because of these risks, reserve concomitant prescribing of opioids and zolpidem for use in patients for whom alternative treatment options are inadequate.

If a decision is made to prescribe zolpidem concomitantly with opioids, prescribe the lowest effective dosages and minimum durations of concomitant use, and follow patients closely for signs and symptoms of respiratory depression and sedation. (see Section 4.5 Interactions with other medicines and other forms of interactions).

### **Severe Anaphylactic and Anaphylactoid Reactions**

Rare cases of angioedema involving the tongue, glottis or larynx have been reported in patients after taking the first or subsequent doses of sedative-hypnotics, including zolpidem. Some patients have had additional symptoms such as dyspnea, throat closing, or nausea and vomiting that suggest anaphylaxis. Some patients have required medical therapy in the emergency department. If angioedema involves the tongue, glottis or larynx, airway obstruction may occur and be fatal. Patients who develop angioedema after treatment with zolpidem should not be rechallenged with the drug.

### **Epilepsy**

Abrupt withdrawal of CNS-depressant drugs in persons with convulsive disorders has been associated with a temporary increase in the frequency and or severity of seizures. As with other sedative/hypnotics, caution is advised when Stilnox is used in these patients.

### **Abuse**

Caution must be exercised in administering Stilnox to individuals known to be addiction prone or those whose history suggests they may increase the dosage on their own initiative. It is desirable to limit repeat prescription without adequate medical supervision.

### **Effects on laboratory tests**

No data available.

### **Lactose**

Patients with rare hereditary problems of galactose intolerance, total lactase deficiency or glucose-galactose malabsorption should not take this medicine.

## **4.5 INTERACTIONS WITH OTHER MEDICINES AND OTHER FORMS OF INTERACTIONS**

### **CNS depressants**

Co-administration of Stilnox with other CNS depressants should be exercised with caution since the central depressant effect may be additive. CNS depressants include alcohol, benzodiazepines, barbiturates, sedative/hypnotics, anxiolytics, antidepressant agents (including tricyclic antidepressants, MAOIs), antipsychotics (neuroleptics), phenothiazines, skeletal muscle relaxants, antihistamines, neuroleptics, antiepileptic drugs, narcotic analgesics or anaesthetics. Concomitant use of zolpidem with these drugs may increase drowsiness and psychomotor impairment, including impaired driving ability. In the case of narcotic analgesics enhancement of euphoria may also occur.

### **Opioids**

The concomitant use of sedative-hypnotic drugs, including zolpidem, and opioids increases the risk of sedation, respiratory depression, coma and death because of additive CNS depressant effect. Limit dosage and duration of concomitant use of zolpidem and opioids (*See section 4.4 Special warnings and precautions for use 'Risks from Concomitant use with Opioids'*).

### **Alcohol**

Prior or concomitant intake with alcohol is contraindicated (*See section 4.3 Contraindications*). Patients should be advised that their tolerance for alcohol and other CNS depressants might be reduced and have an additive effect on psychomotor performance. The use of alcohol and other CNS depressants with zolpidem appears to increase the risk of somnambulism and associated behaviors (*See section 4.4 Special warnings and precautions for use 'Somnambulism and Associated Behaviors'*).

### **Imipramine**

The sedative effects of imipramine 75 mg and zolpidem 20 mg were shown to be additive when the two compounds were given concomitantly in healthy volunteers. No pharmacokinetic interaction was shown between zolpidem and imipramine or its metabolite, desipramine.

## **Chlorpromazine**

The combination of zolpidem 10 mg and chlorpromazine 50 mg in healthy volunteers produced an addition of effects seen in psychometric tests and decreased alertness and psychomotor performance. No pharmacokinetic interaction was observed.

## **Haloperidol**

No evidence of pharmacokinetic interaction between zolpidem 20 mg and haloperidol 2 mg was seen when they were given concurrently to healthy volunteers.

## **Caffeine**

No change in the sleep inducing effect of zolpidem was seen when 300 mg caffeine was given in the evening 45 minutes before administration of zolpidem 20 mg to 8 healthy volunteers.

## **Warfarin**

Prothrombin times were not prolonged in healthy adults when zolpidem 20 mg was administered for 4 consecutive nights concomitantly with warfarin. Warfarin had been given for at least 10 days previously to produce a 1.5 times prolongation of baseline prothrombin time in the volunteers. Zolpidem does not appear to modify the anticoagulant activity of warfarin.

## **Digoxin**

The concurrent administration of zolpidem 10 mg once daily and digoxin 0.25 mg in healthy volunteers did not show any alteration of the pharmacokinetic or pharmacodynamic profile of digoxin.

## **H2 - antagonists**

Simultaneous administration of zolpidem 20 mg and cimetidine 200 mg tds and 400 mg at night or ranitidine 150 mg bd did not cause any significant change in psychometric tests from those produced by zolpidem alone. No change in the pharmacokinetics of zolpidem were caused by concomitant administration of either cimetidine or ranitidine.

## **Hepatic enzyme inhibitors and inducers**

Zolpidem is metabolized via several hepatic cytochrome P450 enzymes: the main enzyme being CYP3A4 with the contribution of CYP1A2. Compounds which inhibit or enhance certain hepatic enzymes (particularly cytochrome P450) may increase or decrease the activity of some hypnotics like zolpidem. The pharmacodynamic effect of zolpidem is decreased when it is administered with a CYP3A4 inducer such as rifampicin and St John's Wort. Co-administration of St John's Wort may decrease blood levels of zolpidem, therefore concurrent use is not recommended.

Ketoconazole has a significant but only quantitatively modest reduction in zolpidem clearance, with an increase in its pharmacodynamic effects. Patients should be advised that use of zolpidem with ketoconazole may enhance the sedative effects of zolpidem. However, when zolpidem is administered with itraconazole (a CYP3A4 inhibitor) its pharmacokinetics

and pharmacodynamics were not significantly modified. The clinical relevance of these results is unknown.

Fluvoxamine is a strong inhibitor of CYP1A2 and a moderate to weak inhibitor of CYP2C9 and CYP3A4. Co-administration of fluvoxamine may increase blood levels of zolpidem, concurrent use is not recommended.

Ciprofloxacin has been shown to be a moderate inhibitor of CYP1A2 and CYP3A4. Co-administration of ciprofloxacin may increase blood levels of zolpidem, concurrent use is not recommended.

## **4.6 FERTILITY, PREGNANCY AND LACTATION**

### **Effects on fertility**

No data available.

### **Use in pregnancy - Category B3**

This drug has been taken by only a limited number of pregnant women and women of childbearing age, without an increase in the frequency of malformation or other direct or indirect harmful effects on the human foetus having been observed. Studies in animals have shown evidence of an increased occurrence of foetal damage, the significance of which is uncertain in humans. The use of zolpidem is not recommended during pregnancy.

Cases of reduced fetal movement and fetal heart rate variability have been described after administration of benzodiazepines or other sedative-hypnotic drugs such as zolpidem during pregnancy.

Administration of zolpidem during the late phase of pregnancy or during labor has been associated with effects on the neonate, such as hypothermia, hypotonia, feeding difficulties, and respiratory depression.

If zolpidem is prescribed to a woman of childbearing potential, she should be warned to contact her physician about stopping the product if she intends to become or suspects that she is pregnant.

### **Teratogenic effects**

In reproductive toxicity studies, rats treated with oral zolpidem with estimated exposures (AUC) to zolpidem and its major metabolite of 41 and 15 times, respectively, the anticipated clinical exposure did not exhibit teratogenic effects but post-implantation survival index and postpartum viability of the offspring were significantly reduced. In rats, delayed ossification of foetal skull bones occurred at zolpidem and metabolite exposure levels of 8 and 3 times, respectively, the anticipated clinical exposure.

Rabbits treated with oral zolpidem with estimated exposure to zolpidem of 0.6-2.6 times the anticipated clinical exposure did not exhibit teratogenic effects, but there was increased post-implantation loss.

Although animal studies have not shown any teratogenic effects with zolpidem, the safety of zolpidem in human pregnancy has not been established.

### **Non teratogenic effects**

Cases of severe neonatal respiratory depression have been reported when zolpidem was used with other CNS depressants at the end of pregnancy.

Infants born to mothers who took hypnotics chronically during the latter stages of pregnancy may have developed physical dependence and may be at some risk for developing withdrawal symptoms in the postnatal period. Appropriate monitoring of the newborn in the postnatal period is recommended.

### **Use in lactation**

The use of Stilnox in nursing women is not recommended as small quantities of zolpidem are excreted into breast milk.

## **4.7 EFFECTS ON ABILITY TO DRIVE AND USE MACHINES**

This preparation is to aid sleep. Patients should not drive or operate machinery for 8 hours after taking Stilnox.

Adverse effects including drowsiness, prolonged reaction time, dizziness, sleepiness, blurred/double vision, reduced alertness and/or impaired driving may continue the following day. In order to minimise this risk a full night of sleep (7-8 hours) is recommended. After ingesting the medicine, patients should be cautioned against engaging in hazardous occupations requiring complete mental alertness or motor co-ordination such as operating machinery or driving a motor vehicle, including potential impairment of the performance of such activities that may occur the day following ingestion of Stilnox. Furthermore, the co-administration of zolpidem with alcohol and other CNS depressants increases the risk of such effects. Patients should be warned not to use alcohol or other psychoactive substances when taking zolpidem.

## **4.8 ADVERSE EFFECTS (UNDESIRABLE EFFECTS)**

### **Clinical trials data**

There is evidence of a dose-relationship for adverse effects associated with Stilnox use, particularly for certain CNS events. These occur most frequently in elderly patients.

#### ***Associated with discontinuation of treatment***

Approximately 4% of 1,701 patients who received zolpidem at all doses (1.25 to 90 mg) in US premarketing clinical trials discontinued treatment because of an adverse clinical event. Events most commonly associated with discontinuation from US trials were daytime drowsiness (0.5%), dizziness (0.4%), headache (0.5%), nausea (0.6%), and vomiting (0.5%). Approximately 6% of 1,320 patients who received zolpidem at all doses (5 to 50 mg) in similar European trials discontinued treatment because of an adverse event. Events most commonly associated with discontinuation from these trials were daytime drowsiness (1.6%), amnesia (0.6%), dizziness (0.6%), headache (0.6%) and nausea (0.6%).

#### ***Incidence in controlled clinical trials***

***Most commonly observed adverse events in controlled trials:*** During short-term treatment (up to 10 nights) with Stilnox at doses up to 10 mg, the most commonly observed adverse events associated with the use of STILNOX and seen at statistically significant differences

from placebo-treated patients were drowsiness (reported by 2% of Stilnox patients), dizziness (1%), and diarrhoea (1%). During longer-term treatment (28 to 35 nights) with Stilnox at doses up to 10 mg, the most commonly observed adverse events associated with the use of STILNOX and seen at statistically significant differences from placebo-treated patients were dizziness (5%) and drugged feelings (3%).

**Adverse events observed at an incidence of  $\geq 1\%$  in controlled trials:** The following tables enumerate treatment-emergent adverse event frequencies that were observed at an incidence equal to 1% or greater among patients with insomnia who received Stilnox in US placebo-controlled trials. Events reported by investigators were classified utilising a modified World Health Organisation (WHO) dictionary of preferred terms for the purpose of establishing event frequencies.

The following table was derived from a pool of 11 placebo-controlled short-term US efficacy trials involving Stilnox in doses ranging from 1.25 to 20 mg. The table is limited to data from doses up to and including 10 mg, the highest dose recommended for use.

**Table 1 - Incidence of Treatment-Emergent Adverse Experiences in Short-term Placebo-Controlled Clinical Trials (Percentage of patients reporting)**

Body System/ Adverse Event*	Zolpidem (<10 mg) (N=685)	Placebo (N=473)
<b>Central and Peripheral Nervous System</b>		
Headache	7	6
Drowsiness	2	-
Dizziness	1	-
<b>Gastrointestinal System</b>		
Nausea	2	3
Diarrhoea	1	-
<b>Musculoskeletal System</b>		
Myalgia	1	2

\*Events reported by at least 1% of Stilnox patients are included.

The following table was derived from a pool of three placebo-controlled long-term efficacy trials involving Stilnox (zolpidem tartrate). These trials involved patients with chronic insomnia who were treated for 28 to 35 nights with zolpidem at doses of 5, 10 or 15 mg. The table is limited to data from doses up to and including 10 mg, the highest dose recommended for use. The table includes only adverse events occurring at an incidence of at least 1% for zolpidem patients.

**Table 2 - Incidence of Treatment-Emergent Adverse Experiences in Long-term Placebo-Controlled Clinical Trials (Percentage of patients reporting)**

Body System/ Adverse Event*	Ambien CR (<10 mg) (N=152)	Placebo (N=161)
<b>Autonomic Nervous System</b>		
Dry Mouth	3	1
<b>Body as a Whole</b>		
Allergy	4	1

Back pain	3	2
Influenza-like symptoms	2	-
Chest pain	1	-
Fatigue	1	2
<b>Cardiovascular System</b>		
Palpitation	2	-
<b>Central and Peripheral Nervous System</b>		
Headache	19	22
Drowsiness	8	5
Dizziness	5	1
Lethargy	3	1
Drugged feeling	3	-
Lightheadedness	2	1
Depression	2	1
Abnormal dreams	1	-
Amnesia	1	-
Anxiety	1	1
Nervousness	1	3
Sleep disorder	1	-
<b>Gastrointestinal System</b>		
Nausea	6	6
Dyspepsia	5	6
Diarrhoea	3	2
Abdominal pain	2	2
Constipation	2	1
Anorexia	1	1
Vomiting	1	1
<b>Immunologic System</b>		
Infection	1	1
<b>Musculoskeletal System</b>		
Myalgia	7	7
Arthralgia	4	4
<b>Respiratory System</b>		
Upper respiratory infection	5	6
Sinusitis	4	2
Pharyngitis	3	1
Rhinitis	1	3
<b>Skin and Appendages</b>		
Rash	2	1
<b>Urogenital System</b>		
Urinary tract infection	2	2

\*Events reported by at least 1% of patients treated with Stilnox.

### **Post Marketing Data**

<b>Infections and infestations</b>	<u>Common:</u> Upper respiratory tract infection, lower respiratory tract infection
<b>Immune system disorders</b>	<u>Rare:</u> Angioneurotic oedema

<b>Metabolism and nutritional disorders</b>	<u>Uncommon:</u> Appetite disorder
<b>Psychiatric disorders</b>	<u>Common:</u> Drowsiness, hallucinations, agitation, nightmare, depression
	<u>Uncommon:</u> Confusion, memory disturbances, reduced alertness, irritability, restlessness, aggressiveness, somnambulism (See section 4.4 – Special warnings and precautions for use-‘Somnambulism and Associated Behaviors’), euphoric mood
	<u>Rare:</u> Perceptual disturbances, aggravated insomnia, libido disorder, delusion, rages, inappropriate behavior, dependence (withdrawal symptoms, or rebound effects may occur after treatment discontinuation), other adverse behavioral effects.
	<u>Not known:</u> Anger, abnormal behavior, and complex sleep behaviors, delirium have been reported ( <i>See section 4.4 Special warnings and precautions for use</i> )
<b>Nervous system disorders</b>	<u>Common:</u> Dizziness, somnolence, headache, exacerbated insomnia, cognitive disorders such as anterograde amnesia (amnesic effects may be associated with inappropriate behavior)
	<u>Uncommon:</u> Paraesthesia, disturbance in attention, tremor, speech disorder
	<u>Rare:</u> Ataxia, dysarthria, depressed level of consciousness
<b>Eye disorders</b>	<u>Uncommon:</u> Diplopia, vision blurred, visual impairment
<b>Respiratory, thoracic and mediastinal disorders</b>	<u>Very rare:</u> Respiratory depression ( <i>See section 4.4 Special warnings and precautions for use – Respiratory function</i> ) has been reported
<b>Gastrointestinal disorders</b>	<u>Common:</u> Diarrhoea, nausea, vomiting, abdominal pain
<b>Hepatobiliary disorders</b>	<u>Uncommon:</u> Elevated liver enzymes
	<u>Rare:</u> Hepatocellular, cholestatic and mixed liver injury ( <i>See section 4.2 Dose and method of administration, section 4.3 Contraindications, and section 4.4 Special warnings and precautions for use</i> )
<b>Skin and subcutaneous tissue</b>	<u>Uncommon:</u> Rash, pruritus, urticaria, hyperhidrosis
<b>Musculoskeletal and connective tissue disorders</b>	<u>Common:</u> Back pain
	<u>Uncommon:</u> Arthralgia, myalgia, muscle spasms, neck pain, muscular weakness

<b>General disorders and administration site conditions</b>	<u>Common:</u> Fatigue
	<u>Rare:</u> Fall, ataxia/gait disturbances, drug tolerance

### Reporting suspected adverse effects

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product.

Any suspected adverse events should be reported to the Ministry of Health according to the National Regulation by using an online form at <https://sideeffects.health.gov.il/>

### 4.9 OVERDOSE

**Signs and Symptoms:** In reports of overdose with Stilnox alone or with other CNS depressant agents (including alcohol), impairment of consciousness has ranged from somnolence to coma and more severe symptomology, including fatal outcomes have been reported. Fatalities have occurred when overdoses of multi CNS depressants were taken.

**Recommended Treatment:** General symptomatic and supportive measures should be used, along with immediate gastric lavage where appropriate. Intravenous fluids should be administered as needed. Sedative drugs should be withheld, even if excitation occurs.

Zolpidem has been shown in trials to be non-dialysable.

Use of flumazenil may be considered when serious symptoms are observed. However, flumazenil administration may contribute to the appearance of neurological symptoms, such as convulsions, since zolpidem does not exhibit the anticonvulsant effects of benzodiazepines.

## 5. PHARMACOLOGICAL PROPERTIES

### 5.1 Mechanism of action

Zolpidem belongs to the imidazopyridine group of compounds and is structurally unrelated to other hypnotic agents. Zolpidem selectively binds the omega-1 receptor subtype (also known as the benzodiazepine-1 subtype) which is the alpha unit of the GABA-A receptor complex. Whereas benzodiazepines non-selectively bind all three omega receptor subtypes, zolpidem preferentially binds the omega-1 subtype. The modulation of the chloride anion channel via this receptor leads to the specific sedative effects demonstrated by zolpidem i.e. the preservation of deep sleep (stage 3 and 4 slow wave sleep).

These effects are reversed by the benzodiazepine antagonist flumazenil.

**In animals:** The selective binding of zolpidem to omega-1 receptors may explain the virtual absence at hypnotic doses of myorelaxant and anti-convulsant effects in animals which are normally exhibited by benzodiazepines which are not selective for omega-1 sites.

**In humans:** The preservation of deep sleep (stages 3 and 4 - slow-wave sleep) may be explained by the selective omega-1 binding by zolpidem. All identified effects of zolpidem are reversed by the benzodiazepine antagonist flumazenil.

### Clinical trials

### ***Insomnia in non-elderly adults***

Short term (1 to 2 nights) placebo controlled studies in 620 volunteers showed that Stilnox 2.5 to 10 mg decreased the latency of persistent sleep in a dose dependent manner. No further increase in efficacy was seen in doses up to STILNOX 40 mg.

The efficacy of Stilnox 2.5 - 20 mg was investigated in 11 placebo-controlled studies in 1606 (513 received Stilnox 10 mg) non-elderly insomniacs over a period of 2-35 nights. Stilnox 10 mg was superior to placebo using both objective (polysomnography) and subjective methods of assessment. Stilnox 20 mg showed little increase in efficacy.

### ***Insomnia in the elderly***

Four studies in 145 elderly (>65 years) patients showed, using objective (2 studies) and subjective (4 studies) methods of assessment that STILNOX 5 mg was the dose giving the optimum efficacy/safety ratio.

*Next day residual effects:* There was no evidence of residual next-day effects seen with Stilnox in several studies utilising the Multiple Sleep Latency Test (MSLT), the Digit Symbol Substitution Test (DSST), and patient ratings of alertness. In one study involving elderly patients, there was a small but statistically significant decrease in one measure of performance, the DSST, but no impairment was seen in the MSLT in this study.

*Rebound Effects:* Although there were no studies to exclude this effect, there was no objective (polysomnographic) evidence of rebound insomnia at recommended doses seen in studies evaluating sleep on the nights following discontinuation of Stilnox. There was subjective evidence of impaired sleep in the elderly on the first post-treatment night at doses above the recommended elderly dose of 5 mg.

*Memory Impairment:* Two small studies (n=6 and n=9) using objective measures of memory yielded little evidence for memory impairment following the administration of Stilnox. There was subjective evidence from adverse event data for anterograde amnesia occurring in association with the administration of Stilnox, predominantly at doses above 10 mg.

*Effects on Sleep Stages:* In studies that measured the percentage of sleep time spent in each sleep stage, Stilnox has generally been shown to preserve sleep stages. Sleep time spent in stages 3 and 4 (deep sleep) was found comparable to placebo with only inconsistent, minor changes in REM (paradoxical) sleep at the recommended dose.

The hypnotic efficacy and safety of Stilnox has not been assessed in children and pregnant women.

## **5.2 PHARMACOKINETIC PROPERTIES**

### **Absorption**

Stilnox has both a rapid absorption and onset of hypnotic action. Peak plasma concentration is reached at between 0.5 and 3 hours. Following oral administration, bioavailability is 70% due to a moderate first-pass metabolism.

The elimination half-life is short, with a mean value of 2.4 hours ( $\pm 0.2$  h) and a duration of action of up to 6 hours. Stilnox pharmacokinetic profile is linear in the therapeutic dose range, and is not modified upon repeated administration.

### **Distribution**

Protein binding amounts to approximately 90%. The distribution volume in adults is  $0.54 \pm 0.02$  L/kg and decreases to  $0.34 \pm 0.05$  L/kg in the very elderly.

## Metabolism

The main cytochrome P450 enzyme involved in the hepatic biotransformation of zolpidem is CYP3A4. CYP1A2 and CYP2D6 contribute minimally to the metabolism of zolpidem (*See section 4.5 Interactions with other medicines and other forms of interactions*).

## Excretion

All metabolites are pharmacologically inactive and are eliminated in the urine (56%) and in the faeces (37%). Furthermore, they do not interfere with zolpidem plasma binding.

## Effect of Food

A food-effect study in 30 healthy male volunteers compared the pharmacokinetics of zolpidem 10 mg when administered while fasting or 20 minutes after a meal. Results demonstrated that with food, mean AUC and  $C_{max}$  were decreased by 15% and 25%, respectively, while mean  $T_{max}$  was prolonged by 60% (from 1.4 to 2.2 hr). The half-life remained unchanged. These results suggest that, for faster sleep onset, Stilnox should not be administered with or immediately after a meal.

## Special Populations

Zolpidem did not accumulate in young adults following nightly dosing with 20 mg zolpidem tartrate tablets for 2 weeks. In the elderly, the recommended dose for Stilnox is 5 mg (*See section 4.2 Dose and method of administration and Section 4.4 Special warnings and precautions for use*). This recommendation is based on several studies in which the mean  $C_{max}$ ,  $T_{1/2}$  and AUC were significantly increased when compared to results in young adults. Zolpidem did not accumulate in elderly subjects following nightly oral dosing of 10 mg for 1 week.

The pharmacokinetics of Stilnox in eight patients with chronic hepatic insufficiency were compared to results in healthy subjects. Following a single 20 mg oral zolpidem dose, mean  $C_{max}$  and AUC were found to be two times (250 vs 499 ng/mL) and five times (788 vs 4,203 ng-hr/mL) higher, respectively, in hepatically compromised patients.  $T_{max}$  did not change.

The mean half-life in cirrhotic patients of 9.9 hr (range: 4.1 to 25.8 hr) was greater than that observed in normal of 2.2 hr (range: 1.6 to 2.4 hr). Dosing should be modified accordingly in patients with hepatic insufficiency (*See section 4.2 Dose and method of administration and Section 4.4 Special warnings and precautions for use*).

In patients with renal insufficiency, whether dialysed or not, there is a moderate reduction in clearance. The other pharmacokinetic parameters are unaffected. Zolpidem has been shown to be non-dialysable.

## 5.3 PRECLINICAL SAFETY DATA

### Genotoxicity

Zolpidem was not genotoxic in assays for gene mutations (*Salmonella typhimurium* histidine reversion assay, L5178Y mouse lymphoma assay), for chromosomal aberrations (human lymphocytes, mouse micronucleus assay) and for DNA repair assays (in human fibroblasts and rat hepatocytes). The mutagenic activity of zolpidem and/or its metabolites was

equivocal in a Chinese hamster V79/HRPT gene mutation assay in the presence of metabolic activation.

### **Carcinogenicity**

Two year dietary carcinogenicity studies on zolpidem were conducted in rats and mice. No evidence of carcinogenic potential was observed in mice at plasma concentrations (AUC) of zolpidem and its major human metabolite of about 2 and 7-12 times, respectively, the anticipated clinical exposure at the maximum recommended clinical dose. An increased incidence of renal liposarcomas was observed in male rats (6% cf. 0 in controls) at plasma concentrations (AUC) of zolpidem and its major metabolite of at least 22 and 9 times, respectively, the anticipated human exposure.

## **6 PHARMACEUTICAL PARTICULARS**

### **6.1 LIST OF EXCIPIENTS**

Lactose monohydrate, microcrystalline cellulose, hypromellose, sodium starch glycollate and magnesium stearate. The coating on the tablets consists of hypromellose, titanium dioxide and macrogol 400.

### **6.2 INCOMPATIBILITIES**

Incompatibilities were either not assessed or not identified as part of the registration of this medicine.

### **6.3 SHELF LIFE**

The expiry date of the product is indicated on the packaging materials.

### **6.4 SPECIAL PRECAUTIONS FOR STORAGE**

Store below 25° C.

### **6.5 NATURE AND CONTENTS OF CONTAINER**

Stilnox 10 mg comes in PVC/Al blister packs of 7 or 14 or 20 tablets


### **6.6 SPECIAL PRECAUTIONS FOR DISPOSAL**

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

## 6.7 PHYSICOCHEMICAL PROPERTIES

Zolpidem tartrate is a white to off white colorless, crystalline powder, sparingly soluble in water.

Chemical structure



Its chemical name is 2-(4-methylphenyl)-N,N,6-trimethylimidazo [1,2,a] pyridine-3-acetamide hemitartrate.

Its molecular formula is (C<sub>19</sub>H<sub>21</sub>N<sub>3</sub>O)<sub>2</sub>, C<sub>4</sub>H<sub>6</sub>O<sub>6</sub>. MW is 764.9.

## 7. IMPORTER AND MARKETING AUTHORISATION HOLDER AND ITS ADDRESS

Sanofi Israel ltd. Greenwork Park, P.O Box 47, Yakum

## 8 MARKETING AUTHORISATION NUMBERS

104-51-27587-00

*Revised in September 2025 according to MoH guidelines.*