

**Patient leaflet in accordance with the Pharmacists' Regulations (Preparations) - 1986**

This medicine is dispensed with a doctor's prescription only

## Pentasa Suppositories

### 1 gram

#### Composition:

Each suppository contains: 1 gram of mesalazine

**Inactive ingredients:** See section 6, 'Additional information'.

**Read the entire leaflet carefully before you start using this medicine.** This leaflet contains concise information about this medicine. If you have any further questions, consult with your doctor or pharmacist.

This medicine has been prescribed to treat your illness. Do not pass it on to others. It may harm them, even if it seems to you that their illness is similar to yours.

#### 1. What is this medicine intended for?

For treatment of ulcerative proctitis.

**Therapeutic group:** Anti-inflammatory medicine of the salicylates class.

#### 2. Before using this medicine

##### Do not use this medicine if you:

- are sensitive (allergic) to mesalazine or to any of the other ingredients of this medicine (see section 6).
- are sensitive (allergic) to other salicylates (such as aspirin).
- have severe liver or kidney problems.

##### Special warnings about using this medicine

##### Before treatment with this medicine, tell your doctor if you:

- are allergic to sulphasalazine (risk of allergy to salicylates)
- currently or previously had liver or kidney disease.
- have a medical condition that can make you prone to bleeding.
- are taking medicines that may affect your kidney function, e.g. nonsteroidal anti-inflammatory drugs (NSAIDs) such as aspirin.
- have a lung problem, particularly asthma.
- have ever developed a severe skin rash or skin peeling, blistering and/or mouth sores after taking mesalazine.

Mesalazine may produce red-brown urine discolouration after a reaction with sodium hypochlorite (household bleach) in toilet water. It occurs due to a chemical reaction and is harmless to your health.

##### Pay special attention to the following conditions:

- Serious skin reactions including drug reaction with eosinophilia and systemic symptoms (DRESS), Stevens-Johnson syndrome, toxic epidermal necrolysis (TEN) have been reported in association with mesalazine treatment. Stop taking the medicine and contact a doctor immediately if you have any of the symptoms of serious skin reactions such as those described in section 4.
- If you experience strong or recurrent headaches, disturbed vision, or ringing or buzzing in the ears – contact your doctor immediately.

##### During treatment with this medicine

- If you suddenly start experiencing abdominal cramps, abdominal pain, fever, severe headache and rash, **stop taking this medicine** and contact a doctor immediately.
- Kidney stones may develop during the treatment with mesalazine. Symptoms may include pain at the sides of your abdomen and blood in urine. Ensure drinking a sufficient amount of liquids during the treatment with mesalazine.
- Avoid dehydration while taking this medicine. Dehydration may occur after prolonged vomiting and/or diarrhoea, high fever or excessive sweating. If this occurs, consult with a doctor or pharmacist as soon as possible.

##### Tests and follow up

During the treatment with the medicine, your doctor may refer you for blood and urine tests to verify that your kidney function is normal, especially at the beginning of treatment.

##### Drug interactions

**If you are taking or have recently taken other medicines, including nonprescription medications and dietary supplements, tell your doctor or pharmacist,** particularly if you are taking:

- azathioprine (given after transplantations or to treat autoimmune diseases).
- 6-mercaptopurine or thioguanine (chemotherapy, used to treat leukaemia).
- certain medicines that inhibit blood clotting (medicines against thrombosis or for blood thinning, such as warfarin).

##### Pregnancy and breastfeeding

If you are pregnant, think you may be pregnant, are planning to become pregnant, or if you are breastfeeding, consult with your doctor or pharmacist before using this medicine. There is limited experience with the use of mesalazine during pregnancy and breastfeeding. Blood disorders have been reported in newborns of mothers who have taken this medicine. The newborn may develop allergic reactions after breastfeeding, such as diarrhoea. If your newborn develops diarrhoea, stop breastfeeding.

##### Driving and using machines

The medicine has no known effect on the ability to drive and/or operate machines.

#### 3. How to use this medicine?

Always use this medicine according to your doctor's instructions. Check with your doctor or pharmacist if you are not sure about your dosage and how you should take this medicine. The dosage and how you should take this medicine will be determined by your doctor only. For use in adults only, not recommended for use in children. The recommended dosage for adults is usually no more than 1 suppository per day before bedtime.

The medicine is intended for rectal treatment (via the anus). To prolong the period during which the medicine remains in the body, it is recommended to insert it before bedtime.

##### Do not exceed the recommended dose.

##### How to use Pentasa Suppositories:

You should make sure that you empty your bowels prior to suppository insertion.

1. Separate a single suppository by tearing the blister along the perforation.
2. Put a finger protector on the finger with which you will insert the suppository.
3. Remove the suppository from the blister before insertion by pushing the suppository through the blister.
4. To make the insertion easier, you can moisten the suppository with some water.
5. Insert the suppository gently and fully into the anus. It may be easier to do it while lying down on one side and raising the other knee.
6. Remain lying for a few minutes after insertion to help keep the suppository in place. If the suppository comes out within the first 10 minutes, a new suppository should be inserted.
7. Dispose of the empty blister and used finger protector and wash your hands.

**If you have accidentally taken a higher dose.** If you have accidentally used more suppositories than you should, or if a child has accidentally swallowed some of the medicine, immediately see a doctor or go to a hospital emergency room and bring the medicine package with you.

**If you forget to take the medicine.** If you forgot to use this medicine at the scheduled time, use it as soon as you remember, but maintain an interval of at least 6 hours until your next dose. Never take a double dose!

Adhere to the treatment as recommended by your doctor.

Even if your health improves, do not stop the treatment with this medicine without consulting with your doctor or pharmacist.

**Do not take medicines in the dark! Check the label and dose every time you take a medicine. Wear glasses if you need them.**

**If you have any further questions about using this medicine, consult with your doctor or pharmacist.**

#### 4. Side effects

Like with all medicines, using Pentasa may cause side effects in some users. Do not be alarmed by this list of side effects; you may not experience any of them.

Following rectal administration, you may experience local side effects such as itching, rectal discomfort and urge for bowel emptying.

##### Stop using this medicine and contact your doctor immediately in the event of:

- reddish non-elevated, target-like or circular patches on the trunk, often with central blisters, skin peeling, ulcers of mouth, throat, nose, genitals and eyes, widespread rash, fever and enlarged lymph nodes. These serious effects can be preceded by fever and flu-like symptoms.
  - itching, skin rash, swelling of the face, lips or throat, difficulties breathing or wheeziness (signs of an allergic reaction).
  - unexplained bleeding, bruising, skin rash, fever or sore throat (signs of a blood disorder).
  - a change in the amount or colour of urine (signs of kidney problems).
- If you experience any of the above side effects, you should contact your doctor or go to the nearest hospital emergency department immediately.

##### Additional side effects

##### Common side effects – side effects affecting 1-10 in 100 users:

- diarrhoea
- abdominal pain
- nausea
- vomiting
- headache
- rash
- flatulence (passing wind)
- rectal discomfort
- irritation at the site of application

##### Rare side effects – side effects affecting 1-10 in 10,000 users:

- inflammation of some areas of the heart (myocarditis and pericarditis) which may cause shortness of breath and chest pain or palpitations (rapid or irregular heartbeats).
- inflammation of the pancreas (symptoms include back and/or abdominal pain).
- increased amylase (enzyme that helps digest carbohydrates).
- dizziness.
- increased sensitivity of your skin to sun and ultraviolet light (photosensitivity).

##### Very rare side effects – side effects affecting less than 1 in 10,000 users:

- anaemia and other blood disorders (decrease in the numbers of certain blood cells, which may cause unexplained bleeding, bruising, fever or sore throat)
- liver disorders (symptoms include jaundice (yellowing of the skin and/or eyes) and/or pale stool)
- kidney disorders (symptoms include blood in the urine and/or oedema (swelling due to build-up of fluid))
- peripheral neuropathy (a condition affecting the nerves of the hands and feet. Symptoms include tingling and numbness)
- allergic and fibrotic lung reactions, inflammation of the lining of the lungs or lung scarring (symptoms include coughing, bronchospasm, chest discomfort or pain on breathing, breathing difficulties, bloody and/or excessive phlegm)
- pancolitis (a kind of inflammatory bowel disorder (IBD) that affects the entire internal lining of the large bowel)
- hair loss (reversible)
- muscle or joint pain
- inflammation which can affect different parts of the body, such as joints, skin, kidneys, heart etc. (symptoms include painful joints, fatigue, fever, abnormal and unexplained bleeding (e.g. nose bleeds), bruising, purple patches on the skin (including severe skin erosions and severe blistering that may affect the skin as the protective barrier of the body))
- reduced sperm count (reversible)
- severe diarrhoea and abdominal pain because of an allergic reaction of the bowel to this medicine
- allergic reactions and fever may also occasionally occur.

##### Side effects of unknown frequency (cannot be estimated from the available information):

- kidney stones and associated kidney pain (see also section 2)
- changes in urine colour
- Inform your doctor immediately if you experience strong or recurrent headaches, disturbed vision, or ringing or buzzing in the ears. These could be symptoms of increased pressure within your skull (idiopathic intracranial hypertension).

**If you experience any side effect, if any side effect gets worse, or if you experience a side effect not mentioned in this leaflet, consult with your doctor.**

##### Reporting side effects

You can report side effects to the Ministry of Health by following the link 'Reporting Side Effects of Drug Treatment' on the Ministry of Health home page ([www.health.gov.il](http://www.health.gov.il)) which links to an online form for reporting side effects. You can also use this link: <https://sideeffects.health.gov.il>

#### 5. How to store the medicine?

- Prevent poisoning! To prevent poisoning, keep this, and all other medicines, in a closed place, out of the reach and sight of children and/or infants. Do not induce vomiting unless explicitly instructed to do so by a doctor.
- Do not use the medicine after the expiry date (exp. date) which is stated on the package. The expiry date refers to the last day of that month.

##### Storage conditions

- Store below 25°C.

#### 6. Additional information

In addition to the active ingredient, the suppositories also contain: povidone, macrogol 6000, magnesium stearate, talc

##### What the medicine looks like and contents of the pack:

Oblong, white to tan speckled suppository. Each pack contains 28 suppositories packed in blisters of 7 suppositories.

##### Registration holder's name and address:

Ferring Pharmaceuticals Ltd., 8 Hashita St., Industrial Park, Caesarea 3088900.

**Manufacturer's name and address:** Ferring International Center SA, Switzerland.

This leaflet was revised in July 2025.

**Registration number of the medicine in the National Drug Registry of the Ministry of Health: 062 73 26904.**