

PATIENT PACKAGE INSERT IN ACCORDANCE WITH THE PHARMACISTS' REGULATIONS (PREPARATIONS) – 1986

The medicine is dispensed with a doctor's prescription only

HEXAKAPRON® TABLETS

Composition

Each tablet contains: Tranexamic Acid 500 mg

For information on inactive ingredients in the preparation – see section 6 – “Further Information”.

Read the leaflet carefully in its entirety before using the medicine. This leaflet contains concise information about the medicine. If you have further questions, refer to the doctor or pharmacist.

This medicine has been prescribed to treat you. Do not pass it on to others. It may harm them even if it seems to you that their medical condition is similar.

1. WHAT IS THE MEDICINE INTENDED FOR?

The medicine is intended to treat hemorrhage.

Therapeutic group

Prophetic blood clotting – antifibrinolytic.

2. BEFORE USING THE MEDICINE

Do not use the medicine if:

- You are sensitive (allergic) to the active ingredient (tranexamic acid) or to any of the additional ingredients contained in the medicine (see section 6).
- You have serious kidney problems (kidney failure).
- You have an active thromboembolic disease (hypercoagulability of the blood).
- You have a history of venous or arterial thrombosis.
- You have a history of convulsions.
- You are at risk of excessive bleeding as a result of a coagulation disorder called consumption coagulopathy.

Special warnings regarding use of the medicine

Before treatment with Hexakapron, tell the doctor if:

- You have blood in the urine, due to an increased risk of blockage of the urinary tract.
- You have ever had uncontrollable bleeding.
- You have disseminated intravascular coagulation (DIC), a disease where the blood starts to clot throughout the body.
- You are taking, every day and for a long period, a medicine to treat a hereditary disease called angioneurotic edema (HANO). If so, you may need to have routine eye tests and blood tests performed to check liver function.
- You have irregular menstrual periods.
- You have had a thromboembolic event in the past.
- One of your family members has suffered from thromboembolic disease (thrombophilia, hypercoagulability).
- You have kidney disease.

Drug interactions

If you are taking, or have recently taken, other medicines, including non-prescription medicines and nutritional supplements, tell the doctor or pharmacist. Especially if you are taking:

- Fibrinolytic medicines (that help to dissolve blood clots), such as streptokinase, since tranexamic acid (the active ingredient in Hexakapron) will prevent the activity of these medicines.
- Oral contraceptives – may increase the risk of formation of blood clots.

Pregnancy, breastfeeding and fertility

If you are pregnant or breastfeeding, think you are pregnant or are planning a pregnancy, consult your doctor or pharmacist before taking this medicine.

Driving and using machines

The medicine has little or no effect on the ability to drive. However, visual disturbance may occur after taking tranexamic acid (the active ingredient in the medicine).

Important information about some of the ingredients of the medicine

This medicine contains less than 23 mg sodium per tablet and is therefore considered sodium-free.

3. HOW SHOULD YOU USE THE MEDICINE?

Always use the preparation according to the doctor's instructions. Check with the doctor or pharmacist if you are uncertain regarding the dosage and treatment regimen of the preparation.

The usual dosage for adults is 2-3 tablets taken 2-3 times per day. With that, the dosage and treatment regimen will be determined by the doctor only, in accordance with the reason for administration.

Children and/or adolescents: The dosage will be determined by the doctor based on body weight.

People suffering from kidney problems: The dosage will be determined by the doctor; the dosage may have to be reduced.

Do not exceed the recommended dose.

Crushing/halving/chewing

The tablet can be halved on the score line.

There is no information regarding crushing or chewing.

Always take the medicine with a glass of water.

If you accidentally took a higher dosage

If you accidentally took an overdose, or if a child has accidentally swallowed the medicine, refer immediately to a doctor or proceed to a hospital

emergency room and bring the package of the medicine with you.

Taking an overdose of tranexamic acid may cause diarrhea, nausea, vomiting, dizziness or lightheadedness when standing up.

If you forgot to take the medicine

If you forgot to take this medicine at the designated time, do not take a double dose to compensate for the forgotten dose. Take the next dose as planned.

Adhere to the treatment regimen as recommended by the doctor.

Do not take medicines in the dark! Check the label and the dose each time you take medicine. Wear glasses if you need them.

If you have further questions regarding use of the medicine, consult the doctor or pharmacist.

4. SIDE EFFECTS

As with any medicine, use of Hexakapron may cause side effects in some users. Do not be alarmed by reading the list of side effects. You may not suffer from any of them.

Stop taking the medicine and proceed immediately to an emergency room if you experience any of the following side effects: Rare side effects (effects that occur in less than 1 user in 1,000):

A blood clot in the leg or lungs, the symptoms of which may include:

- Pain, swelling or redness in the leg
- Difficulty breathing or shortness of breath
- Chest pain, which may get worse while breathing

Very rare side effects (effects that occur in less than 1 user in 10,000):

An allergic reaction, the symptoms of which may include: swelling of your throat and tongue; difficulty breathing or shortness of breath; difficulty swallowing, tightness in the throat or a hoarse voice; wheezing, coughing or noisy breathing; a rash that is swollen, raised or itchy

Stop taking the medicine and refer immediately to a doctor if you experience any of the following side effects:

Rare side effects (effects that occur in less than 1 user in 1,000):

- Itching, redness or swelling of the skin
- Vision problems, especially in seeing colors
- A blood clot in the eye. This may cause bleeding in the eye or loss of vision

Very rare side effects (effects that occur in less than 1 user in 10,000):

- A blood clot in the blood vessels (a condition called thrombosis)
- Nausea, vomiting, diarrhea – these effects are generally mild and pass very quickly, but if they continue, tell your doctor or pharmacist.

Side effect of unknown frequency (frequency cannot be determined from the available data):

Convulsions, especially in cases of misuse

If a side effect occurs, if one of the side effects worsens or if you suffer from a side effect not mentioned in the leaflet, consult with the doctor.

Reporting side effects

Side effects can be reported to the Ministry of Health by clicking on the link “Reporting Side Effects of Drug Treatment” found on the Ministry of Health homepage (www.health.gov.il) that directs you to the online form for reporting side effects, or by entering the link:

<https://sideeffects.health.gov.il>

5. HOW SHOULD THE MEDICINE BE STORED?

- **Avoid poisoning!** This medicine and any other medicine should be kept in a safe place out of the reach and sight of children and/or infants in order to avoid poisoning. Do not induce vomiting unless explicitly instructed to do so by the doctor.
- Do not use the medicine after the expiry date (Exp. Date) that appears on the package. The expiry date refers to the last day of that month.
- **Store in a dry place, below 25°C, in the original package in order to protect from light.**
- Do not discard medicines in the wastewater or household waste. Ask the pharmacist how to dispose of medicines that are not in use. These measures will help protect the environment.

6. FURTHER INFORMATION

In addition to the active ingredient, the medicine also contains:

Microcrystalline cellulose, sodium starch glycolate, magnesium stearate, methylcellulose, talc, colloidal silicon dioxide.

What the medicine looks like and the contents of the package

A white, round, flat beveled tablet, engraved “TEVA” on one side and bisecting line on the other. Each package contains 20 or 30 tablets packed in trays (blister).

Not all package sizes may be marketed.

Name of Manufacturer and License Holder and its Address

Teva Israel Ltd.,

124 Dvora HaNevi'a St., Tel Aviv 6944020

This leaflet was revised in June 2025.

Registration number of the medicine in the National Drug Registry of the Ministry of Health: 016.35.24864

teva

HEXA TAB PL SH 0625