

**PATIENT PACKAGE INSERT IN ACCORDANCE WITH THE PHARMACISTS' REGULATIONS (PREPARATIONS) - 1986**  
The medicine is dispensed with a doctor's prescription only

Medicine name and its form:

## EVRA® Transdermal patch

The active ingredients and their quantities:

**Norelgestromin 6 mg**

**Ethinyl estradiol 600 mcg**

Inactive and allergenic ingredients in the preparation - see in section 6 "Further information".

**Read this leaflet carefully in its entirety before using the medicine.** This leaflet contains concise information about the medicine. If you have further questions, refer to the doctor or pharmacist. This medicine has been prescribed for you. Do not pass it on to others. It may harm them even if it seems to you that their medical condition is similar.

<b>Important things to know about combined hormonal contraceptives (CHCs):</b> <ul style="list-style-type: none"><li>They are one of the most reliable reversible methods of combined hormonal contraceptives if used correctly.</li> <li>They slightly increase the risk of having a blood clot in the veins and arteries, especially in the first year or when restarting a combined hormonal contraceptive following a break of 4 or more weeks.</li> <li>Please be alert and see your doctor if you think you may have symptoms of a blood clot (see in section 2 "Blood clots").</li></ul>
---

### 1. WHAT IS THE MEDICINE INTENDED FOR?

Female contraceptive. EVRA is intended for women of fertile age. EVRA is a transdermal patch that contains two types of sex hormones, a progestogen, called norelgestromin, and an estrogen, called ethinyl estradiol. Since the medicine contains two hormones, it is called a combined hormonal contraceptive used to prevent pregnancy.

**Therapeutic group:** Combined contraceptive patch.

### 2. BEFORE USING THE MEDICINE

Before using EVRA, read the information regarding blood clots found in section 2. It is important to carefully read about the symptoms indicative of blood clots (see in section 2 "Blood clots").

<b>Do not use the medicine if:</b> <ul style="list-style-type: none"><li>you are sensitive (allergic) to norelgestromin, ethinyl estradiol or to any of the other ingredients of the medicine (listed in section 6 in this leaflet "Further information").</li> <li>you are suffering, or have suffered in the past, from a blood clot in a blood vessel of the legs (Deep Vein Thrombosis or DVT), in the lungs (Pulmonary Embolism or PE), or in other organs.</li> <li>you are suffering from a disorder that affects blood clotting – e.g., protein C deficiency, protein S deficiency, antithrombin III deficiency, Factor V Leiden or antiphospholipid antibodies.</li> <li>you are due to undergo surgery or if you will be immobile (you will not be able to walk) for a long time (see in section 2 "Blood clots").</li> <li>you have suffered in the past from a heart attack or a stroke.</li> <li>you are suffering, or have suffered, in the past, from angina pectoris (a condition that causes severe chest pain and may be the first sign of a heart attack) or from transient ischaemic attack (temporary stroke symptoms).</li> <li>you are suffering from a disease that may increase the risk of a blood clot in the arteries, such as: <ul style="list-style-type: none"><li>Severe diabetes with blood vessel damage.</li> <li>Severe hypertension.</li> <li>Very high levels of fats in the blood (cholesterol or triglycerides).</li> <li>A condition called hyperhomocysteinaemia.</li></ul></li> <li>you are suffering, or have suffered in the past, from a type of migraine called 'Migraine with aura'.</li> <li>you have been told that you may be at risk of breast cancer or cancer of the womb, cervix or vagina.</li> <li>you have suffered from liver tumours or from liver disease that caused impaired liver function.</li> <li>you are suffering from unexplained vaginal bleeding.</li> <li>you are suffering from hepatitis C and are taking medicines containing ombitasvir/paritaprevir/ritonavir, dasabuvir, glecaprevir/pibrentasvir or sofosbuvir/velpatasvir/voxilaprevir (also see in section 2 "Drug Interactions").</li></ul> <p>Do not use EVRA if any of the above-mentioned conditions applies to you. If you are unsure, consult the doctor or pharmacist before using EVRA.</p>
---

### Special warnings regarding use of the medicine

Seek urgent medical attention if you notice that you are suffering from symptoms that could be indicative of a blood clot in the leg (Deep Vein Thrombosis or DVT), a blood clot in the lungs (Pulmonary Embolism), a heart attack or a stroke (see in section 2 "Blood clots" below).

For a description of the symptoms of these serious side effects, please see the "How to recognize formation of blood clots" in section 2 later in the leaflet.

### Warnings and precautions

Before starting to use EVRA, see a doctor for a medical check-up.

**Consult a doctor before using EVRA if you suffer from the following conditions:**

- Tell the doctor if you are suffering from any of the following conditions, or if any of the following conditions develops or worsens during the course of treatment with EVRA.
- Crohn's disease or ulcerative colitis (chronic inflammatory bowel disease).
- systemic lupus erythematosus (SLE), an autoimmune disease that affects the immune system.
- a blood clotting problem that causes kidney failure, called Haemolytic Uraemic Syndrome (HUS).
- serious anaemia (an inherited disease of the red blood cells).
- high levels of fats in the blood (hypertriglyceridaemia) or a family history of this condition. Hypertriglyceridaemia is found to be associated with increased risk of developing pancreatitis.
- if you are due to undergo surgery, or if you will be immobile (you will not be able to walk) for a long time (see in section 2 "Blood clots").
- If you have just given birth, you are at an increased risk of blood clots. Consult the doctor regarding how soon after delivery you can start using EVRA.
- inflammation in the veins under the skin (Superficial thrombophlebitis).
- varicose veins.
- symptoms of angioedema such as swollen face, tongue and/or throat and/or difficulty swallowing or hives potentially with difficulty breathing contact a doctor immediately. Products containing estrogens may cause or worsen the symptoms of hereditary and acquired angioedema.

### Blood clots

Using a combined contraceptive, such as EVRA, increases the risk of formation of blood clots compared with the risk in women who do not use such preparations. In rare cases, a blood clot can block blood vessels and cause serious problems. Blood clots can develop:

- in veins (venous thrombosis, venous thromboembolism)
  - in the arteries (arterial thrombosis or arterial thromboembolism)
- Sometimes, recovery from blood clots is not complete. Rarely, there may be lasting effects and, very rarely, there may be fatal effects.

**It is important to remember that the overall risk of formation of harmful blood clots due to use of EVRA is small.**

### How to recognize formation of blood clots?

Seek urgent medical attention if you notice that you are suffering from any of the following signs or symptoms:

<b>Are you experiencing any of the following signs?</b>	<b>What are you likely to be suffering from?</b>
<ul style="list-style-type: none"><li>swelling of one leg or along a vein in the leg or foot, especially when accompanied by: <ul style="list-style-type: none"><li>pain or tenderness in the leg which may only be felt when standing or walking</li> <li>increased warmth in the swollen leg</li> <li>changes in colour of the skin of the leg, e.g., turning pale, red or blue</li></ul></li></ul>	Deep vein thrombosis
<ul style="list-style-type: none"><li>sudden unexplained breathlessness or rapid breathing</li> <li>sudden cough without an obvious cause, which may bring up blood</li> <li>sharp chest pain which can worsen with deep breathing</li> <li>severe dizziness</li> <li>rapid or irregular heartbeat</li> <li>severe stomach pain</li></ul> <p>If you are unsure, please refer to the doctor, as symptoms such as coughing or shortness of breath may be mistaken for a milder condition of respiratory tract infection (e.g., a common cold).</p>	Pulmonary embolism
<ul style="list-style-type: none"><li>symptoms that most commonly occur in one eye: <ul style="list-style-type: none"><li>immediate loss of vision</li> <li>painless blurring of vision which may progress to loss of vision</li></ul></li></ul>	Retinal vein thrombosis (blood clot in the eye)
<ul style="list-style-type: none"><li>chest pain, discomfort, pressure or heaviness</li> <li>sensation of squeezing or fullness in the chest, arm or below the breastbone</li> <li>feeling of fullness, indigestion or choking feeling</li> <li>upper body discomfort radiating to the back, jaw, throat, arm and stomach</li> <li>sweating, nausea, vomiting or dizziness</li> <li>extreme weakness, anxiety or shortness of breath</li> <li>rapid or irregular heartbeat</li></ul>	Heart attack
<ul style="list-style-type: none"><li>sudden weakness or numbness of the face, leg or hand, especially on one side of the body</li> <li>sudden confusion, trouble speaking or understanding</li> <li>sudden trouble seeing in one or both eyes</li> <li>sudden trouble walking, dizziness, loss of balance or coordination</li> <li>sudden, severe or prolonged headache with no known cause</li> <li>seizure</li> <li>loss of consciousness or fainting with or without seizure</li></ul> <p>Sometimes, the symptoms of stroke are brief, with immediate and full recovery, but you should still seek immediate medical attention for fear of an additional stroke.</p>	Stroke
<ul style="list-style-type: none"><li>swelling, slight blue discolouration of the extremities</li> <li>severe stomach pain (acute abdomen)</li></ul>	Blood clots blocking other blood vessels

### Blood clots in a vein

**What can happen when a blood clot forms in a vein?**

- Using a combined hormonal contraceptive has been associated with an increased risk of formation of blood clots in a vein (venous thrombosis). However, these side effects are rare. Mostly, these side effects occur in the first year of use of this type of preparation.
- If a blood clot forms in a vein in the leg or foot, it may cause a deep vein thrombosis (DVT).
- If a blood clot travels from the leg and reaches the lungs, it may cause a pulmonary embolism.
- Very rarely, a blood clot may form in the vein of other organs, such as the eye (retinal vein thrombosis).

**When is the risk of developing a blood clot in a vein highest?**

The risk of developing a blood clot in a vein is highest during the first year of taking a combined hormonal contraceptive for the first time. The risk may also be higher if you restart taking a combined preparation (the same preparation or a different preparation from the same group) after a break of 4 weeks or more.

After the first year, the risk gets smaller, but is always slightly higher than in the population who is not taking a combined hormonal contraceptive.

When you stop using EVRA, the risk of developing blood clots returns to normal within a few weeks.

### What is the risk of developing a blood clot?

The risk depends on your natural risk of venous thromboembolism (VTE) and on the type of combined hormonal contraceptive you are taking.

The overall risk of developing a blood clot in the leg or lung (deep vein thrombosis or pulmonary embolism) with EVRA is low.

- About 2 of 10,000 women who are not taking a combined hormonal contraceptive and are not pregnant will develop a blood clot in a year.
- About 5-7 out of 10,000 women who are taking a combined hormonal contraceptive that contains levonorgestrel, norethisterone or norgestimate will develop a blood clot in a year.
- About 12 out of 10,000 women who are taking a combined hormonal contraceptive that contains etonogestrel or norelgestromin, such as EVRA, will develop a blood clot in a year.

- The risk of developing a blood clot varies according to your medical history (see "Factors that increase the risk of developing a blood clot" further on in this section).

	The risk of developing a blood clot in a year
Women <b>who are not using</b> a combined hormonal contraceptive (pill, patch, ring) and are not pregnant	About 2 out of 10,000 women
Women using a combined hormonal contraceptive pill containing <b>levonorgestrel, norethisterone or norgestimate</b>	About 5-7 out of 10,000 women
Women using EVRA	About 6-12 out of 10,000 women

**Factors that increase the risk of developing a blood clot in a vein**

The risk of developing a blood clot in women using EVRA is low, but some conditions increase this risk. Your risk is higher:

- if you are very overweight (body mass index [BMI] over 30 kg/m²).
- if a member of your immediate family has had a blood clot in the leg, lung or other organ at a young age (below the age of 50). In this case you may have a hereditary blood clotting disorder.
- if you need to undergo surgery, or if you will be immobile (unable to walk) for a long time because of an injury, illness, or if your leg is in a cast. The use of EVRA may need to be stopped several weeks before the surgery or during the period when you are less mobile. If you need to stop treatment with EVRA, please consult the doctor or pharmacist before restarting treatment with EVRA again.
- as you get older (particularly above about the age of 35).
- if you have given birth within the past few weeks.

The risk of developing a blood clot increases as the number of the above-mentioned factors you are suffering from increases.

Travel for over 4 hours may temporarily increase the risk of developing a blood clot, particularly if you are suffering from some of the above-mentioned factors.

Please inform the doctor if you are suffering from any of the above-mentioned factors, even if you are unsure. The doctor may decide that you must stop treatment with EVRA.

If any of the above-mentioned factors changes during the course of treatment with EVRA, for example, a member of your immediate family experiences a thrombosis for an unknown reason, or if you gain a lot of weight, tell the doctor.

### Blood clots in an artery

**What can happen if a blood clot forms in an artery?**

Like a blood clot in a vein, a blood clot in an artery can cause serious problems. For example, a heart attack or a stroke.

**Factors that increase the risk of developing a blood clot in an artery**

It is important to note that the risk of a heart attack or stroke due to use of EVRA is very low but can increase:

- if you are overweight (above the age of 35).

- if you smoke.** When using a combined hormonal contraceptive like EVRA, it is recommended to stop smoking. If you are unable to stop smoking and you are above the age of 35, the doctor may advise you to use a different contraceptive.
- if you are overweight.
- if you suffer from hypertension.
- if a member of your immediate family had a heart attack or a stroke at a young age (below the age of 50). In this case, you could also have a higher risk of heart attack or stroke.

- if you, or someone in your immediate family, have a high level of fat in the blood (cholesterol or triglycerides).
- if you suffer from migraines, especially migraines with aura.
- if you suffer from a heart rhythm problem (a valve problem, a heart rhythm disturbance called atrial fibrillation).
- if you suffer from diabetes.
- if you suffer from more than one of the above-mentioned factors, or if any of them are particularly severe, the risk of developing a blood clot is higher.
- If any of the above-mentioned factors changes during the course of treatment with EVRA, for example, if you gain a lot of weight, tell the doctor.

**Psychiatric disorders**

Some women using hormonal contraceptives including EVRA have reported depression or depressed mood. Depression can be serious and may sometimes lead to suicidal thoughts. If you experience mood changes and depressive symptoms, contact your doctor for further medical advice as soon as possible.

In addition, refer to the doctor, pharmacist or nurse before using EVRA if you suffer from any of the following conditions or if they worsen during the course of treatment with EVRA.

- you might be pregnant.
- headaches that got worse or happen more often.
- you weigh 90 kg or more.
- you suffer from hypertension or your blood pressure gets higher.
- you suffer from gallbladder disease including gallstones or inflammation of the gallbladder.
- you suffer from porphyria (a blood disease).
- you suffer from Sydenham's chorea, a problem of the nervous system involving sudden movements of the body.
- you have suffered from herpes gestationis, a skin rash with blisters during pregnancy.
- you suffer from a decrease in hearing.
- you suffer from diabetes.
- you suffer from depression.
- you suffer from epilepsy or any other condition that causes convulsions.
- you suffer from liver problems, including yellowing of the skin and eyes (jaundice).
- you suffer or have suffered from "pregnancy spots". These are yellowish-brown patches, primarily on the face (called chloasma), that do not go away completely, even after you stop using EVRA. Protect the skin from sunlight or ultraviolet radiation to help prevent you from getting these spots or prevent existing spots from getting worse.
- you suffer from kidney problems.

If you are not sure if any of the above-mentioned conditions applies to you, consult the doctor or pharmacist before using EVRA.

### Sexually transmitted diseases

EVRA does not protect against infection with HIV (AIDS) or other sexually transmitted diseases. These diseases include: chlamydia, genital herpes, genital warts, gonorrhoea, hepatitis B (jaundice), syphilis. Always use a condom to protect yourself from these diseases.

### Medical tests

If you undergo any blood or urine tests, tell the doctor/laboratory staff that you are using EVRA, because the results of some laboratory tests can be affected by it.

### Children and adolescents

EVRA has not been tested in young and adolescent girls under 18 years of age. EVRA must not be used in young and adolescent girls who have not yet had their first menstrual period.

### Drug Interactions

**If you are taking, or have recently taken, other medicines, including non-prescription medicines and nutritional supplements, tell the doctor or pharmacist.**

Do not use EVRA if you have hepatitis C and are taking the medicines containing ombitasvir/paritaprevir/ritonavir, dasabuvir, glecaprevir/pibrentasvir or sofosbuvir/velpatasvir/voxilaprevir as these products may increase liver function blood test results (increase in ALT liver enzyme). Your doctor will prescribe another type of contraceptive prior to starting treatment with these medicines. You can take EVRA again approximately 2 weeks after completion of this treatment. See in section 2 "Do not use the medicine if".

Certain medicines and herbal therapies may impair the effectiveness of EVRA. When EVRA does not work, you may become pregnant or experience unexpected bleeding. These include medicines used for the treatment of:

- certain medicines for the treatment of HIV infection (AIDS) and hepatitis C virus infections (called protease inhibitors and non-nucleoside analog reverse transcriptase inhibitors e.g., ritonavir, nevirapine and efavirenz)
- medicines for the treatment of infections (e.g., rifampicin and griseofulvin)
- anti-seizure medicines (e.g., barbiturates, topiramate, phenytoin, carbamazepine, primidone, oxcarbazepine, and felbamate)

- a medicine to treat pulmonary hypertension (bosentan)

- St. John's wort (an herbal therapy).

If you are taking any of the above-mentioned medicines, you may need a different contraceptive method (e.g., condom, diaphragm or foam). The interfering effect of some of the above-mentioned medicines may last for up to 28 days after discontinuing taking them. Consult the doctor or pharmacist about using a different contraceptive if you use EVRA and are concomitantly taking any of the above-mentioned medicines.

EVRA may affect the effectiveness of the following medicines:

- medicines containing ciclosporin
  - lamotrigine for the treatment of epilepsy (may increase the risk of convulsions).
- The doctor may need to adjust the dosage of the additional medicine. Consult the doctor or pharmacist before taking any medicine.

### Pregnancy and breastfeeding

- Do not use the preparation if you are pregnant or may be pregnant.

- Stop using EVRA immediately if you become pregnant.
- Do not use the preparation if you are breastfeeding or are planning to breastfeed. If you may be pregnant or are planning to become pregnant, consult the doctor or pharmacist before using the medicine.

### Driving and use of machines

You can drive and operate machines when using EVRA.

### Risks in using combined hormonal contraceptives

The information provided below is based on data about combined birth control pills. Since EVRA is a transdermal patch containing hormones similar to those used in combined birth control pills, it is likely to have the same risks. All combined birth control pills have risks, that may lead to disability or death. It has not been shown that a transdermal patch like EVRA is safer than combined birth control pills taken by mouth.

### Cancer and combined hormonal contraceptives

#### Cervical cancer

The prevalence of cervical cancer has been found to be higher in women who used combined hormonal contraceptives. However, this increased prevalence may be associated with other causes including sexually transmitted disease.

#### Breast cancer

The prevalence of breast cancer has been found to be higher in women who used combined hormonal contraceptives. However, it is possible that the combined hormonal contraceptives are not the cause of more women having breast cancer; rather, it may be that women using combined hormonal contraceptives are examined more often, and there is therefore a greater chance of diagnosing breast cancer. The increased risk gradually declines after discontinuing use of combined hormonal contraceptives. After 10 years, the risk is the same as for women who have never used combined hormonal contraceptives.

#### Liver cancer

In rare cases, liver tumours which are not cancerous have been found in women who used combined hormonal contraceptives. More rarely, cancerous liver tumours have been found. Such a tumour can cause internal bleeding with strong pain in the stomach area.

**If this happens, report to your doctor immediately.**

### 3. HOW SHOULD EVRA BE USED?

Always use EVRA according to the doctor's or pharmacist's instructions.

Check with the doctor or pharmacist if you are uncertain about the medicine dosage and treatment regimen. The dosage and treatment regimen will be determined by the doctor only.

**Do not exceed the recommended dose.** Do not swallow!

This medicine is intended for external use only.

- If you do not use it as per the instructions, you increase the risk of getting pregnant.
- Ask the doctor or pharmacist if you are uncertain.
- Always keep non-hormonal contraceptives (e.g., condoms, foam or contraceptive sponge) as a back-up in case of any mistake when using the patch.

### How many patches to use:

Weeks 1, 2, 3 - Put on one patch and leave it on for exactly seven days.

Week 4 - **Do not put on a patch this week.**

**If you have not used a hormonal contraceptive during your previous cycle:**

- Start using EVRA on the first day of your next period.
- If one or more days have elapsed since the start of your period, please refer to the doctor about temporarily using a non-hormonal contraceptive.
- If you switch from an oral contraceptive pill to EVRA:**
  - Wait until you get your menstrual period.
  - Put on your first patch during the first 24 hours of your period.

If the patch is applied after Day 1 of your period:

- Use a non-hormonal contraceptive until Day 8, on which you change the patch.

If you do not get your period within 5 days of taking the last contraceptive pill, check with the doctor before starting treatment with EVRA.

**Switching from a progestogen-only pill, an implant or a contraceptive injection to EVRA:**

- You may start using EVRA any day after stopping the progestogen-only pill or on the day of removal of the implant or on the day you are supposed to inject the contraceptive preparation.**
- Put on the patch the first day after stopping the progestogen-only pill, the day of removing the implant or when you were supposed to inject the contraceptive preparation.
- Use a non-hormonal contraceptive until Day 8, on which you change a patch.

**After a miscarriage or abortion before 20 weeks of pregnancy**

- Please consult the doctor.
- You may start treatment with EVRA right away.

If one or more days have elapsed since your miscarriage or abortion until you start using EVRA, consult the doctor about temporarily using a non-hormonal contraceptive.

**After miscarriage or abortion after 20 weeks of pregnancy**

- Please consult the doctor.
- You may start using EVRA on Day 21 following the miscarriage or abortion, or on the first day of your next period, whichever comes first.

#### After delivery

- Please consult the doctor.
- If you have had a baby and are not breastfeeding, EVRA can only be used after 4 weeks from the delivery have passed.
- If you start using EVRA more than 4 weeks after delivery, please use another non-hormonal contraceptive in addition to EVRA for the first 7 days.

If you have had sex since the delivery, please wait for your first period or see your doctor to make sure you are not pregnant before you start using EVRA.

#### If you are breastfeeding

- Please consult the doctor.
- Do not use this preparation if you are breastfeeding or are planning to breastfeed (see also in section 2 "Pregnancy and breastfeeding").


#### Important information to follow when using the patch:

- Change the EVRA patch on the same day of the week, since the patch is designed to work over 7 days.
- Do not stop using the EVRA patch and remove it for more than 7 days in a row (in the week without a patch).
- Always put on only one patch at a time.
- Do not cut or tamper with the patch.
- Do not put the patch on skin that is red, irritated or cut.
- To work properly, the patch must stick firmly to your skin.
- Press the patch down firmly until the edges stick well.
- Do not use creams, oils, lotions, powder or make-up on the skin where you are placing the patch or near a patch you are already wearing, since this may make the patch fall off the skin.
- Do not put a new patch on the same area of skin from which you removed the old patch. If you put on the patch on the same area, you may cause irritation.
- Check each day that the patch has not fallen off.
- Keep using the patches even if you do not have sex very often.

#### How to use EVRA:


If this is the first time you are using EVRA, wait until the day you get your menstrual period.

- Apply the first patch during the first 24 hours of your period.
- If the patch is first put on after the first day of your period, use a non-hormonal contraceptive until Day 8, on which you change the patch.
- The day you apply the first patch will be Day 1. The "Patch Change Day" will be on this day of the week, every week.***




Choose a place on your body to put the patch.

- Always put the patch on clean, dry and hairless skin.
- The patch can be put on the buttock, abdomen, upper outer arm or upper back only, in places where the patch will not be rubbed by tight clothing.
- Never put the patch on the chest.**



Using your fingers, open the aluminium sachet.


- Open it by tearing it along the edge (do not use scissors).
- Firmly grasp a corner of the patch and gently take it from the sachet.
- There is a clear protective covering on the patch.
- Sometimes patches can stick to the inside of the sachet – be careful not to accidentally remove the clear protective covering as you remove the patch.**



- Take off half of the clear protective covering (see picture).
- Avoid touching the sticky surface of the patch.

Put the patch on your skin.

- Take off the other half of the clear covering.
- Press down firmly on the patch with the palm of your hand for 10 seconds.
- Make sure that the edges stick well.



Leave the patch on for 7 days (one week).

- On the first "Patch Change Day", Day 8, take off the used patch.
- Put on a new patch immediately.
- On Day 15 (Week 3), take off the used patch.
- Put on a new patch immediately.

This makes a total of 3 weeks using the patches.

Day 1	<table border="1"> <tr><td>○</td><td>calendar</td><td>○</td></tr> <tr><td>☑</td><td></td><td>→</td></tr> </table>	○	calendar	○	☑		→	Week 1	<table border="1"> <tr><td>○</td><td>calendar</td><td>○</td></tr> <tr><td>☑</td><td></td><td>→</td></tr> <tr><td>☑</td><td></td><td>→</td></tr> <tr><td>☑</td><td></td><td>→</td></tr> </table>	○	calendar	○	☑		→	☑		→	☑		→	Week 4	<table border="1"> <tr><td>○</td><td>calendar</td><td>○</td></tr> <tr><td>☑</td><td></td><td>→</td>&lt;/</tr></table>	○	calendar	○	☑		→
○	calendar	○																											
☑		→																											
○	calendar	○																											
☑		→																											
☑		→																											
☑		→																											
○	calendar	○																											
☑		→																											