

## 1. NAME OF THE MEDICINAL PRODUCT

AMGEVITA 100 mg/mL - 20 mg solution for injection in pre-filled syringe.

AMGEVITA 100 mg/mL - 40 mg solution for injection in pre-filled syringe.

AMGEVITA 100 mg/mL - 40 mg solution for injection in pre-filled pen.

## 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

### AMGEVITA 100 mg/mL - 20 mg solution for injection in pre-filled syringe

Each single dose pre-filled syringe contains 20 mg of adalimumab in 0.2 mL (100 mg/mL) solution.

### AMGEVITA 100 mg/mL - 40 mg solution for injection in pre-filled syringe

Each single dose pre-filled syringe contains 40 mg of adalimumab in 0.4 mL (100 mg/mL) solution.

### AMGEVITA 100 mg/mL - 40 mg solution for injection in pre-filled pen

Each single dose pre-filled pen contains 40 mg of adalimumab in 0.4 mL (100 mg/mL) solution.

Adalimumab is a recombinant human monoclonal antibody produced in Chinese Hamster Ovary cells.

### Excipients with known effect

AMGEVITA 100 mg/mL - 20 mg: This medicinal product contains 0.2 mg of polysorbate 80 in each 20 mg dose.

AMGEVITA 100 mg/mL - 40 mg: This medicinal product contains 0.4 mg of polysorbate 80 in each 40 mg dose.

For the full list of excipients, see section 6.1.

### **Patient safety information card**

The marketing of AMGEVITA 100 mg/mL is subject to a risk management plan (RMP) including a 'Patient safety information card'. The 'Patient safety information card' emphasizes important safety information that the patient should be aware of before and during treatment. Please explain to the patient the need to review the card before starting treatment.

## 3. PHARMACEUTICAL FORM

AMGEVITA 100 mg/mL - 20 mg solution for injection in pre-filled syringe

AMGEVITA 100 mg/mL - 40 mg solution for injection in pre-filled syringe

Solution for injection (injection).

AMGEVITA 100 mg/mL - 40 mg solution for injection in pre-filled pen

Solution for injection (injection).

Clear and colorless to slightly yellow solution.

AMGEVITA 100 mg/mL is a biosimilar medicinal product, that has been demonstrated to be similar in quality, safety and efficacy to the reference medicinal product Humira. Please be aware of any differences in the indications between the biosimilar medicinal product and the reference medicinal product. Information regarding biosimilar products can be found on the website of the Ministry of Health: <https://www.gov.il/he/pages/biosimilar>.

## 4. CLINICAL PARTICULARS

### 4.1 Therapeutic indications

#### Rheumatoid arthritis

AMGEVITA 100 mg/mL in combination with methotrexate is indicated for:

- The treatment of moderate to severe, active rheumatoid arthritis in adult patients when the response to disease-modifying anti-rheumatic drugs (DMARDs) including methotrexate has been inadequate.
- The treatment of severe, active and progressive rheumatoid arthritis in adults not previously treated with methotrexate.

AMGEVITA 100 mg/mL can be given as monotherapy in case of intolerance to methotrexate or when continued treatment with methotrexate is inappropriate.

AMGEVITA 100 mg/mL has been shown to reduce the rate of progression of joint damage as measured by X-ray and to improve physical function, when given in combination with methotrexate.

#### Juvenile idiopathic arthritis

##### *Polyarticular juvenile idiopathic arthritis*

AMGEVITA 100 mg/mL in combination with methotrexate is indicated for the treatment of active polyarticular juvenile idiopathic arthritis, in patients from the age of 2 years who have had an inadequate response to one or more disease-modifying anti-rheumatic drugs. AMGEVITA 100 mg/mL can be given as monotherapy in case of intolerance to methotrexate or when continued treatment with methotrexate is inappropriate (for the efficacy in monotherapy, see section 5.1). AMGEVITA 100 mg/mL has not been studied in patients aged less than 2 years.

##### *Enthesitis-related arthritis*

AMGEVITA 100 mg/mL is indicated for the treatment of active enthesitis-related arthritis in patients, 6 years of age and older, who have had an inadequate response to, or who are intolerant of, conventional therapy (see section 5.1).

#### Axial spondyloarthritis

##### *Ankylosing spondylitis (AS):*

AMGEVITA 100 mg/mL is indicated for the treatment of adults with severe active ankylosing spondylitis who have had an inadequate response to conventional therapy.

##### *Axial spondyloarthritis without radiographic evidence of AS:*

AMGEVITA 100 mg/mL is indicated for the treatment of adults with severe axial spondyloarthritis without radiographic evidence of AS, but with objective signs of inflammation by radiological and/or laboratory tests including MRI and serum CRP levels, who have had an inadequate response to, or are intolerant to, non - steroidal anti-inflammatory drugs.

### Psoriatic arthritis

AMGEVITA 100 mg/mL is indicated for the treatment of active and progressive psoriatic arthritis in adults when the response to previous disease-modifying anti-rheumatic drug therapy has been inadequate. Adalimumab has been shown to reduce the rate of progression of peripheral joint damage as measured by X-ray in patients with polyarticular symmetrical subtypes of the disease (see section 5.1) and to improve physical function.

### Psoriasis

AMGEVITA 100 mg/mL is indicated for the treatment of moderate to severe chronic plaque psoriasis in adult patients who are candidates for systemic therapy.

### Pediatric plaque psoriasis

AMGEVITA 100 mg/mL is indicated for the treatment of severe chronic plaque psoriasis in children and adolescents from 4 years of age who have had an inadequate response to or are inappropriate candidates for topical therapy and phototherapies.

### Hidradenitis suppurativa (HS)

AMGEVITA 100 mg/mL is indicated for the treatment of active moderate to severe hidradenitis suppurativa (acne inversa) in adults and adolescents from 12 years of age with an inadequate response to conventional systemic HS therapy (see sections 5.1 and 5.2).

### Crohn's disease

AMGEVITA 100 mg/mL is indicated for reducing signs and symptoms and inducing and maintaining clinical remission in adult patients with moderately to severely active Crohn's disease who have had an inadequate response to conventional therapy. AMGEVITA 100 mg/mL is indicated for reducing signs and symptoms and inducing clinical remission in these patients if they have also lost response to or are intolerant to infliximab.

### Pediatric Crohn's disease

AMGEVITA 100 mg/mL is indicated for the treatment of moderately to severely active Crohn's disease in pediatric patients (from 6- years of age) who have had an inadequate response to conventional therapy including primary nutrition therapy and corticosteroid, and/or an immunomodulator, or who are intolerant to or have contraindications for such therapies.

### Ulcerative colitis

AMGEVITA 100 mg/mL is indicated for treatment of moderately to severely active ulcerative colitis in adult patients who have had an inadequate response to conventional therapy including corticosteroids and 6-mercaptopurine (6-MP) or azathioprine (AZA), or who are intolerant to or have medical contraindications for such therapies.

### Pediatric ulcerative colitis

AMGEVITA 100 mg/mL is indicated for the treatment of moderately to severely active ulcerative colitis in pediatric patients (from 6 years of age) who have had an inadequate response to conventional therapy including corticosteroids and/or 6-mercaptopurine (6-MP) or azathioprine (AZA), or who are intolerant to or have medical contraindications for such therapies.

### Uveitis

AMGEVITA 100 mg/mL is indicated for the treatment of non-infectious intermediate, posterior and panuveitis in adult patients who have had an inadequate response to corticosteroids, in patients in need of corticosteroid-sparing, or in whom corticosteroid treatment is inappropriate.

### Pediatric Uveitis

AMGEVITA 100 mg/mL is indicated for the treatment of chronic non-infectious uveitis in pediatric patients from 2 years of age who have had an inadequate response to or are intolerant to conventional therapy, or in whom conventional therapy is inappropriate.

### Intestinal Behcet's disease

AMGEVITA 100 mg/mL is indicated for the treatment of intestinal Behcet's disease in patients who have had an inadequate response to conventional therapy.

## **4.2 Posology and method of administration**

AMGEVITA 100 mg/mL treatment should be initiated and supervised by specialist physicians experienced in the diagnosis and treatment of conditions for which AMGEVITA 100 mg/mL is indicated. Ophthalmologists are advised to consult with an appropriate specialist before initiation of treatment with AMGEVITA 100 mg/mL (see section 4.4). Patients treated with AMGEVITA 100 mg/mL should be given the 'Patient safety information card'.

After proper training in injection technique, patients may self-inject with AMGEVITA 100 mg/mL if their physician determines that it is appropriate and with medical follow-up as necessary.

During treatment with AMGEVITA 100 mg/mL, other concomitant therapies (e.g. corticosteroids and/or immunomodulatory agents) should be optimized.

### Posology

#### *Rheumatoid arthritis*

The recommended dose of AMGEVITA 100 mg/mL for adult patients with rheumatoid arthritis is 40 mg adalimumab administered every other week as a single dose via subcutaneous injection. Methotrexate should be continued during treatment with AMGEVITA 100 mg/mL.

Glucocorticoids, salicylates, non-steroidal anti-inflammatory drugs, or analgesics can be continued during treatment with AMGEVITA 100 mg/mL. Regarding combination with disease-modifying anti-rheumatic drugs other than methotrexate, see sections 4.4 and 5.1.

In monotherapy, some patients who experience a decrease in their response to AMGEVITA 100 mg/mL 40 mg every other week may benefit from an increase in dosage to 40 mg adalimumab every week or 80 mg every other week.

Available data suggest that the clinical response is usually achieved within 12 weeks of treatment. Continued therapy should be reconsidered in a patient not responding within this time period.

#### *Dose interruption*

There may be a need for dose interruption, for instance before surgery or if a serious infection occurs.

Available data suggest that re-introduction of AMGEVITA 100 mg/mL after discontinuation for 70 days or longer resulted in the same magnitudes of clinical response and similar safety profile as before dose interruption.

#### *Ankylosing spondylitis, axial spondyloarthritis without radiographic evidence of AS and psoriatic arthritis*

The recommended dose of AMGEVITA 100 mg/mL for patients with ankylosing spondylitis, axial spondyloarthritis without radiographic evidence of AS and for patients with psoriatic arthritis is 40 mg adalimumab administered every other week as a single dose via subcutaneous injection.

Available data suggest that the clinical response is usually achieved within 12 weeks of treatment. Continued therapy should be reconsidered in a patient not responding within this time period.

### *Psoriasis*

The recommended dose of AMGEVITA 100 mg/mL for adult patients is an initial dose of 80 mg administered subcutaneously, followed by 40 mg subcutaneously given every other week starting one week after the initial dose.

Continued therapy beyond 16 weeks should be carefully reconsidered in a patient not responding within this time period.

Beyond 16 weeks, patients with inadequate response to AMGEVITA 100 mg/mL 40 mg every other week may benefit from an increase in dosage to 40 mg every week or 80 mg every other week. The benefits and risks of continued 40 mg weekly or 80 mg every other week therapy should be carefully reconsidered in a patient with an inadequate response after the increase in dosage (see section 5.1). If adequate response is achieved with 40 mg every week or 80 mg every other week, the dosage may subsequently be reduced to 40 mg every other week.

### *Hidradenitis suppurativa*

The recommended AMGEVITA 100 mg/mL dose regimen for adult patients with hidradenitis suppurativa (HS) is 160 mg initially at day 1 (given as four 40 mg or two 80 mg injections in one day or as two 40 mg or one 80 mg injections per day for two consecutive days), followed by 80 mg two weeks later at day 15 (given as two 40 mg or one 80 mg injections in one day). Two weeks later (day 29) continue with a dose of 40 mg every week or 80 mg every other week (given as two 40 mg injections in one day). Antibiotics may be continued during treatment with AMGEVITA 100 mg/mL if necessary. It is recommended that the patient should use a topical antiseptic wash on their HS lesions on a daily basis during treatment with AMGEVITA 100 mg/mL.

Continued therapy beyond 12 weeks should be carefully reconsidered in a patient with no improvement within this time period.

Should treatment be interrupted, AMGEVITA 100 mg/mL 40 mg every week or 80 mg every other week may be re-introduced (see section 5.1).

The benefit and risk of continued long-term treatment should be periodically evaluated (see section 5.1).

### *Crohn's disease*

The recommended AMGEVITA 100 mg/mL induction dose regimen for adult patients with moderately to severely active Crohn's disease is 160 mg at week 0 (given as four 40 mg or two 80 mg injections in one day or as two 40 mg or one 80 mg injections per day for two consecutive days), followed by 80 mg at week 2 (given as two 40 mg or one 80 mg injections in one day), followed by a maintenance dose of 40 mg every other week via subcutaneous injection beginning at week 4.

Aminosalicylates, corticosteroids and/or immunomodulatory agents (e.g. 6-mercaptopurine and azathioprine) may be continued during treatment with AMGEVITA 100 mg/mL.

Some patients who experience decrease in their response to AMGEVITA 100 mg/mL 40 mg every other week may benefit from an increase in dosage to 40 mg AMGEVITA 100 mg/mL every week or 80 mg every other week.

Some patients who have not responded by week 4 may benefit from continued maintenance therapy through week 12. Continued therapy should be carefully reconsidered in a patient not responding within this time period.

### *Ulcerative colitis*

The recommended AMGEVITA 100 mg/mL induction dose regimen for adult patients with moderate to severe ulcerative colitis is 160 mg at week 0 (given as four 40 mg or two 80 mg injections in one

day or as two 40 mg or one 80 mg injections per day for two consecutive days) and 80 mg at week 2 (given as two 40 mg or one 80 mg injections in one day). After induction treatment, the recommended dose is 40 mg every other week via subcutaneous injection.

During maintenance treatment, corticosteroids may be tapered in accordance with clinical practice guidelines.

Some patients who experience decrease in their response to AMGEVITA 100 mg/mL 40 mg every other week may benefit from an increase in dosage to 40 mg AMGEVITA 100 mg/mL every week or 80 mg every other week.

Available data suggest that clinical response is usually achieved within 2-8 weeks of treatment. AMGEVITA 100 mg/mL therapy should not be continued in patients failing to respond within this time period.

#### *Uveitis*

The recommended dose of AMGEVITA 100 mg/mL for adult patients with uveitis is an initial dose of 80 mg, followed by 40 mg given every other week starting one week after the initial dose. There is limited experience in the initiation of treatment with adalimumab alone. Treatment with AMGEVITA 100 mg/mL can be initiated in combination with corticosteroids and/or with other non-biologic immunomodulatory agents. Concomitant corticosteroids may be tapered in accordance with clinical practice starting two weeks after initiating treatment with AMGEVITA 100 mg/mL.

It is recommended that the benefit and risk of continued long-term treatment should be evaluated on a yearly basis (see section 5.1).

#### *Intestinal Behcet's disease*

The initial dose of AMGEVITA 100 mg/mL for adult intestinal Behcet's disease patients is 160 mg as subcutaneous injection. The initial dose is followed by 80 mg two weeks later. From four weeks after the initial dose, 40 mg is administered every other week.

AMGEVITA 100 mg/mL should be used when the signs and symptoms caused by intestinal Behcet's disease remain clearly even if patients have appropriate treatment with existing drug (steroids or immunomodulator, etc.).

Continued therapy with the same regimen should be carefully reconsidered in a patient not responding such as clinical symptoms and/or endoscopic findings within 12 weeks of treatment.

#### Special populations

##### Elderly

No dose adjustment is required.

##### Renal and/or hepatic impairment

Adalimumab has not been studied in these patient populations. No dose recommendations can be made.

##### Pediatric population:

##### *Juvenile idiopathic arthritis*

##### *Polyarticular juvenile idiopathic arthritis from 2 years of age*

The recommended dose of AMGEVITA 100 mg/mL for patients with polyarticular juvenile idiopathic arthritis from 2 years of age is based on body weight (Table 1). AMGEVITA 100 mg/mL is administered every other week via subcutaneous injection.

**Table 1. AMGEVITA 100 mg/mL dose for patients with Polyarticular Juvenile Idiopathic Arthritis**

<b>Patient Weight</b>	<b>Dosing Regimen</b>
10 kg to < 30 kg	20 mg every other week
≥ 30 kg	40 mg every other week

Available data suggest that clinical response is usually achieved within 12 weeks of treatment. Continued therapy should be carefully reconsidered in a patient not responding within this time period.

There is no relevant use of AMGEVITA 100 mg/mL in patients aged less than 2 years for this indication.

*Enthesitis-related arthritis*

The recommended dose of AMGEVITA 100 mg/mL for patients with enthesitis-related arthritis from 6 years of age is based on body weight (Table 2). AMGEVITA 100 mg/mL is administered every other week via subcutaneous injection.

**Table 2. AMGEVITA 100 mg/mL dose for patients with Enthesitis-Related Arthritis**

<b>Patient Weight</b>	<b>Dosing Regimen</b>
15 kg to < 30 kg	20 mg every other week
≥ 30 kg	40 mg every other week

AMGEVITA 100 mg/mL has not been studied in patients with enthesitis-related arthritis aged less than 6 years.

*Psoriatic arthritis and axial spondyloarthritis including ankylosing spondylitis*

There is no relevant use of AMGEVITA 100 mg/mL in the pediatric population for the indications of ankylosing spondylitis and psoriatic arthritis.

*Pediatric plaque psoriasis*

The recommended AMGEVITA 100 mg/mL dose for patients with plaque psoriasis from 4 to 17 years of age is based on body weight (Table 3). AMGEVITA 100 mg/mL is administered via subcutaneous injection.

**Table 3. AMGEVITA 100 mg/mL dose for pediatric patients with Plaque Psoriasis**

<b>Patient Weight</b>	<b>Dosing Regimen</b>
15 kg to < 30 kg	Initial dose of 20 mg, followed by 20 mg given every other week starting one week after the initial dose
≥ 30 kg	Initial dose of 40 mg, followed by 40 mg given every other week starting one week after the initial dose

Continued therapy beyond 16 weeks should be carefully considered in a patient not responding within this time period.

If retreatment with AMGEVITA 100 mg/mL is indicated, the above guidance on dose and treatment duration should be followed.

The safety of AMGEVITA 100 mg/mL in pediatric patients with plaque psoriasis has been assessed for a mean of 13 months.

There is no relevant use of AMGEVITA 100 mg/mL in children aged less than 4 years for this indication.

*Adolescent hidradenitis suppurativa (from 12 years of age, weighing at least 30 kg)*

There are no clinical trials with AMGEVITA 100 mg/mL in adolescent patients with HS. The posology of AMGEVITA 100 mg/mL in these patients has been determined from pharmacokinetic modelling and simulation (see section 5.2).

The recommended AMGEVITA 100 mg/mL dose is 80 mg at Week 0 followed by 40 mg every other week starting at Week 1 via subcutaneous injection.

In adolescent patients with inadequate response to AMGEVITA 100 mg/mL 40 mg every other week, an increase in dosage to 40 mg every week or 80 mg every other week may be considered.

Antibiotics may be continued during treatment with AMGEVITA 100 mg/mL if necessary. It is recommended that the patient should use a topical antiseptic wash on their HS lesions on a daily basis during treatment with AMGEVITA 100 mg/mL.

Continued therapy beyond 12 weeks should be carefully reconsidered in a patient with no improvement within this time period.

Should treatment be interrupted, AMGEVITA 100 mg/mL may be re-introduced as appropriate.

The benefit and risk of continued long-term treatment should be periodically evaluated (see adult data in section 5.1)

There is no relevant use of AMGEVITA 100 mg/mL in children aged less than 12 years in this indication.

*Pediatric Crohn's disease*

The recommended dose of AMGEVITA 100 mg/mL for patients with Crohn's disease from 6 to 17 years of age is based on body weight (Table 4). AMGEVITA 100 mg/mL is administered via subcutaneous injection.

**Table 4. AMGEVITA 100 mg/mL dose for pediatric patients with Crohn's disease**

<b>Patient Weight</b>	<b>Induction Dose</b>	<b>Maintenance Dose Starting at Week 4</b>
< 40 kg	<ul style="list-style-type: none"><li>• 40 mg at week 0 and 20 mg at week 2</li></ul> <p>In case there is a need for a more rapid response to therapy with the awareness that the risk for adverse events may be higher with use of the higher induction dose, the following dose may be used:</p> <ul style="list-style-type: none"><li>• 80 mg at week 0 and 40 mg at week 2</li></ul>	20 mg every other week
≥ 40 kg	<ul style="list-style-type: none"><li>• 80 mg at week 0 and 40 mg at week 2</li></ul> <p>In case there is a need for a more rapid response to therapy with the awareness that the risk for adverse events may be higher with use of the higher induction dose, the following dose may be used:</p> <ul style="list-style-type: none"><li>• 160 mg at week 0 and 80 mg at week 2</li></ul>	40 mg every other week

Patients who experience insufficient response may benefit from an increase in dosage:

- < 40 kg: 20 mg every week
- ≥ 40 kg: 40 mg every week or 80 mg every other week

Continued therapy should be carefully considered in a subject not responding by week 12.

There is no relevant use of AMGEVITA 100 mg/mL in children aged less than 6 years for this indication.

### *Pediatric ulcerative colitis*

The recommended dose of AMGEVITA 100 mg/mL for patients from 6 to 17 years of age with ulcerative colitis is based on body weight (Table 5). AMGEVITA 100 mg/mL is administered via subcutaneous injection.

**Table 5. AMGEVITA 100 mg/mL dose for pediatric patients with Ulcerative Colitis**

<b>Patient Weight</b>	<b>Induction Dose</b>	<b>Maintenance Dose Starting at Week 4*</b>
< 40 kg	<ul style="list-style-type: none"><li>• 80 mg at Week 0 (given as two 40 mg injections in one day) and</li><li>• 40 mg at Week 2 (given as one 40 mg injection)</li></ul>	<ul style="list-style-type: none"><li>• 40 mg every other week</li></ul>
≥ 40 kg	<ul style="list-style-type: none"><li>• 160 mg at Week 0 (given as four 40 mg injections in one day or two 40 mg injections per day for two consecutive days) and</li><li>• 80 mg at Week 2 (given as two 40 mg injections in one day)</li></ul>	<ul style="list-style-type: none"><li>• 80 mg every other week</li></ul>

\* Pediatric patients who turn 18 years of age while on AMGEVITA 100 mg/mL should continue their prescribed maintenance dose.

Continued therapy beyond 8 weeks should be carefully considered in patients not showing signs of response within this time period.

There is no relevant use of AMGEVITA 100 mg/mL in children aged less than 6 years in this indication.

### *Pediatric Uveitis*

The recommended dose of AMGEVITA 100 mg/mL for pediatric patients with uveitis from 2 years of age is based on body weight (Table 6). AMGEVITA 100 mg/mL is administered via subcutaneous injection.

In pediatric uveitis, there is no experience in the treatment with AMGEVITA 100 mg/mL without concomitant treatment with methotrexate.

**Table 6. AMGEVITA 100 mg/mL dose for pediatric patients with Uveitis**

<b>Patient Weight</b>	<b>Dosing Regimen</b>
< 30 kg	20 mg every other week in combination with methotrexate
≥ 30 kg	40 mg every other week in combination with methotrexate

When AMGEVITA 100 mg/mL therapy is initiated, a loading dose of 40 mg for patients < 30 kg or 80 mg for patients ≥ 30 kg may be administered one week prior to the start of maintenance therapy. No clinical data are available on the use of AMGEVITA 100 mg/mL loading dose in children < 6 years of age (see section 5.2).

There is no relevant use of AMGEVITA 100 mg/mL in children aged less than 2 years in this indication.

It is recommended that the benefit and risk of continued long-term treatment should be evaluated on a yearly basis (see section 5.1).

### Method of administration

AMGEVITA 100 mg/mL is administered by subcutaneous injection. Full instructions for use are provided in the package leaflet.

### 4.3 Contraindications

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

Active tuberculosis or other severe infections such as sepsis, and opportunistic infections (see section 4.4).

Moderate to severe heart failure (NYHA class III/IV) (see section 4.4).

### 4.4 Special warnings and precautions for use

#### Traceability

In order to improve traceability of biological medicinal products, the name and the batch number of the administered product should be clearly recorded.

#### Infections

Patients taking TNF-antagonists are more susceptible to serious infections. Impaired lung function may increase the risk for developing infections. Patients must therefore be monitored closely for infections, including tuberculosis, before, during and after treatment with AMGEVITA 100 mg/mL. Because the elimination of adalimumab may take up to four months, monitoring should be continued throughout this period.

Treatment with AMGEVITA 100 mg/mL should not be initiated in patients with active infections including chronic or localized infections until infections are controlled. In patients who have been exposed to tuberculosis and patients who have travelled in areas of high risk of tuberculosis or endemic mycoses, such as histoplasmosis, coccidioidomycosis, or blastomycosis, the risk and benefits of treatment with AMGEVITA 100 mg/mL should be considered prior to initiating therapy (see Other opportunistic infections).

Patients who develop a new infection while undergoing treatment with AMGEVITA 100 mg/mL, should be monitored closely and undergo a complete diagnostic evaluation. Administration of AMGEVITA 100 mg/mL should be discontinued if a patient develops a new serious infection or sepsis, and appropriate antimicrobial or antifungal therapy should be initiated until the infection is controlled. Physicians should exercise caution when considering the use of AMGEVITA 100 mg/mL in patients with a history of recurring infection or with underlying conditions which may predispose patients to infections, including the use of concomitant immunosuppressive medications.

#### *Serious infections*

Serious infections, including sepsis, due to bacterial, mycobacterial, invasive fungal, parasitic, viral, or other opportunistic infections such as listeriosis, legionellosis and pneumocystis have been reported in patients receiving adalimumab.

Other serious infections seen in clinical trials include pneumonia, pyelonephritis, septic arthritis and septicemia. Hospitalization or fatal outcomes associated with infections have been reported.

#### *Tuberculosis*

Tuberculosis, including reactivation and new onset of tuberculosis, has been reported in patients receiving adalimumab. Reports included cases of pulmonary and extra-pulmonary (i.e. disseminated) tuberculosis.

Before initiation of therapy with AMGEVITA 100 mg/mL, all patients must be evaluated for both active or inactive ("latent") tuberculosis infection. This evaluation should include a detailed medical assessment of patient history of tuberculosis or possible previous exposure to people with active tuberculosis and previous and/or current immunosuppressive therapy. Appropriate screening tests (i.e. tuberculin skin test and chest X-ray) should be performed in all patients (local recommendations may

apply). It is recommended that the conduct and results of these tests are recorded in the 'Patient safety information card'. Prescribers are reminded of the risk of false negative tuberculin skin test results, especially in patients who are severely ill or immunocompromised.

If active tuberculosis is diagnosed, AMGEVITA 100 mg/mL therapy must not be initiated (see section 4.3).

In all situations described below, the benefit/risk balance of therapy should be very carefully considered.

If latent tuberculosis is suspected, a physician with expertise in the treatment of tuberculosis should be consulted.

If latent tuberculosis is diagnosed, appropriate treatment must be started with anti-tuberculosis prophylaxis treatment before the initiation of AMGEVITA 100 mg/mL, and in accordance with local recommendations.

Use of anti-tuberculosis prophylaxis treatment should also be considered before the initiation of AMGEVITA 100 mg/mL in patients with several or significant risk factors for tuberculosis despite a negative test for tuberculosis and in patients with a past history of latent or active tuberculosis in whom an adequate course of treatment cannot be confirmed.

Despite prophylactic treatment for tuberculosis, cases of reactivated tuberculosis have occurred in patients treated with adalimumab. Some patients who have been successfully treated for active tuberculosis have redeveloped tuberculosis while being treated with adalimumab.

Patients should be instructed to seek medical advice if signs/symptoms suggestive of a tuberculosis infection (e.g. persistent cough, wasting/weight loss, low grade fever, listlessness) occur during or after therapy with AMGEVITA 100 mg/mL.

#### *Other opportunistic infections*

Opportunistic infections, including invasive fungal infections have been observed in patients receiving adalimumab. These infections have not consistently been recognized in patients taking TNF-antagonists and this has resulted in delays in appropriate treatment, sometimes resulting in fatal outcomes.

For patients who develop the signs and symptoms such as fever, malaise, weight loss, sweats, cough, dyspnea, and/or pulmonary infiltrates or other serious systemic illness with or without concomitant shock an invasive fungal infection should be suspected and administration of AMGEVITA 100 mg/mL should be promptly discontinued. Diagnosis and administration of empiric antifungal therapy in these patients should be made in consultation with a physician with expertise in the care of patients with invasive fungal infections.

#### Hepatitis B reactivation

Reactivation of hepatitis B has occurred in patients receiving a TNF-antagonist including adalimumab, who are chronic carriers of this virus (i.e. surface antigen positive). Some cases have had a fatal outcome. Patients should be tested for HBV infection before initiating treatment with AMGEVITA 100 mg/mL. For patients who test positive for hepatitis B infection, consultation with a physician with expertise in the treatment of hepatitis B is recommended.

Carriers of HBV who require treatment with AMGEVITA 100 mg/mL should be closely monitored for signs and symptoms of active HBV infection throughout therapy and for several months following termination of therapy. Adequate data from treating patients who are carriers of HBV with anti-viral therapy in conjunction with TNF-antagonist therapy to prevent HBV reactivation are not available. In

patients who develop HBV reactivation, AMGEVITA 100 mg/mL should be stopped and effective anti-viral therapy with appropriate supportive treatment should be initiated.

### Neurological events

TNF-antagonists including adalimumab have been associated in rare instances with new onset or exacerbation of clinical symptoms and/or radiographic evidence of central nervous system demyelinating disease including multiple sclerosis and optic neuritis, and peripheral demyelinating disease, including Guillain-Barré syndrome. Prescribers should exercise caution in considering the use of AMGEVITA 100 mg/mL in patients with pre-existing or recent-onset central or peripheral nervous system demyelinating disorders; discontinuation of AMGEVITA 100 mg/mL should be considered if any of these disorders develop. There is a known association between intermediate uveitis and central demyelinating disorders. Neurologic evaluation should be performed in patients with non-infectious intermediate uveitis prior to the initiation of AMGEVITA 100 mg/mL therapy and regularly during treatment to assess for pre-existing or developing central demyelinating disorders.

### Allergic reactions

Serious allergic reactions associated with adalimumab were rare during clinical trials. Non-serious allergic reactions associated with adalimumab were uncommon during clinical trials. Reports of serious allergic reactions including anaphylaxis have been received following adalimumab administration. If an anaphylactic reaction or other serious allergic reaction occurs, administration of AMGEVITA 100 mg/mL should be discontinued immediately and appropriate therapy initiated.

### Immunosuppression

In a study of 64 patients with rheumatoid arthritis that were treated with adalimumab, there was no evidence of depression of delayed-type hypersensitivity, depression of immunoglobulin levels, or change in enumeration of effector T-, B-, NK-cells, monocyte/macrophages, and neutrophils.

### Malignancies and lymphoproliferative disorders

In the controlled portions of clinical trials of TNF-antagonists, more cases of malignancies including lymphoma have been observed among patients receiving a TNF-antagonist compared with control patients. However, the occurrence was rare. In the post-marketing setting, cases of leukemia have been reported in patients treated with a TNF-antagonist. There is an increased background risk for lymphoma and leukemia in rheumatoid arthritis patients with long-standing highly active, inflammatory disease, which complicates the risk estimation. With the current knowledge, a possible risk for the development of lymphomas, leukemia, and other malignancies in patients treated with a TNF-antagonist cannot be excluded.

Malignancies, some fatal, have been reported among children, adolescents and young adults (up to 22 years of age) treated with TNF-antagonists (initiation of therapy  $\leq$  18 years of age), including adalimumab in the post-marketing setting. Approximately half the cases were lymphomas. The other cases represented a variety of different malignancies and included rare malignancies usually associated with immunosuppression. A risk for the development of malignancies in children and adolescents treated with TNF-antagonists cannot be excluded.

Rare post-marketing cases of hepatosplenic T-cell lymphoma have been identified in patients treated with adalimumab. This rare type of T-cell lymphoma has a very aggressive disease course and is usually fatal. Some of these hepatosplenic T-cell lymphomas with adalimumab have occurred in young adult patients on concomitant treatment with azathioprine or 6-mercaptopurine used for inflammatory bowel disease. The potential risk with the combination of azathioprine or 6-mercaptopurine and AMGEVITA 100 mg/mL should be carefully considered. A risk for the development of hepatosplenic T-cell lymphoma in patients treated with AMGEVITA 100 mg/mL cannot be excluded (see section 4.8).

No studies have been conducted that include patients with a history of malignancy or in whom treatment with adalimumab is continued following development of malignancy. Thus additional caution should be exercised in considering AMGEVITA 100 mg/mL treatment of these patients (see section 4.8).

All patients, and in particular patients with a medical history of extensive immunosuppressant therapy or psoriasis patients with a history of PUVA treatment should be examined for the presence of non-melanoma skin cancer prior to and during treatment with AMGEVITA 100 mg/mL. Melanoma and Merkel cell carcinoma have also been reported in patients treated with TNF-antagonists including adalimumab (see section 4.8).

In an exploratory clinical trial evaluating the use of another TNF-antagonist, infliximab, in patients with moderate to severe chronic obstructive pulmonary disease (COPD), more malignancies, mostly in the lung or head and neck, were reported in infliximab-treated patients compared with control patients. All patients had a history of heavy smoking. Therefore, caution should be exercised when using any TNF-antagonist in COPD patients, as well as in patients with increased risk for malignancy due to heavy smoking.

With current data it is not known if adalimumab treatment influences the risk for developing dysplasia or colon cancer. All patients with ulcerative colitis who are at increased risk for dysplasia or colon carcinoma (for example, patients with long-standing ulcerative colitis or primary sclerosing cholangitis), or who had a prior history of dysplasia or colon carcinoma should be screened for dysplasia at regular intervals before therapy and throughout their disease course. This evaluation should include colonoscopy and biopsies per local recommendations.

#### Hematologic reactions

Rare reports of pancytopenia including aplastic anemia have been reported with TNF-antagonists. Adverse events of the hematologic system, including medically significant cytopenia (e.g. thrombocytopenia, leukopenia) have been reported with adalimumab. All patients should be advised to seek immediate medical attention if they develop signs and symptoms suggestive of blood dyscrasias (e.g. persistent fever, bruising, bleeding, pallor) while on AMGEVITA 100 mg/mL. Discontinuation of AMGEVITA 100 mg/mL therapy should be considered in patients with confirmed significant hematologic abnormalities.

#### Vaccinations

Similar antibody responses to the standard 23-valent pneumococcal vaccine and the influenza trivalent virus vaccination were observed in a study in 226 adult subjects with rheumatoid arthritis who were treated with adalimumab or placebo. No data are available on the secondary transmission of infection by live vaccines in patients receiving adalimumab.

It is recommended that pediatric patients, if possible, be brought up to date with all immunizations in agreement with current immunization guidelines prior to initiating AMGEVITA 100 mg/mL therapy.

Patients on AMGEVITA 100 mg/mL may receive concurrent vaccinations, except for live vaccines. Administration of live vaccines (e.g. BCG vaccine) to infants exposed to AMGEVITA 100 mg/mL *in utero* is not recommended for 5 months following the mother's last AMGEVITA 100 mg/mL injection during pregnancy.

#### Congestive heart failure

In a clinical trial with another TNF-antagonist worsening congestive heart failure and increased mortality due to congestive heart failure have been observed. Cases of worsening congestive heart failure have also been reported in patients receiving adalimumab. AMGEVITA 100 mg/mL should be used with caution in patients with mild heart failure (NYHA class I/II). AMGEVITA 100 mg/mL is contraindicated in moderate to severe heart failure (see section 4.3). Treatment with AMGEVITA

100 mg/mL must be discontinued in patients who develop new or worsening symptoms of congestive heart failure.

#### Autoimmune processes

Treatment with AMGEVITA 100 mg/mL may result in the formation of autoimmune antibodies. The impact of long-term treatment with AMGEVITA 100 mg/mL on the development of autoimmune diseases is unknown. If a patient develops symptoms suggestive of a lupus-like syndrome following treatment with AMGEVITA 100 mg/mL and is positive for antibodies against double-stranded DNA, further treatment with AMGEVITA 100 mg/mL should not be given (see section 4.8).

#### Concurrent administration of biologic DMARDs or TNF-antagonists

Serious infections were seen in clinical studies with concurrent use of anakinra and another TNF-antagonist, etanercept, with no added clinical benefit compared to etanercept alone. Because of the nature of the adverse events seen with the combination of etanercept and anakinra therapy, similar toxicities may also result from the combination of anakinra and other TNF-antagonists. Therefore, the combination of AMGEVITA 100 mg/mL and anakinra is not recommended (see section 4.5).

Concomitant administration of AMGEVITA 100 mg/mL with other biologic DMARDs (e.g. anakinra and abatacept) or other TNF-antagonists is not recommended based upon the possible increased risk for infections, including serious infections and other potential pharmacological interactions (see section 4.5).

#### Surgery

There is limited safety experience of surgical procedures in patients treated with adalimumab. The long half-life of adalimumab should be taken into consideration if a surgical procedure is planned. A patient who requires surgery while on AMGEVITA 100 mg/mL should be closely monitored for infections and appropriate actions should be taken. There is limited safety experience in patients undergoing arthroplasty while receiving adalimumab.

#### Small bowel obstruction

Failure to respond to treatment for Crohn's disease may indicate the presence of fixed fibrotic stricture that may require surgical treatment. Available data suggest that adalimumab does not worsen or cause strictures.

#### Elderly

The frequency of serious infections among adalimumab-treated subjects over 65 years of age (3.7%) was higher than for those under 65 years of age (1.5%). Some of those had a fatal outcome. Particular attention regarding the risk for infection should be paid when treating the elderly.

#### Excipients with known effects

This medicinal product contains less than 1 mmol of sodium (23 mg) per 0.8 mL dose, i.e. essentially 'sodium-free'.

#### Pediatric population

See Vaccinations above.

#### Excipients with known effect

AMGEVITA 100 mg/mL - 20 mg: This medicinal product contains 0.2 mg of polysorbate 80 in each 20 mg dose. Polysorbates may cause allergic reactions.

AMGEVITA 100 mg/mL - 40 mg: This medicinal product contains 0.4 mg of polysorbate 80 in each 40 mg dose. Polysorbates may cause allergic reactions.

#### 4.5 Interaction with other medicinal products and other forms of interaction

Adalimumab has been studied in rheumatoid arthritis, polyarticular juvenile idiopathic arthritis and psoriatic arthritis patients taking adalimumab as monotherapy and those taking concomitant methotrexate. Antibody formation was lower when adalimumab was given together with methotrexate in comparison with use as monotherapy. Administration of adalimumab without methotrexate resulted in increased formation of antibodies, increased clearance and reduced efficacy of adalimumab (see section 5.1).

The combination of AMGEVITA 100 mg/mL and anakinra is not recommended (see section 4.4 “Concurrent administration of biologic DMARDs or TNF-antagonists”).

The combination of AMGEVITA 100 mg/mL and abatacept is not recommended (see section 4.4 “Concurrent administration of biologic DMARDs or TNF-antagonists”).

#### 4.6 Fertility, pregnancy and lactation

##### Women of childbearing potential

Women of childbearing potential should consider the use of adequate contraception to prevent pregnancy and continue its use for at least five months after the last AMGEVITA 100 mg/mL treatment.

##### Pregnancy

A large number (approximately 2,100) of prospectively collected pregnancies exposed to adalimumab resulting in live birth with known outcomes, including more than 1,500 exposed during the first trimester, does not indicate an increase in the rate of malformation in the newborn.

In a prospective cohort registry, 257 women with rheumatoid arthritis (RA) or Crohn’s disease (CD) treated with adalimumab at least during the first trimester and 120 women with RA or CD not treated with adalimumab were enrolled. The primary endpoint was the birth prevalence of major birth defects. The rate of pregnancies ending with at least one live born infant with a major birth defect was 6/69 (8.7%) in the adalimumab-treated women with RA and 5/74 (6.8%) in the untreated women with RA (unadjusted OR 1.31, 95% CI 0.38-4.52) and 16/152 (10.5%) in the adalimumab-treated women with CD and 3/32 (9.4%) in the untreated women with CD (unadjusted OR 1.14, 95% CI 0.31-4.16). The adjusted OR (accounting for baseline differences) was 1.10 (95% CI 0.45-2.73) with RA and CD combined. There were no distinct differences between adalimumab-treated and untreated women for the secondary endpoints spontaneous abortions, minor birth defects, preterm delivery, birth size and serious or opportunistic infections and no stillbirths or malignancies were reported. The interpretation of data may be impacted due to methodological limitations of the study, including small sample size and non-randomized design.

In a developmental toxicity study conducted in monkeys, there was no indication of maternal toxicity, embryotoxicity or teratogenicity. Preclinical data on postnatal toxicity of adalimumab are not available (see section 5.3).

Due to its inhibition of TNF $\alpha$ , adalimumab administered during pregnancy could affect normal immune responses in the newborn. AMGEVITA 100 mg/mL should only be used during pregnancy if clearly needed.

Adalimumab may cross the placenta into the serum of infants born to women treated with adalimumab during pregnancy. Consequently, these infants may be at increased risk for infection. Administration of live vaccines (e.g. BCG vaccine) to infants exposed to adalimumab *in utero* is not recommended for 5 months following the mother’s last adalimumab injection during pregnancy.

### Breast-feeding

Limited information from the published literature indicates that adalimumab is excreted in breast milk at very low concentrations with the presence of adalimumab in human milk at concentrations of 0.1% to 1% of the maternal serum level. Given orally, immunoglobulin G proteins undergo intestinal proteolysis and have poor bioavailability. No effects on the breastfed newborns/infants are anticipated. Consequently, AMGEVITA 100 mg/mL can be used during breast-feeding.

### Fertility

Preclinical data on fertility effects of adalimumab are not available.

#### **4.7 Effects on ability to drive and use machines**

AMGEVITA 100 mg/mL may have a minor influence on the ability to drive and use machines. Vertigo and visual impairment may occur following administration of AMGEVITA 100 mg/mL (see section 4.8).

#### **4.8 Undesirable effects**

##### Summary of the safety profile

Adalimumab was studied in 9,506 patients in pivotal controlled and open-label trials for up to 60 months or more. These trials included rheumatoid arthritis patients with short term and long-standing disease, juvenile idiopathic arthritis (polyarticular juvenile idiopathic arthritis and enthesitis-related arthritis) as well as axial spondyloarthritis (ankylosing spondylitis and axial spondyloarthritis without radiographic evidence of AS), psoriatic arthritis, Crohn's disease, ulcerative colitis, psoriasis, hidradenitis suppurativa and uveitis patients. The pivotal controlled studies involved 6,089 patients receiving adalimumab and 3,801 patients receiving placebo or active comparator during the controlled period.

The proportion of patients who discontinued treatment due to adverse events during the double-blind, controlled portion of pivotal studies was 5.9% for patients taking adalimumab and 5.4% for control-treated patients.

The most commonly reported adverse reactions are infections (such as nasopharyngitis, upper respiratory tract infection and sinusitis), injection site reactions (erythema, itching, hemorrhage, pain or swelling), headache and musculoskeletal pain.

Serious adverse reactions have been reported for adalimumab. TNF-antagonists, such as AMGEVITA 100 mg/mL affect the immune system and their use may affect the body's defence against infection and cancer.

Fatal and life-threatening infections (including sepsis, opportunistic infections and TB), HBV reactivation and various malignancies (including leukemia, lymphoma and HSTCL) have also been reported with use of adalimumab.

Serious hematological, neurological and autoimmune reactions have also been reported. These include rare reports of pancytopenia, aplastic anemia, central and peripheral demyelinating events and reports of lupus, lupus-related conditions and Stevens-Johnson syndrome.

##### Pediatric population

In general, the adverse events in pediatric patients were similar in frequency and type to those seen in adult patients.

##### Tabulated list of adverse reactions

The following list of adverse reactions is based on experience from clinical trials and on post-marketing experience and are displayed by system organ class and frequency in table 7 below:

very common ( $\geq 1/10$ ); common ( $\geq 1/100$  to  $< 1/10$ ); uncommon ( $\geq 1/1,000$  to  $< 1/100$ ); rare ( $\geq 1/10,000$  to  $< 1/1,000$ ); and not known (cannot be estimated from the available data). Within each frequency grouping, undesirable effects are presented in order of decreasing seriousness. The highest frequency seen among the various indications has been included. An asterisk (\*) appears in the SOC column if further information is found elsewhere in sections 4.3, 4.4 and 4.8.

**Table 7. Undesirable effects**

<b>System Organ Class</b>	<b>Frequency</b>	<b>Adverse Reaction</b>
Infections and infestations*	Very common	Respiratory tract infections (including lower and upper respiratory tract infection, pneumonia, sinusitis, pharyngitis, nasopharyngitis and pneumonia herpes viral)
	Common	Systemic infections (including sepsis, candidiasis and influenza), Intestinal infections (including gastroenteritis viral), Skin and soft tissue infections (including paronychia, cellulitis, impetigo, necrotizing fasciitis and herpes zoster), Ear infections, Oral infections (including herpes simplex, oral herpes and tooth infections), Reproductive tract infections (including vulvovaginal mycotic infection), Urinary tract infections (including pyelonephritis), Fungal infections, Joint infections
	Uncommon	Neurological infections (including viral meningitis), Opportunistic infections and tuberculosis (including coccidioidomycosis, histoplasmosis and mycobacterium avium complex infection), Bacterial infections, Eye infections, Diverticulitis <sup>1)</sup>
Neoplasms benign, malignant and unspecified (including cysts and polyps)*	Common	Skin cancer excluding melanoma (including basal cell carcinoma and squamous cell carcinoma), Benign neoplasm
	Uncommon	Lymphoma**, Solid organ neoplasm (including breast cancer, lung neoplasm and thyroid neoplasm), Melanoma**
	Rare	Leukemia <sup>1)</sup>
	Not known	Hepatosplenic T-cell lymphoma <sup>1)</sup> , Merkel cell carcinoma (neuroendocrine carcinoma of the skin) <sup>1)</sup> , Kaposi's sarcoma
Blood and the lymphatic system disorders*	Very common	Leukopenia (including neutropenia and agranulocytosis), Anemia
	Common	Leukocytosis, Thrombocytopenia
	Uncommon	Idiopathic thrombocytopenic purpura
	Rare	Pancytopenia
Immune system disorders*	Common	Hypersensitivity, Allergies (including seasonal allergy)

<b>System Organ Class</b>	<b>Frequency</b>	<b>Adverse Reaction</b>
	Uncommon	Sarcoidosis <sup>1)</sup> , Vasculitis
	Rare	Anaphylaxis <sup>1)</sup>
Metabolism and nutrition disorders	Very common	Lipids increased
	Common	Hypokalemia, Uric acid increased, Blood sodium abnormal, Hypocalcemia, Hyperglycemia, Hypophosphatemia, Dehydration
Psychiatric disorders	Common	Mood alterations (including depression), Anxiety, Insomnia
Nervous system disorders*	Very common	Headache
	Common	Paresthesias (including hypoesthesia), Migraine, Nerve root compression
	Uncommon	Cerebrovascular accident <sup>1)</sup> , Tremor, Neuropathy
	Rare	Multiple sclerosis, Demyelinating disorders (e.g. optic neuritis, Guillain-Barré syndrome) <sup>1)</sup>
Eye disorders	Common	Visual impairment, Conjunctivitis, Blepharitis, Eye swelling
	Uncommon	Diplopia
Ear and labyrinth disorders	Common	Vertigo
	Uncommon	Deafness, Tinnitus
Cardiac disorders*	Common	Tachycardia
	Uncommon	Myocardial infarction <sup>1)</sup> , Arrhythmia, Congestive heart failure
	Rare	Cardiac arrest
Vascular disorders	Common	Hypertension, Flushing, Hematoma
	Uncommon	Aortic aneurysm, Vascular arterial occlusion, Thrombophlebitis
Respiratory, thoracic and mediastinal disorders*	Common	Asthma, Dyspnea, Cough
	Uncommon	Pulmonary embolism <sup>1)</sup> , Interstitial lung disease, Chronic obstructive pulmonary disease, Pneumonitis, Pleural effusion <sup>1)</sup>

<b>System Organ Class</b>	<b>Frequency</b>	<b>Adverse Reaction</b>
	Rare	Pulmonary fibrosis <sup>1)</sup>
Gastrointestinal disorders	Very common	Abdominal pain, Nausea and vomiting
	Common	GI hemorrhage, Dyspepsia, Gastroesophageal reflux disease, Sicca syndrome
	Uncommon	Pancreatitis, Dysphagia, Face edema
	Rare	Intestinal perforation <sup>1)</sup>
Hepato-biliary disorders*	Very common	Elevated liver enzymes
	Uncommon	Cholecystitis and cholelithiasis, Hepatic steatosis, Bilirubin increased
	Rare	Hepatitis, Reactivation of hepatitis B <sup>1)</sup> , Autoimmune hepatitis <sup>1)</sup>
	Not known	Liver failure <sup>1)</sup>
Skin and subcutaneous tissue disorders	Very common	Rash (including exfoliative rash)
	Common	Worsening or new onset of psoriasis (including palmoplantar pustular psoriasis) <sup>1)</sup> , Urticaria, Bruising (including purpura), Dermatitis (including eczema), Onychoclasis, Hyperhidrosis, Alopecia <sup>1)</sup> , Pruritus
	Uncommon	Night sweats, Scar
	Rare	Erythema multiforme <sup>1)</sup> , Stevens-Johnson syndrome <sup>1)</sup> , Angioedema <sup>1)</sup> , Cutaneous vasculitis <sup>1)</sup> , Lichenoid skin reaction <sup>1)</sup>
	Not known	Worsening of symptoms of dermatomyositis <sup>1)</sup>
Musculoskeletal and connective tissue disorders	Very common	Musculoskeletal pain
	Common	Muscle spasms (including blood creatine phosphokinase increased)
	Uncommon	Rhabdomyolysis, Systemic lupus erythematosus
	Rare	Lupus-like syndrome <sup>1)</sup>
Renal and urinary disorders	Common	Renal impairment, Hematuria
	Uncommon	Nocturia
Reproductive system and breast disorders	Uncommon	Erectile dysfunction
General disorders and	Very common	Injection site reaction (including injection site erythema)

<b>System Organ Class</b>	<b>Frequency</b>	<b>Adverse Reaction</b>
administration site conditions*	Common	Chest pain, Edema, Pyrexia <sup>1)</sup>
	Uncommon	Inflammation
Investigations*	Common	Coagulation and bleeding disorders (including activated partial thromboplastin time prolonged), Autoantibody test positive (including double- stranded DNA antibody), Blood lactate dehydrogenase increased
	Not known	Weight increased <sup>2)</sup>
Injury, poisoning and procedural complications	Common	Impaired healing

\* further information is found elsewhere in sections 4.3, 4.4 and 4.8

\*\* including open-label extension studies

<sup>1)</sup> including spontaneous reporting data

<sup>2)</sup> The mean weight change from baseline for adalimumab ranged from 0.3 kg to 1.0 kg across adult indications compared to (minus) -0.4 kg to 0.4 kg for placebo over a treatment period of 4-6 months. Weight increase of 5-6 kg has also been observed in long-term extension studies with mean exposures of approximately 1-2 years without control group, particularly in patients with Crohn's disease and Ulcerative colitis. The mechanism behind this effect is unclear but could be associated with the anti-inflammatory effect of adalimumab.

#### Hidradenitis suppurativa

The safety profile for patients with HS treated with adalimumab weekly was consistent with the known safety profile of adalimumab.

#### Uveitis

The safety profile for patients with uveitis treated with adalimumab every other week was consistent with the known safety profile of adalimumab.

#### Description of selected adverse reactions

##### *Injection site reactions*

In the pivotal controlled trials in adults and children, 12.9% of patients treated with adalimumab developed injection site reactions (erythema and/or itching, hemorrhage, pain or swelling), compared to 7.2% of patients receiving placebo or active control. Injection site reactions generally did not necessitate discontinuation of the medicinal product.

##### *Infections*

In the pivotal controlled trials in adults and children, the rate of infection was 1.51 per patient year in the adalimumab-treated patients and 1.46 per patient year in the placebo and active control-treated patients. The infections consisted primarily of nasopharyngitis, upper respiratory tract infection, and sinusitis. Most patients continued on adalimumab after the infection resolved.

The incidence of serious infections was 0.04 per patient year in adalimumab-treated patients and 0.03 per patient year in placebo and active control-treated patients.

In controlled and open-label adult and pediatric studies with adalimumab, serious infections (including fatal infections, which occurred rarely) have been reported, which include reports of tuberculosis (including miliary and extra-pulmonary locations) and invasive opportunistic infections (e.g. disseminated or extra-pulmonary histoplasmosis, blastomycosis, coccidioidomycosis, pneumocystis,

candidiasis, aspergillosis and listeriosis). Most of the cases of tuberculosis occurred within the first eight months after initiation of therapy and may reflect recrudescence of latent disease.

#### *Malignancies and lymphoproliferative disorders*

No malignancies were observed in 249 pediatric patients with an exposure of 655.6 patient-years during adalimumab trials in patients with juvenile idiopathic arthritis (polyarticular juvenile idiopathic arthritis and enthesitis-related arthritis). In addition, no malignancies were observed in 192 pediatric patients with an exposure of 498.1 patient-years during adalimumab trials in pediatric patients with Crohn's disease. No malignancies were observed in 77 pediatric patients with an exposure of 80.0 patient-years during a adalimumab trial in pediatric patients with chronic plaque psoriasis. No malignancies were observed in 93 pediatric patients with an exposure of 65.3 patient-years during a adalimumab trial in pediatric patients with ulcerative colitis. No malignancies were observed in 60 pediatric patients with an exposure of 58.4 patient-years during a adalimumab trial in pediatric patients with uveitis.

During the controlled portions of pivotal adalimumab trials in adults of at least 12 weeks in duration in patients with moderately to severely active rheumatoid arthritis, ankylosing spondylitis, axial spondyloarthritis without radiographic evidence of AS, psoriatic arthritis, psoriasis, hidradenitis suppurativa, Crohn's disease, ulcerative colitis and uveitis, malignancies, other than lymphoma and non-melanoma skin cancer, were observed at a rate (95% confidence interval) of 6.8 (4.4, 10.5) per 1,000 patient-years among 5,291 adalimumab-treated patients versus a rate of 6.3 (3.4, 11.8) per 1,000 patient-years among 3,444 control patients (median duration of treatment was 4.0 months for adalimumab and 3.8 months for control-treated patients).

The rate (95% confidence interval) of non-melanoma skin cancers was 8.8 (6.0, 13.0) per 1,000 patient-years among adalimumab-treated patients and 3.2 (1.3, 7.6) per 1,000 patient-years among control patients. Of these skin cancers, squamous cell carcinomas occurred at rates (95% confidence interval) of 2.7 (1.4, 5.4) per 1,000 patient-years among adalimumab-treated patients and 0.6 (0.1, 4.5) per 1,000 patient-years among control patients. The rate (95% confidence interval) of lymphomas was 0.7 (0.2, 2.7) per 1,000 patient-years among adalimumab-treated patients and 0.6 (0.1, 4.5) per 1,000 patient-years among control patients.

When combining controlled portions of these trials and ongoing and completed open-label extension studies with a median duration of approximately 3.3 years including 6,427 patients and over 26,439 patient-years of therapy, the observed rate of malignancies, other than lymphoma and non-melanoma skin cancers is approximately 8.5 per 1,000 patient-years. The observed rate of non-melanoma skin cancers is approximately 9.6 per 1,000 patient-years, and the observed rate of lymphomas is approximately 1.3 per 1,000 patient-years.

In post-marketing experience from January 2003 to December 2010, predominantly in patients with rheumatoid arthritis, the spontaneously reported rate of malignancies is approximately 2.7 per 1,000 patient treatment years. The spontaneously reported rates for non-melanoma skin cancers and lymphomas are approximately 0.2 and 0.3 per 1,000 patient treatment years, respectively (see section 4.4).

Rare post-marketing cases of hepatosplenic T-cell lymphoma have been reported in patients treated with adalimumab (see section 4.4).

#### *Autoantibodies*

Patients had serum samples tested for autoantibodies at multiple time points in rheumatoid arthritis studies I–V. In these trials, 11.9% of patients treated with adalimumab and 8.1% of placebo and active control-treated patients that had negative baseline anti-nuclear antibody titres reported positive titres at week 24. Two patients out of 3,441 treated with adalimumab in all rheumatoid arthritis and psoriatic arthritis studies developed clinical signs suggestive of new onset lupus-like syndrome. The patients

improved following discontinuation of therapy. No patients developed lupus nephritis or central nervous system symptoms.

#### *Hepato-biliary events*

In controlled phase 3 trials of adalimumab in patients with rheumatoid arthritis and psoriatic arthritis with a control period duration ranging from 4 to 104 weeks, ALT elevations  $\geq 3 \times \text{ULN}$  occurred in 3.7% of adalimumab-treated patients and 1.6% of control-treated patients.

In controlled Phase 3 trials of adalimumab in patients with polyarticular juvenile idiopathic arthritis who were 4 to 17 years and enthesitis-related arthritis who were 6 to 17 years, ALT elevations  $\geq 3 \times \text{ULN}$  occurred in 6.1% of adalimumab-treated patients and 1.3% of control-treated patients. Most ALT elevations occurred with concomitant methotrexate use. No ALT elevations  $\geq 3 \times \text{ULN}$  occurred in the Phase 3 trial of adalimumab in patients with polyarticular juvenile idiopathic arthritis who were 2 to < 4 years.

In controlled phase 3 trials of adalimumab in patients with Crohn's disease and ulcerative colitis with a control period ranging from 4 to 52 weeks. ALT elevations  $\geq 3 \times \text{ULN}$  occurred in 0.9% of adalimumab-treated patients and 0.9% of controlled-treated patients.

In the Phase 3 trial of adalimumab in patients with pediatric Crohn's disease which evaluated efficacy and safety of two body weight adjusted maintenance dose regimens following body weight adjusted induction therapy up to 52 weeks of treatment, ALT elevations  $\geq 3 \times \text{ULN}$  occurred in 2.6% (5/192) of patients, of whom 4 were receiving concomitant immunosuppressants at baseline.

In controlled phase 3 trials of adalimumab in patients with plaque psoriasis with a control period duration ranging from 12 to 24 weeks, ALT elevations  $\geq 3 \times \text{ULN}$  occurred in 1.8% of adalimumab-treated patients and 1.8% of control-treated patients.

No ALT elevations  $\geq 3 \times \text{ULN}$  occurred in the Phase 3 trial of adalimumab in pediatric patients with plaque psoriasis.

In controlled trials of adalimumab (initial doses of 160 mg at week 0 and 80 mg at week 2, followed by 40 mg every week starting at week 4), in patients with hidradenitis suppurativa with a control period duration ranging from 12 to 16 weeks, ALT elevations  $\geq 3 \times \text{ULN}$  occurred in 0.3% of adalimumab-treated patients and 0.6% of control-treated patients.

In controlled trials of adalimumab (initial doses of 80 mg at week 0 followed by 40 mg every other week starting at week 1) in adult patients with uveitis up to 80 weeks with a median exposure of 166.5 days and 105.0 days in adalimumab-treated and control-treated patients, respectively, ALT elevations  $\geq 3 \times \text{ULN}$  occurred in 2.4% of adalimumab-treated patients and 2.4% of control-treated patients.

In the controlled Phase 3 trial of adalimumab in patients with pediatric ulcerative colitis (N = 93) which evaluated efficacy and safety of a maintenance dose of 0.6 mg/kg (maximum of 40 mg) every other week (N = 31) and a maintenance dose of 0.6 mg/kg (maximum of 40 mg) every Week (N = 32), following body weight adjusted induction dosing of 2.4 mg/kg (maximum of 160 mg) at Week 0 and Week 1, and 1.2 mg/kg (maximum of 80 mg) at Week 2 (N = 63), or an induction dose of 2.4 mg/kg (maximum of 160 mg) at Week 0, placebo at Week 1, and 1.2 mg/kg (maximum of 80 mg) at Week 2 (N = 30), ALT elevations  $\geq 3 \times \text{ULN}$  occurred in 1.1% (1/93) of patients.

Across all indications in clinical trials patients with raised ALT were asymptomatic and in most cases elevations were transient and resolved on continued treatment. However, there have also been post-marketing reports of liver failure as well as less severe liver disorders that may precede liver failure, such as hepatitis including autoimmune hepatitis in patients receiving adalimumab.

### Concurrent treatment with azathioprine/6-mercaptopurine

In adult Crohn's disease studies, higher incidences of malignant and serious infection-related adverse events were seen with the combination of adalimumab and azathioprine/6-mercaptopurine compared with adalimumab alone.

### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorization of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Any suspected adverse events should be reported to the Ministry of Health according to the National Regulation by using an online form <https://sideeffects.health.gov.il/>

## **4.9 Overdose**

No dose-limiting toxicity was observed during clinical trials. The highest dose level evaluated has been multiple intravenous doses of 10 mg/kg, which is approximately 15 times the recommended dose.

## **5. PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: Immunosuppressants, Tumor Necrosis Factor alpha (TNF $\alpha$ ) inhibitors.  
ATC code: L04AB04

#### Mechanism of action

Adalimumab binds specifically to TNF and neutralizes the biological function of TNF by blocking its interaction with the p55 and p75 cell surface TNF receptors.

Adalimumab also modulates biological responses that are induced or regulated by TNF, including changes in the levels of adhesion molecules responsible for leukocyte migration (ELAM-1, VCAM-1, and ICAM-1 with an IC<sub>50</sub> of 0.1-0.2 nM).

#### Pharmacodynamic effects

After treatment with adalimumab, a rapid decrease in levels of acute phase reactants of inflammation (C-reactive protein (CRP) and erythrocyte sedimentation rate (ESR)) and serum cytokines (IL-6) was observed, compared to baseline in patients with rheumatoid arthritis. Serum levels of matrix metalloproteinases (MMP-1 and MMP-3) that produce tissue remodeling responsible for cartilage destruction were also decreased after adalimumab administration. Patients treated with adalimumab usually experienced improvement in hematological signs of chronic inflammation.

A rapid decrease in CRP levels was also observed in patients with polyarticular juvenile idiopathic arthritis, Crohn's disease, ulcerative colitis and hidradenitis suppurativa after treatment with adalimumab. In patients with Crohn's disease, a reduction of the number of cells expressing inflammatory markers in the colon including a significant reduction of expression of TNF $\alpha$  was seen. Endoscopic studies in intestinal mucosa have shown evidence of mucosal healing in adalimumab-treated patients.

#### Clinical efficacy and safety

##### *Rheumatoid arthritis*

Adalimumab was evaluated in over 3,000 patients in all rheumatoid arthritis clinical trials. The efficacy and safety of adalimumab were assessed in five randomized, double-blind and well-controlled

studies. Some patients were treated for up to 120 months duration. Injection site pain of adalimumab was assessed in two randomized, active control, single-blind, two-period crossover studies.

RA study I evaluated 271 patients with moderately to severely active rheumatoid arthritis who were  $\geq 18$  years old, had failed therapy with at least one disease-modifying, anti-rheumatic drug and had insufficient efficacy with methotrexate at doses of 12.5 to 25 mg (10 mg if methotrexate-intolerant) every week and whose methotrexate dose remained constant at 10 to 25 mg every week. Doses of 20, 40 or 80 mg of adalimumab or placebo were given every other week for 24 weeks.

RA study II evaluated 544 patients with moderately to severely active rheumatoid arthritis who were  $\geq 18$  years old and had failed therapy with at least one disease-modifying, anti-rheumatic drugs. Doses of 20 or 40 mg of adalimumab were given by subcutaneous injection every other week with placebo on alternative weeks or every week for 26 weeks; placebo was given every week for the same duration. No other disease-modifying anti-rheumatic drugs were allowed.

RA study III evaluated 619 patients with moderately to severely active rheumatoid arthritis who were  $\geq 18$  years old, and who had an ineffective response to methotrexate at doses of 12.5 to 25 mg or have been intolerant to 10 mg of methotrexate every week. There were three groups in this study. The first received placebo injections every week for 52 weeks. The second received 20 mg of adalimumab every week for 52 weeks. The third group received 40 mg of adalimumab every other week with placebo injections on alternate weeks. Upon completion of the first 52 weeks, 457 patients enrolled in an open-label extension phase in which 40 mg of adalimumab/MTX was administered every other week up to 10 years.

RA study IV primarily assessed safety in 636 patients with moderately to severely active rheumatoid arthritis who were  $\geq 18$  years old. Patients were permitted to be either disease-modifying, anti-rheumatic drug-naïve or to remain on their pre-existing rheumatologic therapy provided that therapy was stable for a minimum of 28 days. These therapies include methotrexate, leflunomide, hydroxychloroquine, sulfasalazine and/or gold salts. Patients were randomized to 40 mg of adalimumab or placebo every other week for 24 weeks.

RA study V evaluated 799 methotrexate-naïve, adult patients with moderate to severely active early rheumatoid arthritis (mean disease duration less than 9 months). This study evaluated the efficacy of adalimumab 40 mg every other week/methotrexate combination therapy, adalimumab 40 mg every other week monotherapy and methotrexate monotherapy in reducing the signs and symptoms and rate of progression of joint damage in rheumatoid arthritis for 104 weeks. Upon completion of the first 104 weeks, 497 patients enrolled in an open-label extension phase in which 40 mg of adalimumab was administered every other week up to 10 years.

RA studies VI and VII each evaluated 60 patients with moderately to severely active rheumatoid arthritis who were  $\geq 18$  years old. Enrolled patients were either current users of adalimumab 40 mg/0.8 mL and rated their average injection site pain as at least 3 cm (on a 0-10 cm VAS) or were biologic-naïve subjects who were starting adalimumab 40 mg/0.8 mL. Patients were randomized to receive a single dose of adalimumab 40 mg/0.8 mL or adalimumab 40 mg/0.4 mL, followed by a single injection of the opposite treatment at their next dose.

The primary end point in RA studies I, II and III and the secondary endpoint in RA study IV was the percent of patients who achieved an ACR 20 response at week 24 or 26. The primary endpoint in RA study V was the percent of patients who achieved an ACR 50 response at week 52. RA studies III and V had an additional primary endpoint at 52 weeks of retardation of disease progression (as detected by X-ray results). RA study III also had a primary endpoint of changes in quality of life. The primary endpoint in RA studies VI and VII was injection site pain immediately after injection as measured by a 0-10 cm VAS.

## ACR response

The percent of adalimumab-treated patients achieving ACR 20, 50 and 70 responses was consistent across RA studies I, II and III. The results for the 40 mg every other week dose are summarized in table 8.

**Table 8. ACR responses in placebo-controlled trials (percent of patients)**

Response	RA study I <sup>a**</sup>		RA study II <sup>a**</sup>		RA study III <sup>a**</sup>	
	Placebo/ MTX <sup>c</sup> n = 60	Adalimuma b <sup>b</sup> / MTX <sup>c</sup> n = 63	Placebo n = 110	Adalimumab <sup>b</sup> n = 113	Placebo/ MTX <sup>c</sup> n = 200	Adalimuma b <sup>b</sup> / MTX <sup>c</sup> n = 207
ACR 20						
6 months	13.3%	65.1%	19.1%	46.0%	29.5%	63.3%
12 months	NA	NA	NA	NA	24.0%	58.9%
ACR 50						
6 months	6.7%	52.4%	8.2%	22.1%	9.5%	39.1%
12 months	NA	NA	NA	NA	9.5%	41.5%
ACR 70						
6 months	3.3%	23.8%	1.8%	12.4%	2.5%	20.8%
12 months	NA	NA	NA	NA	4.5%	23.2%

<sup>a</sup> RA study I at 24 weeks, RA study II at 26 weeks, and RA study III at 24 and 52 weeks

<sup>b</sup> 40 mg adalimumab administered every other week

<sup>c</sup> MTX = methotrexate

\*\* p < 0.01, adalimumab versus placebo

In RA studies I-IV, all individual components of the ACR response criteria (number of tender and swollen joints, physician and patient assessment of disease activity and pain, disability index (HAQ) scores and CRP (mg/dL) values) improved at 24 or 26 weeks compared to placebo. In RA study III, these improvements were maintained throughout 52 weeks.

In the open-label extension for RA study III, most patients who were ACR responders maintained response when followed for up to 10 years. Of 207 patients who were randomized to adalimumab 40 mg every other week, 114 patients continued on adalimumab 40 mg every other week for 5 years. Among those, 86 patients (75.4%) had ACR 20 responses; 72 patients (63.2%) had ACR 50 responses; and 41 patients (36%) had ACR 70 responses. Of 207 patients, 81 patients continued on adalimumab 40 mg every other week for 10 years. Among those, 64 patients (79.0%) had ACR 20 responses; 56 patients (69.1%) had ACR 50 responses; and 43 patients (53.1%) had ACR 70 responses.

In RA study IV, the ACR 20 response of patients treated with adalimumab plus standard of care was statistically significantly better than patients treated with placebo plus standard of care (p < 0.001).

In RA studies I-IV, adalimumab-treated patients achieved statistically significant ACR 20 and 50 responses compared to placebo as early as one to two weeks after initiation of treatment.

In RA study V with early rheumatoid arthritis patients who were methotrexate-naïve, combination therapy with adalimumab and methotrexate led to faster and significantly greater ACR responses than methotrexate monotherapy and adalimumab monotherapy at week 52 and responses were sustained at week 104 (see table 9).

**Table 9. ACR responses in RA study V (percent of patients)**

Response	MTX n = 257	Adalimumab n = 274	Adalimumab /MTX n = 268	p-value <sup>a</sup>	p-value <sup>b</sup>	p-value <sup>c</sup>
ACR 20						

<b>Response</b>	<b>MTX n = 257</b>	<b>Adalimumab n = 274</b>	<b>Adalimumab /MTX n = 268</b>	<b>p-value<sup>a</sup></b>	<b>p-value<sup>b</sup></b>	<b>p-value<sup>c</sup></b>
Week 52	62.6%	54.4%	72.8%	0.013	< 0.001	0.043
Week 104	56.0%	49.3%	69.4%	0.002	< 0.001	0.140
<b>ACR 50</b>						
Week 52	45.9%	41.2%	61.6%	< 0.001	< 0.001	0.317
Week 104	42.8%	36.9%	59.0%	< 0.001	< 0.001	0.162
<b>ACR 70</b>						
Week 52	27.2%	25.9%	45.5%	< 0.001	< 0.001	0.656
Week 104	28.4%	28.1%	46.6%	< 0.001	< 0.001	0.864

<sup>a</sup> p-value is from the pairwise comparison of methotrexate monotherapy and adalimumab/methotrexate combination therapy using the Mann-Whitney U test

<sup>b</sup> p-value is from the pairwise comparison of adalimumab monotherapy and adalimumab/methotrexate combination therapy using the Mann-Whitney U test

<sup>c</sup> p-value is from the pairwise comparison of adalimumab monotherapy and methotrexate monotherapy using the Mann-Whitney U test

In the open-label extension for RA study V, ACR response rates were maintained when followed for up to 10 years. Of 542 patients who were randomized to adalimumab 40 mg every other week, 170 patients continued on adalimumab 40 mg every other week for 10 years. Among those, 154 patients (90.6%) had ACR 20 responses; 127 patients (74.7%) had ACR 50 responses; and 102 patients (60.0%) had ACR 70 responses.

At week 52, 42.9% of patients who received adalimumab/methotrexate combination therapy achieved clinical remission (DAS28 (CRP) < 2.6) compared to 20.6% of patients receiving methotrexate monotherapy and 23.4% of patients receiving adalimumab monotherapy. Adalimumab/methotrexate combination therapy was clinically and statistically superior to methotrexate ( $p < 0.001$ ) and adalimumab monotherapy ( $p < 0.001$ ) in achieving a low disease state in patients with recently diagnosed moderate to severe rheumatoid arthritis. The response for the two monotherapy arms was similar ( $p = 0.447$ ). Of 342 subjects originally randomized to adalimumab monotherapy or adalimumab/methotrexate combination therapy who entered the open-label extension study, 171 subjects completed 10 years of adalimumab treatment. Among those, 109 subjects (63.7%) were reported to be in remission at 10 years.

### Radiographic response

In RA study III, where adalimumab-treated patients had a mean duration of rheumatoid arthritis of approximately 11 years, structural joint damage was assessed radiographically and expressed as change in modified Total Sharp Score (TSS) and its components, the erosion score and joint space narrowing score. Adalimumab/methotrexate patients demonstrated significantly less radiographic progression than patients receiving methotrexate alone at 6 and 12 months (see table 10).

In the open-label extension of RA study III, the reduction in rate of progression of structural damage is maintained for 8 and 10 years in a subset of patients. At 8 years, 81 of 207 patients originally treated with 40 mg adalimumab every other week were evaluated radiographically. Among those, 48 patients showed no progression of structural damage defined by a change from baseline in the mTSS of 0.5 or less. At 10 years, 79 of 207 patients originally treated with 40 mg adalimumab every other week were evaluated radiographically. Among those, 40 patients showed no progression of structural damage defined by a change from baseline in the mTSS of 0.5 or less.

**Table 10. Radiographic mean changes over 12 months in RA study III**

	Placebo/ MTX <sup>a</sup>	Adalimumab /MTX 40 mg every other week	Placebo/MTX- adalimumab /MTX (95% Confidence Interval <sup>b</sup> )	p-value
Total Sharp Score	2.7	0.1	2.6 (1.4, 3.8)	< 0.001 <sup>c</sup>
Erosion score	1.6	0.0	1.6 (0.9, 2.2)	< 0.001
JSN <sup>d</sup> score	1.0	0.1	0.9 (0.3, 1.4)	0.002

<sup>a</sup> methotrexate

<sup>b</sup> 95% confidence intervals for the differences in change scores between methotrexate and adalimumab

<sup>c</sup> Based on rank analysis

<sup>d</sup> Joint Space Narrowing

In RA study V, structural joint damage was assessed radiographically and expressed as change in modified Total Sharp Score (see table 11).

**Table 11. Radiographic mean changes at week 52 in RA study V**

	MTX n = 257 (95% Confidence Interval)	Adalimumab n = 274 (95% Confidence Interval)	Adalimumab/MTX n = 268 (95% Confidence Interval)	p-value <sup>a</sup>	p-value <sup>b</sup>	p-value <sup>c</sup>
Total sharp score	5.7 (4.2-7.3)	3.0 (1.7-4.3)	1.3 (0.5-2.1)	< 0.001	0.0020	< 0.001
Erosion score	3.7 (2.7-4.7)	1.7 (1.0-2.4)	0.8 (0.4-1.2)	< 0.001	0.0082	< 0.001
JSN score	2.0 (1.2-2.8)	1.3 (0.5-2.1)	0.5 (0-1.0)	< 0.001	0.0037	0.151

<sup>a</sup> p-value is from the pairwise comparison of methotrexate monotherapy and adalimumab/methotrexate combination therapy using the Mann-Whitney U test

<sup>b</sup> p-value is from the pairwise comparison of adalimumab monotherapy and adalimumab/methotrexate combination therapy using the Mann-Whitney U test

<sup>c</sup> p-value is from the pairwise comparison of adalimumab monotherapy and methotrexate monotherapy using the Mann-Whitney U test

Following 52 weeks and 104 weeks of treatment, the percentage of patients without progression (change from baseline in modified Total Sharp Score  $\leq$  0.5) was significantly higher with adalimumab/methotrexate combination therapy (63.8% and 61.2% respectively) compared to methotrexate monotherapy (37.4% and 33.5% respectively,  $p < 0.001$ ) and adalimumab monotherapy (50.7%,  $p < 0.002$  and 44.5%,  $p < 0.001$  respectively).

In the open-label extension of RA study V, the mean change from baseline at year 10 in the modified Total Sharp Score was 10.8, 9.2 and 3.9 in patients originally randomized to methotrexate monotherapy, adalimumab monotherapy and adalimumab/methotrexate combination therapy, respectively. The corresponding proportions of patients with no radiographic progression were 31.3%, 23.7% and 36.7% respectively.

#### Quality of life and physical function

Health-related quality of life and physical function were assessed using the disability index of the Health Assessment Questionnaire (HAQ) in the four original adequate and well-controlled trials, which was a pre-specified primary endpoint at week 52 in RA study III. All doses/schedules of adalimumab in all four studies showed statistically significantly greater improvement in the disability index of the HAQ from baseline to month 6 compared to placebo and in RA study III the same was seen at week 52. Results from the Short Form Health Survey (SF-36) for all doses/schedules of

adalimumab in all four studies support these findings, with statistically significant physical component summary (PCS) scores, as well as statistically significant pain and vitality domain scores for the 40 mg every other week dose. A statistically significant decrease in fatigue as measured by functional assessment of chronic illness therapy (FACIT) scores was seen in all three studies in which it was assessed (RA studies I, III, IV).

In RA study III, most subjects who achieved improvement in physical function and continued treatment maintained improvement through week 520 (120 months) of open-label treatment. Improvement in quality of life was measured up to week 156 (36 months) and improvement was maintained through that time.

In RA study V, the improvement in the HAQ disability index and the physical component of the SF-36 showed greater improvement ( $p < 0.001$ ) for adalimumab/methotrexate combination therapy versus methotrexate monotherapy and adalimumab monotherapy at week 52, which was maintained through week 104. Among the 250 subjects who completed the open-label extension study, improvements in physical function were maintained through 10 years of treatment.

#### Injection site pain

For the pooled crossover RA studies VI and VII, a statistically significant difference for injection site pain immediately after dosing was observed between adalimumab 40 mg/0.8 mL and adalimumab 40 mg/0.4 mL (mean VAS of 3.7 cm versus 1.2 cm, scale of 0-10 cm,  $P < 0.001$ ). This represented an 84% median reduction in injection site pain.

#### *Axial spondyloarthritis*

##### Ankylosing spondylitis (AS)

Adalimumab 40 mg every other week was assessed in 393 patients in two randomized, 24 week double-blind, placebo-controlled studies in patients with active ankylosing spondylitis (mean baseline score of disease activity [Bath Ankylosing Spondylitis Disease Activity Index (BASDAI)] was 6.3 in all groups) who have had an inadequate response to conventional therapy. Seventy-nine (20.1%) patients were treated concomitantly with disease-modifying anti-rheumatic drugs, and 37 (9.4%) patients with glucocorticoids. The blinded period was followed by an open-label period during which patients received adalimumab 40 mg every other week subcutaneously for up to an additional 28 weeks. Subjects ( $n = 215$ , 54.7%) who failed to achieve ASAS 20 at weeks 12, or 16 or 20 received early escape open-label adalimumab 40 mg every other week subcutaneously and were subsequently treated as non-responders in the double-blind statistical analyzes.

In the larger AS study I with 315 patients, results showed statistically significant improvement of the signs and symptoms of ankylosing spondylitis in patients treated with adalimumab compared to placebo. Significant response was first observed at week 2 and maintained through 24 weeks (Table 12).

**Table 12. Efficacy responses in placebo-controlled AS study – study I reduction of signs and symptoms**

<b>Response</b>	<b>Placebo N = 107</b>	<b>Adalimumab N = 208</b>
ASAS <sup>a</sup> 20		
Week 2	16%	42%***
Week 12	21%	58%***
Week 24	19%	51%***
ASAS 50		
Week 2	3%	16%***
Week 12	10%	38%***

<b>Response</b>	<b>Placebo N = 107</b>	<b>Adalimumab N = 208</b>
Week 24	11%	35%***
ASAS 70		
Week 2	0%	7%**
Week 12	5%	23%***
Week 24	8%	24%***
BASDAI <sup>b</sup> 50		
Week 2	4%	20%***
Week 12	16%	45%***
Week 24	15%	42%***

\*\*\*, \*\* Statistically significant at  $p < 0.001$ ,  $< 0.01$  for all comparisons between adalimumab and placebo at weeks 2, 12 and 24

<sup>a</sup> Assessments in Ankylosing Spondylitis

<sup>b</sup> Bath Ankylosing Spondylitis Disease Activity Index

Adalimumab-treated patients had significantly greater improvement at week 12 which was maintained through week 24 in both the SF-36 and Ankylosing Spondylitis Quality of Life Questionnaire (ASQoL).

Similar trends (not all statistically significant) were seen in the smaller randomized, double-blind, placebo controlled AS study II of 82 adult patients with active ankylosing spondylitis.

#### Axial spondyloarthritis without radiographic evidence of AS

The safety and efficacy of adalimumab were assessed in two randomized, double-blind placebo-controlled studies in patients with non-radiographic axial spondyloarthritis (nr-axSpA). Study nr-axSpA I evaluated patients with active nr-axSpA. Study nr-axSpA II was a treatment withdrawal study in active nr-axSpA patients who achieved remission during open-label treatment with adalimumab.

#### Study nr-axSpA I

In Study nr-axSpA I, adalimumab 40 mg every other week was assessed in 185 patients in a randomized, 12 week double-blind, placebo-controlled study in patients with active nr-axSpA (mean baseline score of disease activity [Bath Ankylosing Spondylitis Disease Activity Index (BASDAI)] was 6.4 for patients treated with adalimumab and 6.5 for those on placebo) who have had an inadequate response to or intolerance to  $\geq 1$  NSAIDs, or a contraindication for NSAIDs.

Thirty-three (18%) patients were treated concomitantly with disease-modifying anti-rheumatic drugs, and 146 (79%) patients with NSAIDs at baseline. The double-blind period was followed by an open-label period during which patients receive adalimumab 40 mg every other week subcutaneously for up to an additional 144 weeks. Week 12 results showed statistically significant improvement of the signs and symptoms of active nr-axSpA in patients treated with adalimumab compared to placebo (Table 13).

**Table 13. Efficacy response in placebo-controlled Study nr-axSpA I**

<b>Double-Blind Response at week 12</b>	<b>Placebo N = 94</b>	<b>Adalimumab N = 91</b>
ASAS <sup>a</sup> 40	15%	36%***
ASAS 20	31%	52%**
ASAS 5/6	6%	31%***
ASAS Partial Remission	5%	16%*
BASDAI <sup>b</sup> 50	15%	35%**
ASDAS <sup>c,d,e</sup>	-0.3	-1.0***

ASDAS Inactive Disease	4%	24%***
hs-CRP <sup>d,f,g</sup>	-0.3	-4.7***
SPARCC <sup>h</sup> MRI Sacroiliac Joints <sup>d,i</sup>	-0.6	-3.2**
SPARCC MRI Spine <sup>d,j</sup>	-0.2	-1.8**

<sup>a</sup> Assessment of Spondyloarthritis International Society

<sup>b</sup> Bath Ankylosing Spondylitis Disease Activity Index

<sup>c</sup> Ankylosing Spondylitis Disease Activity Score

<sup>d</sup> Mean change from baseline

<sup>e</sup> n = 91 placebo and n = 87 adalimumab

<sup>f</sup> High sensitivity C-Reactive Protein (mg/L)

<sup>g</sup> n = 73 placebo and n = 70 adalimumab

<sup>h</sup> Spondyloarthritis Research Consortium of Canada

<sup>i</sup> n = 84 placebo and adalimumab

<sup>j</sup> n = 82 placebo and n = 85 adalimumab

\*\*\*, \*\*, \* Statistically significant at p < 0.001, < 0.01, and < 0.05, respectively, for all comparisons between adalimumab and placebo

In the open-label extension, improvement in the signs and symptoms was maintained with adalimumab therapy through week 156.

#### Inhibition of inflammation

Significant improvement of signs of inflammation as measured by hs-CRP and MRI of both Sacroiliac Joints and the Spine was maintained in adalimumab-treated patients through week 156 and week 104, respectively.

#### Quality of life and physical function


Health-related quality of life and physical function were assessed using the HAQ-S and the SF-36 questionnaires. Adalimumab showed statistically significantly greater improvement in the HAQ-S total score and the SF-36 Physical Component Score (PCS) from baseline to week 12 compared to placebo. Improvement in health-related quality of life and physical function was maintained during the open-label extension through week 156.

#### Study nr-axSpA II

673 patients with active nr-axSpA (mean baseline disease activity [BASDAI] was 7.0) who had an inadequate response to  $\geq 2$  NSAIDs, or an intolerance to or a contraindication for NSAIDs enrolled into the open-label period of Study nr-axSpA II during which they received adalimumab 40 mg eow for 28 weeks. These patients also had objective evidence of inflammation in the sacroiliac joints or spine on MRI or elevated hs-CRP. Patients who achieved sustained remission for at least 12 weeks (N = 305) (ASDAS < 1.3 at weeks 16, 20, 24, and 28) during the open-label period were then randomized to receive either continued treatment with adalimumab 40 mg eow (N = 152) or placebo (N = 153) for an additional 40 weeks in a double-blind, placebo-controlled period (total study duration 68 weeks). Subjects who flared during the double-blind period were allowed adalimumab 40 mg eow rescue therapy for at least 12 weeks.

The primary efficacy endpoint was the proportion of patients with no flare by week 68 of the study. Flare was defined as ASDAS  $\geq 2.1$  at two consecutive visits four weeks apart. A greater proportion of patients on adalimumab had no disease flare during the double-blind period, when compared with those on placebo (70.4% vs. 47.1%, p < 0.001) (see figure 1).

**Figure 1. Kaplan-Meier curves summarizing time to flare in Study nr-axSpA II**



Note: P = Placebo (Number at Risk (flared)); A = Adalimumab (Number at Risk (flared)).

Among the 68 patients who flared in the group allocated to treatment withdrawal, 65 completed 12 weeks of rescue therapy with adalimumab, out of which 37 (56.9%) had regained remission (ASDAS < 1.3) after 12 weeks of restarting the open-label treatment.

By Week 68, patients receiving continuous adalimumab treatment showed statistically significant greater improvement of the signs and symptoms of active nr-axSpA as compared to patients allocated to treatment withdrawal during the double-blind period of the study (Table 14).

**Table 14. Efficacy response in placebo-controlled period for Study nr-axSpA II**

Double-Blind Response at Week 68	Placebo N = 153	Adalimumab N = 152
ASAS <sup>a,b</sup> 20	47.1%	70.4%***
ASAS <sup>a,b</sup> 40	45.8%	65.8%***
ASAS <sup>a</sup> Partial Remission	26.8%	42.1%**
ASDAS <sup>c</sup> Inactive Disease	33.3%	57.2%***
Partial Flare <sup>d</sup>	64.1%	40.8%***

<sup>a</sup> Assessment of SpondyloArthritis international Society

<sup>b</sup> Baseline is defined as open-label baseline when patients have active disease

<sup>c</sup> Ankylosing Spondylitis Disease Activity Score

<sup>d</sup> Partial flare is defined as ASDAS  $\geq$  1.3 but < 2.1 at 2 consecutive visits

\*\*\*, \*\* Statistically significant at  $p < 0.001$  and  $< 0.01$ , respectively, for all comparisons between adalimumab and placebo

### Psoriatic arthritis

Adalimumab, 40 mg every other week, was studied in patients with moderately to severely active psoriatic arthritis in two placebo-controlled studies, PsA studies I and II. PsA study I with 24 week duration, treated 313 adult patients who had an inadequate response to non-steroidal anti-inflammatory drug therapy and of these, approximately 50% were taking methotrexate. PsA study II with 12-week duration, treated 100 patients who had an inadequate response to DMARD therapy. Upon completion

of both studies, 383 patients enrolled in an open-label extension study, in which 40 mg adalimumab was administered every other week (eow).

There is insufficient evidence of the efficacy of adalimumab in patients with ankylosing spondylitis-like psoriatic arthropathy due to the small number of patients studied.

**Table 15. ACR response in placebo-controlled psoriatic arthritis studies (percent of patients)**

Response	PsA study I		PsA study II	
	Placebo N = 162	Adalimumab N = 151	Placebo N = 49	Adalimumab N = 51
ACR 20				
Week 12	14%	58%***	16%	39%*
Week 24	15%	57%***	N/A	N/A
ACR 50				
Week 12	4%	36%***	2%	25%***
Week 24	6%	39%***	N/A	N/A
ACR 70				
Week 12	1%	20%***	0%	14%*
Week 24	1%	23%***	N/A	N/A

\*\*\* p < 0.001 for all comparisons between adalimumab and placebo

\* p < 0.05 for all comparisons between adalimumab and placebo

N/A not applicable

ACR responses in PsA study I were similar with and without concomitant methotrexate therapy. ACR responses were maintained in the open-label extension study for up to 136 weeks.

Radiographic changes were assessed in the psoriatic arthritis studies. Radiographs of hands, wrists, and feet were obtained at baseline and week 24 during the double-blind period when patients were on adalimumab or placebo and at week 48 when all patients were on open-label adalimumab. A modified Total Sharp Score (mTSS), which included distal interphalangeal joints (i.e. not identical to the TSS used for rheumatoid arthritis), was used.

Adalimumab treatment reduced the rate of progression of peripheral joint damage compared with placebo treatment as measured by change from baseline in mTSS (mean ± SD) 0.8 ± 2.5 in the placebo group (at week 24) compared with 0.0 ± 1.9 (p < 0.001) in the adalimumab group (at week 48).

In subjects treated with adalimumab with no radiographic progression from baseline to week 48 (n = 102), 84% continued to show no radiographic progression through 144 weeks of treatment.

Adalimumab-treated patients demonstrated statistically significant improvement in physical function as assessed by HAQ and Short Form Health Survey (SF-36) compared to placebo at week 24. Improved physical function continued during the open-label extension up to week 136.

### *Psoriasis*

The safety and efficacy of adalimumab were studied in adult patients with chronic plaque psoriasis (≥ 10% BSA involvement and Psoriasis Area and Severity Index (PASI) ≥ 12 or ≥ 10) who were candidates for systemic therapy or phototherapy in randomized, double-blind studies. 73% of patients enrolled in psoriasis studies I and II had received prior systemic therapy or phototherapy. The safety and efficacy of adalimumab were also studied in adult patients with moderate to severe chronic plaque psoriasis with concomitant hand and/or foot psoriasis who were candidates for systemic therapy in a randomized double-blind study (psoriasis study III).

Psoriasis study I (REVEAL) evaluated 1,212 patients within three treatment periods. In period A, patients received placebo or adalimumab at an initial dose of 80 mg followed by 40 mg every other week starting one week after the initial dose. After 16 weeks of therapy, patients who achieved at least

a PASI 75 response (PASI score improvement of at least 75% relative to baseline), entered period B and received open-label 40 mg adalimumab every other week. Patients who maintained  $\geq$  PASI 75 response at week 33 and were originally randomized to active therapy in period A, were re-randomized in period C to receive 40 mg adalimumab every other week or placebo for an additional 19 weeks. Across all treatment groups, the mean baseline PASI score was 18.9 and the baseline Physician's Global Assessment (PGA) score ranged from "moderate" (53% of subjects included) to "severe" (41%) to "very severe" (6%).

Psoriasis study II (CHAMPION) compared the efficacy and safety of adalimumab versus methotrexate and placebo in 271 patients. Patients received placebo, an initial dose of MTX 7.5 mg and thereafter dose increases up to week 12, with a maximum dose of 25 mg or an initial dose of 80 mg adalimumab followed by 40 mg every other week (starting one week after the initial dose) for 16 weeks. There are no data available comparing adalimumab and MTX beyond 16 weeks of therapy. Patients receiving MTX who achieved a  $\geq$  PASI 50 response at week 8 and/or 12 did not receive further dose increases. Across all treatment groups, the mean baseline PASI score was 19.7 and the baseline PGA score ranged from "mild" (< 1%) to "moderate" (48%) to "severe" (46%) to "very severe" (6%).

Patients participating in all phase 2 and phase 3 psoriasis studies were eligible to enroll into an open-label extension trial, where adalimumab was given for at least an additional 108 weeks.

In psoriasis studies I and II, a primary endpoint was the proportion of patients who achieved a PASI 75 response from baseline at week 16 (see tables 16 and 17).

**Table 16. Ps study I (REVEAL) - efficacy results at 16 weeks**

	<b>Placebo</b> <b>N = 398</b> <b>n (%)</b>	<b>Adalimumab 40 mg eow</b> <b>N = 814</b> <b>n (%)</b>
$\geq$ PASI 75 <sup>a</sup>	26 (6.5)	578 (70.9) <sup>b</sup>
PASI 100	3 (0.8)	163 (20.0) <sup>b</sup>
PGA: Clear/minimal	17 (4.3)	506 (62.2) <sup>b</sup>

<sup>a</sup> Percent of patients achieving PASI 75 response was calculated as center-adjusted rate

<sup>b</sup> p < 0.001, adalimumab vs. placebo

**Table 17. Ps study II (CHAMPION) efficacy results at 16 weeks**

	<b>Placebo</b> <b>N = 53</b> <b>n (%)</b>	<b>MTX</b> <b>N = 110</b> <b>n (%)</b>	<b>Adalimumab 40 mg eow</b> <b>N = 108</b> <b>n (%)</b>
$\geq$ PASI 75	10 (18.9)	39 (35.5)	86 (79.6) <sup>a, b</sup>
PASI 100	1 (1.9)	8 (7.3)	18 (16.7) <sup>c, d</sup>
PGA: Clear/minimal	6 (11.3)	33 (30.0)	79 (73.1) <sup>a, b</sup>

<sup>a</sup> p < 0.001 adalimumab vs. placebo

<sup>b</sup> p < 0.001 adalimumab vs. methotrexate

<sup>c</sup> p < 0.01 adalimumab vs. placebo

<sup>d</sup> p < 0.05 adalimumab vs. methotrexate

In psoriasis study I, 28% of patients who were PASI 75 responders and were re-randomized to placebo at week 33 compared to 5% continuing on adalimumab, p < 0.001, experienced "loss of adequate response" (PASI score after week 33 and on or before week 52 that resulted in a < PASI 50 response relative to baseline with a minimum of a 6-point increase in PASI score relative to week 33). Of the patients who lost adequate response after re-randomization to placebo who then enrolled into the open-label extension trial, 38% (25/66) and 55% (36/66) regained PASI 75 response after 12 and 24 weeks of retreatment, respectively.

A total of 233 PASI 75 responders at week 16 and week 33 received continuous adalimumab therapy for 52 weeks in psoriasis study I, and continued adalimumab in the open-label extension trial. PASI 75 and PGA of clear or minimal response rates in these patients were 74.7% and 59.0%, respectively, after an additional 108 weeks of open-label therapy (total of 160 weeks). In an analysis in which all patients who dropped out of the study for adverse events or lack of efficacy, or who dose-escalated, were considered non-responders, PASI 75 and PGA of clear or minimal response rates in these patients were 69.6% and 55.7%, respectively, after an additional 108 weeks of open-label therapy (total of 160 weeks).

A total of 347 stable responders participated in a withdrawal and retreatment evaluation in an open-label extension study. During the withdrawal period, symptoms of psoriasis returned over time with a median time to relapse (decline to PGA “moderate” or worse) of approximately 5 months. None of these patients experienced rebound during the withdrawal period. A total of 76.5% (218/285) of patients who entered the retreatment period had a response of PGA “clear” or “minimal” after 16 weeks of retreatment, irrespective of whether they relapsed during withdrawal (69.1% [123/178] and 88.8% [95/107] for patients who relapsed and who did not relapse during the withdrawal period, respectively). A similar safety profile was observed during retreatment as before withdrawal.

Significant improvements at week 16 from baseline compared to placebo (studies I and II) and MTX (study II) were demonstrated in the DLQI (Dermatology Life Quality Index). In study I, improvements in the physical and mental component summary scores of the SF-36 were also significant compared to placebo.

In an open-label extension study, for patients who dose escalated from 40 mg every other week to 40 mg weekly due to a PASI response below 50%, 26.4% (92/349) and 37.8% (132/349) of patients achieved PASI 75 response at week 12 and 24, respectively.

Psoriasis study III (REACH) compared the efficacy and safety of adalimumab versus placebo in 72 patients with moderate to severe chronic plaque psoriasis and hand and/or foot psoriasis. Patients received an initial dose of 80 mg adalimumab followed by 40 mg every other week (starting one week after the initial dose) or placebo for 16 weeks. At week 16, a statistically significantly greater proportion of patients who received adalimumab achieved PGA of “clear” or “almost clear” for the hands and/or feet compared to patients who received placebo (30.6% versus 4.3%, respectively [ $p = 0.014$ ]).

Psoriasis study IV compared efficacy and safety of adalimumab versus placebo in 217 adult patients with moderate to severe nail psoriasis. Patients received an initial dose of 80 mg adalimumab followed by 40 mg every other week (starting one week after the initial dose) or placebo for 26 weeks followed by open-label adalimumab treatment for an additional 26 weeks. Nail psoriasis assessments included the Modified Nail Psoriasis Severity Index (mNAPSI), the Physician’s Global Assessment of Fingernail Psoriasis (PGA-F) and the Nail Psoriasis Severity Index (NAPSI) (see table 18). Adalimumab demonstrated a treatment benefit in nail psoriasis patients with different extents of skin involvement (BSA  $\geq 10\%$  (60% of patients) and BSA  $< 10\%$  and  $\geq 5\%$  (40% of patients)).

**Table 18. Ps study IV efficacy results at 16, 26 and 52 weeks**

Endpoint	Week 16 Placebo-controlled		Week 26 Placebo-controlled		Week 52 Open-label
	Placebo N = 108	Adalimumab 40 mg eow N = 109	Placebo N = 108	Adalimumab 40 mg eow N = 109	Adalimumab 40 mg eow N = 80
$\geq$ mNAPSI 75 (%)	2.9	26.0 <sup>a</sup>	3.4	46.6 <sup>a</sup>	65.0
PGA-F clear/minimal and $\geq$ 2-grade improvement (%)	2.9	29.7 <sup>a</sup>	6.9	48.9 <sup>a</sup>	61.3

Endpoint	Week 16 Placebo-controlled		Week 26 Placebo-controlled		Week 52 Open-label
	Placebo N = 108	Adalimumab 40 mg eow N = 109	Placebo N = 108	Adalimumab 40 mg eow N = 109	Adalimumab 40 mg eow N = 80
Percent change in total fingernail NAPSI (%)	-7.8	-44.2 <sup>a</sup>	-11.5	-56.2 <sup>a</sup>	-72.2

<sup>a</sup> p < 0.001, adalimumab vs. placebo

Adalimumab-treated patients showed statistically significant improvements at week 26 compared with placebo in the DLQI.

### *Hidradenitis suppurativa*

The safety and efficacy of adalimumab were assessed in randomized, double-blind, placebo-controlled studies and an open-label extension study in adult patients with moderate to severe hidradenitis suppurativa (HS) who were intolerant, had a contraindication or an inadequate response to at least a 3-month trial of systemic antibiotic therapy. The patients in HS-I and HS-II had Hurley Stage II or III disease with at least 3 abscesses or inflammatory nodules.

Study HS-I (PIONEER I) evaluated 307 patients with 2 treatment periods. In Period A, patients received placebo or adalimumab at an initial dose of 160 mg at week 0, 80 mg at week 2, and 40 mg every week starting at week 4 to week 11. Concomitant antibiotic use was not allowed during the study. After 12 weeks of therapy, patients who had received adalimumab in Period A were re-randomized in Period B to 1 of 3 treatment groups (adalimumab 40 mg every week, adalimumab 40 mg every other week, or placebo from week 12 to week 35). Patients who had been randomized to placebo in Period A were assigned to receive adalimumab 40 mg every week in Period B.

Study HS-II (PIONEER II) evaluated 326 patients with 2 treatment periods. In Period A, patients received placebo or adalimumab at an initial dose of 160 mg at week 0 and 80 mg at week 2 and 40 mg every week starting at week 4 to week 11. 19.3% of patients had continued baseline oral antibiotic therapy during the study. After 12 weeks of therapy, patients who had received adalimumab in Period A were re-randomized in Period B to 1 of 3 treatment groups (adalimumab 40 mg every week, adalimumab 40 mg every other week, or placebo from week 12 to week 35). Patients who had been randomized to placebo in Period A were assigned to receive placebo in Period B.

Patients participating in Studies HS-I and HS-II were eligible to enroll into an open-label extension study in which adalimumab 40 mg was administered every week. Mean exposure in all adalimumab population was 762 days. Throughout all 3 studies patients used topical antiseptic wash daily.

### Clinical response

Reduction of inflammatory lesions and prevention of worsening of abscesses and draining fistulas was assessed using Hidradenitis Suppurativa clinical response (HiSCR; at least a 50% reduction in total abscess and inflammatory nodule count with no increase in abscess count and no increase in draining fistula count relative to Baseline). Reduction in HS-related skin pain was assessed using a Numeric Rating Scale in patients who entered the study with an initial baseline score of 3 or greater on a 11 point scale.

At week 12, a significantly higher proportion of patients treated with adalimumab versus placebo achieved HiSCR. At week 12, a significantly higher proportion of patients in study HS-II experienced a clinically relevant decrease in HS-related skin pain (see table 19). Patients treated with adalimumab had significantly reduced risk of disease flare during the initial 12 weeks of treatment.

**Table 19. Efficacy results at 12 weeks, HS studies I and II**

	HS study I		HS study II	
	Placebo	Adalimumab 40 mg weekly	Placebo	Adalimumab 40 mg weekly
Hidradenitis Suppurativa Clinical Response (HiSCR) <sup>a</sup>	N = 154 40 (26.0%)	N = 153 64 (41.8%)*	N = 163 45 (27.6%)	N = 163 96 (58.9%)*
≥ 30% Reduction in Skin Pain <sup>b</sup>	N = 109 27 (24.8%)	N = 122 34 (27.9%)	N = 111 23 (20.7%)	N = 105 48 (45.7%)*

\*P < 0.05, \*\*P < 0.001, adalimumab versus placebo

<sup>a</sup> Among all randomized patients

<sup>b</sup> Among patients with baseline HS-related skin pain assessment ≥ 3, based on Numeric Rating Scale 0 – 10; 0 = no skin pain, 10 = skin pain as bad as you can imagine

Treatment with adalimumab 40 mg every week significantly reduced the risk of worsening of abscesses and draining fistulas. Approximately twice the proportion of patients in the placebo group in the first 12 weeks of Studies HS-I and HS-II, compared with those in the adalimumab group experienced worsening of abscesses (23.0% vs. 11.4%, respectively) and draining fistulas (30.0% vs. 13.9%, respectively).

Greater improvements at week 12 from baseline compared to placebo were demonstrated in skin-specific health-related quality of life, as measured by the Dermatology Life Quality Index (DLQI; Studies HS-I and HS-II), patient global satisfaction with medication treatment as measured by the Treatment Satisfaction Questionnaire-medication (TSQM; Studies HS-I and HS-II), and physical health as measured by the physical component summary score of the SF-36 (study HS-I).

In patients with at least a partial response to adalimumab 40 mg weekly at week 12, the HiSCR rate at week 36 was higher in patients who continued weekly adalimumab than in patients in whom dosing frequency was reduced to every other week, or in whom treatment was withdrawn (see table 20).

**Table 20. Proportion of patients<sup>a</sup> achieving HiSCR<sup>b</sup> at weeks 24 and 36 after treatment reassignment from weekly adalimumab at week 12**

	Placebo (treatment withdrawal) N = 73	Adalimumab 40 mg every other week N = 70	Adalimumab 40 mg weekly N = 70
Week 24	24 (32.9%)	36 (51.4%)	40 (57.1%)
Week 36	22 (30.1%)	28 (40.0%)	39 (55.7%)

<sup>a</sup> Patients with at least a partial response to adalimumab 40 mg weekly after 12 weeks of treatment

<sup>b</sup> Patients meeting protocol-specified criteria for loss of response or no improvement were required to discontinue from the studies and were counted as non-responders

Among patients who were at least partial responders at week 12, and who received continuous weekly adalimumab therapy, the HiSCR rate at week 48 was 68.3% and at week 96 was 65.1%. Longer term treatment with adalimumab 40 mg weekly for 96 weeks identified no new safety findings.

Among patients whose adalimumab treatment was withdrawn at week 12 in Studies HS-I and HS-II, the HiSCR rate 12 weeks after re-introduction of adalimumab 40 mg weekly returned to levels similar to that observed before withdrawal (56.0%).

### *Crohn's disease*

The safety and efficacy of adalimumab were assessed in over 1,500 patients with moderately to severely active Crohn's disease (Crohn's Disease Activity Index (CDAI) ≥ 220 and ≤ 450) in randomized, double-blind, placebo-controlled studies. Concomitant stable doses of aminosaliclates,

corticosteroids, and/or immunomodulatory agents were permitted and 80% of patients continued to receive at least one of these medications.

Induction of clinical remission (defined as CDAI < 150) was evaluated in two studies, CD study I (CLASSIC I) and CD study II (GAIN). In CD study I, 299 TNF-antagonist naïve patients were randomized to one of four treatment groups; placebo at weeks 0 and 2, 160 mg adalimumab at week 0 and 80 mg at week 2, 80 mg at week 0 and 40 mg at week 2, and 40 mg at week 0 and 20 mg at week 2. In CD study II, 325 patients who had lost response or were intolerant to infliximab were randomized to receive either 160 mg adalimumab at week 0 and 80 mg at week 2 or placebo at weeks 0 and 2. The primary non-responders were excluded from the studies and therefore these patients were not further evaluated.

Maintenance of clinical remission was evaluated in CD study III (CHARM). In CD study III, 854 patients received open-label 80 mg at week 0 and 40 mg at week 2. At week 4 patients were randomized to 40 mg every other week, 40 mg every week, or placebo with a total study duration of 56 weeks. Patients in clinical response (decrease in CDAI  $\geq$  70) at week 4 were stratified and analyzed separately from those not in clinical response at week 4. Corticosteroid taper was permitted after week 8.

CD study I and CD study II induction of remission and response rates are presented in table 21.

**Table 21. Induction of clinical remission and response (percent of patients)**

	CD study I: infliximab-naïve patients			CD study II: infliximab experienced patients	
	Placebo N = 74	Adalimumab 80/40 mg N = 75	Adalimumab 160/80 mg N = 76	Placebo N = 166	Adalimumab 160/80 mg N = 159
Week 4					
Clinical remission	12%	24%	36%*	7%	21%*
Clinical response (CR-100)	24%	37%	49%**	25%	38%**

All p-values are pairwise comparisons of proportions for adalimumab versus placebo

\* p < 0.001

\*\* p < 0.01

Similar remission rates were observed for the 160/80 mg and 80/40 mg induction regimens by week 8 and adverse events were more frequently noted in the 160/80 mg group.

In CD study III, at week 4, 58% (499/854) of patients were in clinical response and were assessed in the primary analysis. Of those in clinical response at week 4, 48% had been previously exposed to other TNF-antagonists. Maintenance of remission and response rates are presented in table 22. Clinical remission results remained relatively constant irrespective of previous TNF-antagonist exposure.

Disease-related hospitalizations and surgeries were statistically significantly reduced with adalimumab compared with placebo at week 56.

**Table 22. Maintenance of clinical remission and response (percent of patients)**

	Placebo	40 mg adalimumab every other week	40 mg adalimumab every week
Week 26	N = 170	N = 172	N = 157
Clinical remission	17%	40%*	47%*
Clinical response (CR-100)	27%	52%*	52%*
Patients in steroid-free remission for $\geq$ 90 days <sup>a</sup>	3% (2/66)	19% (11/58)**	15% (11/74)**

<b>Week 56</b>	<b>N = 170</b>	<b>N = 172</b>	<b>N = 157</b>
Clinical remission	12%	36%*	41%*
Clinical response (CR-100)	17%	41%*	48%*
Patients in steroid-free remission for $\geq 90$ days <sup>a</sup>	5% (3/66)	29% (17/58)*	20% (15/74)**

\*  $p < 0.001$  for adalimumab versus placebo pairwise comparisons of proportions

\*\*  $p < 0.02$  for adalimumab versus placebo pairwise comparisons of proportions

<sup>a</sup> Of those receiving corticosteroids at baseline

Among patients who were not in response at week 4, 43% of adalimumab maintenance patients responded by week 12 compared to 30% of placebo maintenance patients. These results suggest that some patients who have not responded by week 4 benefit from continued maintenance therapy through week 12. Therapy continued beyond 12 weeks did not result in significantly more responses (see section 4.2).

117/276 patients from CD study I and 272/777 patients from CD studies II and III were followed through at least 3 years of open-label adalimumab therapy. 88 and 189 patients, respectively, continued to be in clinical remission. Clinical response (CR-100) was maintained in 102 and 233 patients, respectively.

### Quality of life

In CD study I and CD study II, statistically significant improvement in the disease-specific inflammatory bowel disease questionnaire (IBDQ) total score was achieved at week 4 in patients randomized to adalimumab 80/40 mg and 160/80 mg compared to placebo and was seen at weeks 26 and 56 in CD study III as well among the adalimumab treatment groups compared to the placebo group.

### Ulcerative colitis

The safety and efficacy of multiple doses of adalimumab were assessed in adult patients with moderately to severely active ulcerative colitis (Mayo score 6 to 12 with endoscopy subscore of 2 to 3) in randomized, double-blind, placebo-controlled studies.

In study UC-I, 390 TNF-antagonist naïve patients were randomized to receive either placebo at weeks 0 and 2, 160 mg adalimumab at week 0 followed by 80 mg at week 2, or 80 mg adalimumab at week 0 followed by 40 mg at week 2. After week 2, patients in both adalimumab arms received 40 mg eow. Clinical remission (defined as Mayo score  $\leq 2$  with no subscore  $> 1$ ) was assessed at week 8.

In study UC-II, 248 patients received 160 mg of adalimumab at week 0, 80 mg at week 2 and 40 mg eow thereafter, and 246 patients received placebo. Clinical results were assessed for induction of remission at week 8 and for maintenance of remission at week 52.

Patients induced with 160/80 mg adalimumab achieved clinical remission versus placebo at week 8 in statistically significantly greater percentages in study UC-I (18% vs. 9% respectively,  $p = 0.031$ ) and study UC-II (17% vs. 9% respectively,  $p = 0.019$ ). In study UC-II, among those treated with adalimumab who were in remission at week 8, 21/41 (51%) were in remission at week 52.

Results from the overall UC-II study population are shown in table 23.

**Table 23. Response, remission and mucosal healing in study UC-II (percent of patients)**

	<b>Placebo</b>	<b>Adalimumab 40 mg eow</b>
<b>Week 52</b>	<b>N = 246</b>	<b>N = 248</b>
Clinical response	18%	30%*
Clinical remission	9%	17%*

Mucosal healing	15%	25%*
Steroid-free remission for $\geq 90$ days <sup>a</sup>	6% (N = 140)	13%* (N = 150)
Week 8 and 52		
Sustained response	12%	24%**
Sustained remission	4%	8%*
Sustained mucosal healing	11%	19%*

Clinical remission is Mayo score  $\leq 2$  with no subscore  $> 1$ ;

Clinical response is decrease from baseline in Mayo score  $\geq 3$  points and  $\geq 30\%$  plus a decrease in the rectal bleeding subscore [RBS]  $\geq 1$  or an absolute RBS of 0 or 1;

\*  $p < 0.05$  for adalimumab vs. placebo pairwise comparison of proportions

\*\*  $p < 0.001$  for adalimumab vs. placebo pairwise comparison of proportions

<sup>a</sup> Of those receiving corticosteroids at baseline

Of those patients who had a response at week 8, 47% were in response, 29% were in remission, 41% had mucosal healing, and 20% were in steroid-free remission for  $\geq 90$  days at week 52.

Approximately 40% of patients in study UC-II had failed prior anti-TNF treatment with infliximab. The efficacy of adalimumab in those patients was reduced compared to that in anti-TNF naïve patients. Among patients who had failed prior anti-TNF treatment, week 52 remission was achieved by 3% on placebo and 10% on adalimumab.

Patients from studies UC-I and UC-II had the option to roll over into an open-label long-term extension study (UC III). Following 3 years of adalimumab therapy, 75% (301/402) continued to be in clinical remission per partial Mayo score.

#### Hospitalization rates

During 52 weeks of studies UC-I and UC-II, lower rates of all-cause hospitalizations and UC-related hospitalizations were observed for the adalimumab-treated arm compared to the placebo arm. The number of all cause hospitalizations in the adalimumab treatment group was 0.18 per patient year vs. 0.26 per patient year in the placebo group and the corresponding figures for UC-related hospitalizations were 0.12 per patient year vs. 0.22 per patient year.

#### Quality of life

In study UC-II, treatment with adalimumab resulted in improvements in the Inflammatory Bowel Disease Questionnaire (IBDQ) score.

#### Intestinal Behcet's disease

Phase 3 Clinical study in Japan

In an open-label and uncontrolled study in 20 patients\* with intestinal Behcet's disease who have had an inadequate response to conventional therapy (steroid or immunomodulator), marked improvement rate at week 24 (the proportion of the subjects whose global assessment of gastrointestinal symptoms and endoscopic improvement are both  $\leq 1$ ) was 45.0% (9/20).

Common adverse events (at week 52) were nasopharyngitis 9 cases (45.0%), diarrhea, Behcet's syndrome (exacerbation of original disease), contused wound and cough 3 cases (15.0%) each.

\* The patients who were diagnosed to have the complete type, incomplete type or suspected according to the diagnostic criteria for Behcet’s disease by the research division of the Ministry of Health, Labor and Welfare and were observed to have a typical ulcer of 1 cm or larger in longer diameter in the ileocecal region.

### *Uveitis*

The safety and efficacy of adalimumab were assessed in adult patients with non-infectious intermediate, posterior, and panuveitis, excluding patients with isolated anterior uveitis, in two randomized, double-masked, placebo-controlled studies (UV I and II). Patients received placebo or adalimumab at an initial dose of 80 mg followed by 40 mg every other week starting one week after the initial dose. Concomitant stable doses of one non-biologic immunosuppressant were permitted.

Study UV I evaluated 217 patients with active uveitis despite treatment with corticosteroids (oral prednisone at a dose of 10 to 60 mg/day). All patients received a 2 week standardized dose of prednisone 60 mg/day at study entry followed by a mandatory taper schedule, with complete corticosteroid discontinuation by week 15.

Study UV II evaluated 226 patients with inactive uveitis requiring chronic corticosteroid treatment (oral prednisone 10 to 35 mg/day) at baseline to control their disease. Patients subsequently underwent a mandatory taper schedule, with complete corticosteroid discontinuation by week 19.

The primary efficacy endpoint in both studies was ‘time to treatment failure’. Treatment failure was defined by a multi-component outcome based on inflammatory chorioretinal and/or inflammatory retinal vascular lesions, anterior chamber (AC) cell grade, vitreous haze (VH) grade and best corrected visual acuity (BCVA).

Patients who completed Studies UV I and UV II were eligible to enroll in an uncontrolled long-term extension study with an originally planned duration of 78 weeks. Patients were allowed to continue on study medication beyond week 78 until they had access to adalimumab.

### *Clinical response*

Results from both studies demonstrated statistically significant reduction of the risk of treatment failure in patients treated with adalimumab versus patients receiving placebo (see table 24). Both studies demonstrated an early and sustained effect of adalimumab on the treatment failure rate versus placebo (see figure 2).

**Table 24. Time to treatment failure in studies UV I and UV II**

<b>Analysis Treatment</b>	<b>N</b>	<b>Failure N (%)</b>	<b>Median time to failure (months)</b>	<b>HR<sup>a</sup></b>	<b>95% CI for HR<sup>a</sup></b>	<b>p-value<sup>b</sup></b>
<b>Time to treatment failure at or after week 6 in study UV I</b>						
Primary analysis (ITT)						
Placebo	107	84 (78.5)	3.0	--	--	--
Adalimumab	110	60 (54.5)	5.6	0.50	0.36, 0.70	< 0.001
<b>Time to treatment failure at or after week 2 in study UV II</b>						
Primary analysis (ITT)						
Placebo	111	61 (55.0)	8.3	--	--	--
Adalimumab	115	45 (39.1)	NE <sup>c</sup>	0.57	0.39, 0.84	0.004


Note: Treatment failure at or after week 6 (study UV I), or at or after week 2 (study UV II), was counted as event. Drop outs due to reasons other than treatment failure were censored at the time of dropping out

<sup>a</sup> HR of adalimumab vs. placebo from proportional hazards regression with treatment as factor

<sup>b</sup> 2-sided p-value from log rank test

<sup>c</sup> NE = not estimable. Fewer than half of at risk subjects had an event

**Figure 2. Kaplan-Meier curves summarizing time to treatment failure on or after week 6 (study UV I) or week 2 (study UV II)**



Note: P# = Placebo (number of events/number at risk); A# = Adalimumab (number of events/number at risk).

In study UV I statistically significant differences in favor of adalimumab versus placebo were observed for each component of treatment failure. In study UV II, statistically significant differences were observed for visual acuity only, but the other components were numerically in favor of adalimumab.

Of the 424 subjects included in the uncontrolled long-term extension of Studies UV I and UV II, 60 subjects were regarded ineligible (e.g. due to deviations or due to complications secondary to diabetic retinopathy, due to cataract surgery or vitrectomy) and were excluded from the primary analysis of efficacy. Of the 364 remaining patients, 269 evaluable patients (74%) reached 78 weeks of open-label adalimumab treatment. Based on the observed data approach, 216 (80.3%) were in quiescence (no active inflammatory lesions, AC cell grade  $\leq 0.5+$ , VH grade  $\leq 0.5+$ ) with a concomitant steroid dose  $\leq 7.5$  mg per day, and 178 (66.2 %) were in steroid-free quiescence. BCVA was either improved or maintained ( $< 5$  letters deterioration) in 88.6% of the eyes at week 78. Data beyond week 78 were

generally consistent with these results but the number of enrolled subjects declined after this time. Overall, among the patients who discontinued the study, 18% discontinued due to adverse events, and 8% due to insufficient response to adalimumab treatment.

### Quality of life

Patient reported outcomes regarding vision-related functioning were measured in both clinical studies, using the NEI VFQ-25. Adalimumab was numerically favored for the majority of subscores with statistically significant mean differences for general vision, ocular pain, near vision, mental health, and total score in study UV I, and for general vision and mental health in study UV II. Vision related effects were not numerically in favor of adalimumab for color vision in study UV I and for color vision, peripheral vision and near vision in study UV II.

### Immunogenicity

Formation of anti-adalimumab antibodies is associated with increased clearance and reduced efficacy of adalimumab. There is no apparent correlation between the presence of anti-adalimumab antibodies and the occurrence of adverse events.

Patients in rheumatoid arthritis Studies I, II and III were tested at multiple time points for anti-adalimumab antibodies during the 6 to 12 month period. In the pivotal trials, anti-adalimumab antibodies were identified in 5.5 % (58/1053) of patients treated with adalimumab, compared to 0.5% (2/370) on placebo. In patients not given concomitant methotrexate, the incidence was 12.4%, compared to 0.6% when adalimumab was used as add-on to methotrexate.

In patients with polyarticular juvenile idiopathic arthritis who were 4 to 17 years, anti-adalimumab antibodies were identified in 15.8% (27/171) of patients treated with adalimumab. In patients not given concomitant methotrexate, the incidence was 25.6% (22/86) compared to 5.9% (5/85) when adalimumab was used as add-on to methotrexate. In patients with polyarticular juvenile idiopathic arthritis who were 2 to < 4 years old or aged 4 and above weighing < 15 kg, anti-adalimumab antibodies were identified in 7% (1/15) of patients, and the one patient was receiving concomitant methotrexate.

In patients with enthesitis-related arthritis, anti-adalimumab antibodies were identified in 10.9% (5/46) of patients treated with adalimumab. In patients not given concomitant methotrexate, the incidence was 13.6% (3/22), compared to 8.3% (2/24) when adalimumab was used as add-on to methotrexate.

In patients with psoriatic arthritis, anti-adalimumab antibodies were identified in 38/376 subjects (10%) treated with adalimumab. In patients not given concomitant methotrexate, the incidence was 13.5% (24/178 subjects), compared to 7% (14 of 198 subjects) when adalimumab was used as add-on to methotrexate.

In patients with ankylosing spondylitis anti-adalimumab antibodies were identified in 17/204 subjects (8.3%) treated with adalimumab. In patients not given concomitant methotrexate, the incidence was 16/185 (8.6%), compared to 1/19 (5.3%) when adalimumab was used as add-on to methotrexate.

In patients with non-radiographic axial spondyloarthritis, anti-adalimumab antibodies were identified in 8/152 subjects (5.3%) who were treated continuously with adalimumab.

In patients with Crohn's disease, anti-adalimumab antibodies were identified in 7/269 subjects (2.6 %) and in 19/487 subjects (3.9%) with ulcerative colitis.

In adult patients with psoriasis, anti-adalimumab antibodies were identified in 77/920 subjects (8.4%) treated with adalimumab monotherapy.

In adult plaque psoriasis patients on long-term adalimumab monotherapy who participated in a withdrawal and retreatment study, the rate of antibodies to adalimumab after retreatment (11 of 482 subjects, 2.3%) was similar to the rate observed prior to withdrawal (11 of 590 subjects, 1.9%).

In patients with pediatric psoriasis, anti-adalimumab antibodies were identified in 5/38 subjects (13%) treated with 0.8 mg/kg adalimumab monotherapy.

In patients with moderate to severe hidradenitis suppurativa, anti-adalimumab antibodies were identified in 10/99 subjects (10.1%) treated with adalimumab.

In patients with moderately to severely active pediatric Crohn's disease, the rate of anti-adalimumab antibody development in patients receiving adalimumab was 3.3%.

In adult patients with non-infectious uveitis, anti-adalimumab antibodies were identified in 4.8% (12/249) of patients treated with adalimumab.

In Japanese patients with intestinal Behcet's disease, anti-adalimumab antibodies were identified in 5% (1/20) of patients treated with adalimumab.

In patients with moderately to severely active pediatric ulcerative colitis, the rate of anti-adalimumab antibody development in patients receiving adalimumab was 3%.

Because immunogenicity analyzes are product-specific, comparison of antibody rates with those from other products is not appropriate.

### Pediatric population

#### *Juvenile idiopathic arthritis (JIA)*

#### *Polyarticular juvenile idiopathic arthritis (pJIA)*

The safety and efficacy of Adalimumab was assessed in two studies (pJIA I and II) in children with active polyarticular or polyarticular course juvenile idiopathic arthritis, who had a variety of JIA onset types (most frequently rheumatoid-factor negative or positive polyarthritis and extended oligoarthritis).

#### pJIA I

The safety and efficacy of Adalimumab were assessed in a multicenter, randomized, double-blind, parallel-group study in 171 children (4-17 years old) with polyarticular JIA. In the open-label lead in phase (OL LI) patients were stratified into two groups, MTX (methotrexate)-treated or non-MTX-treated. Patients who were in the non-MTX stratum were either naïve to or had been withdrawn from MTX at least two weeks prior to study drug administration. Patients remained on stable doses of NSAIDs and or prednisone ( $\leq 0.2$  mg/kg/day or 10 mg/day maximum). In the OL LI phase all patients received 24 mg/m<sup>2</sup> up to a maximum of 40 mg AMGEVITA 100 mg/mL every other week for 16 weeks. The distribution of patients by age and minimum, median and maximum dose received during the OL LI phase is presented in table 25.

**Table 25. Distribution of patients by age and adalimumab dose received during the OL LI phase**

Age Group	Number of patients at Baseline n (%)	Minimum, median and maximum dose
4 to 7 years	31 (18.1)	10, 20 and 25 mg
8 to 12 years	71 (41.5)	20, 25 and 40 mg
13 to 17 years	69 (40.4)	25, 40 and 40 mg

Patients demonstrating a Pediatric ACR 30 response at Week 16 were eligible to be randomized into the double blind (DB) phase and received either Adalimumab 24 mg/m<sup>2</sup> up to a maximum of 40 mg, or placebo every other week for an additional 32 weeks or until disease flare. Disease flare criteria were defined as a worsening of  $\geq 30\%$  from baseline in  $\geq 3$  of 6 Pediatric ACR core criteria,  $\geq 2$  active joints, and improvement of  $> 30\%$  in no more than 1 of the 6 criteria. After 32 weeks or at disease flare, patients were eligible to enroll into the open-label extension phase.

**Table 26. Ped ACR 30 responses in the JIA study**

Stratum	MTX		Without MTX	
Phase				
OL-LI 16 weeks				
Ped ACR 30 response (n/N)	94.1% (80/85)		74.4% (64/86)	
Efficacy Outcomes				
Double Blind 32 weeks	Adalimumab / MTX (N = 38)	Placebo / MTX (N = 37)	Adalimumab (N = 30)	Placebo (N = 28)
Disease flares at the end of 32 weeks <sup>a</sup> (n/N)	36.8% (14/38)	64.9% (24/37) <sup>b</sup>	43.3% (13/30)	71.4% (20/28) <sup>c</sup>
Median time to disease flare	> 32 weeks	20 weeks	> 32 weeks	14 weeks

<sup>a</sup> Ped ACR 30/50/70 responses Week 48 significantly greater than those of placebo treated patients

<sup>b</sup> p = 0.015

<sup>c</sup> p = 0.031

Amongst those who responded at Week 16 (n = 144), the Pediatric ACR 30/50/70/90 responses were maintained for up to six years in the OLE phase in patients who received Adalimumab throughout the study. Over all 19 subjects, of which 11 of the baseline age group 4 to 12 and 8 of the baseline age group 13 to 17 years were treated 6 years or longer.

Overall responses were generally better and, fewer patients developed antibodies when treated with the combination of Adalimumab and MTX compared to AMGEVITA 100 mg/mL alone. Taking these results into consideration, Adalimumab is recommended for use in combination with MTX and for use as monotherapy in patients for whom MTX use is not appropriate (see section 4.2).

## pJIA II

The safety and efficacy of Adalimumab was assessed in an open-label, multicenter study in 32 children (2 < 4 years old or aged 4 and above weighing < 15 kg) with moderately to severely active polyarticular JIA. The patients received 24 mg/m<sup>2</sup> body surface area (BSA) of Adalimumab up to a maximum of 20 mg every other week as a single dose via SC injection for at least 24 weeks. During the study, most subjects used concomitant MTX, with fewer reporting use of corticosteroids or NSAIDs.

At Week 12 and Week 24, PedACR30 response was 93.5% and 90.0%, respectively, using the observed data approach. The proportions of subjects with PedACR50/70/90 at Week 12 and Week 24 were 90.3%/61.3%/38.7% and 83.3%/73.3%/36.7%, respectively. Amongst those who responded (Pediatric ACR 30) at Week 24 (n = 27 out of 30 patients), the Pediatric ACR 30 responses were maintained for up to 60 weeks in the OLE phase in patients who received Adalimumab throughout this time period. Overall, 20 subjects were treated for 60 weeks or longer.

### Enthesitis-related arthritis

The safety and efficacy of Adalimumab were assessed in a multicenter, randomized,

double-blind study in 46 pediatric patients (6 to 17 years old) with moderate enthesitis-related arthritis. Patients were randomized to receive either 24 mg/m<sup>2</sup> body surface area (BSA) of Adalimumab up to a maximum of 40 mg, or placebo every other week for 12 weeks. The double-blind period is followed by an open-label (OL) period during which patients received 24 mg/m<sup>2</sup> BSA of Adalimumab up to a maximum of 40 mg every other week subcutaneously for up to an additional 192 weeks. The primary endpoint was the percent change from Baseline to Week 12 in the number of active joints with arthritis (swelling not due to deformity or joints with loss of motion plus pain and/or tenderness), which was achieved with mean percent decrease of -62.6% (median percent change -88.9%) in patients in the Adalimumab group compared to -11.6% (median percent change -50.0%) in patients in the placebo group. Improvement in number of active joints with arthritis was maintained during the OL period through Week 156 for the 26 of 31 (84%) patients in the Adalimumab group who remained in the study. Although not statistically significant, the majority of patients demonstrated clinical improvement in secondary endpoints such as number of sites of enthesitis, tender joint count (TJC), swollen joint count (SJC), Pediatric ACR 50 response, and Pediatric ACR 70 response.

#### *Pediatric plaque psoriasis*

The efficacy of Adalimumab was assessed in a randomized, double-blind, controlled study of 114 pediatric patients from 4 years of age with severe chronic plaque psoriasis (as defined by a PGA  $\geq$  4 or  $>$  20% BSA involvement or  $>$  10% BSA involvement with very thick lesions or PASI  $\geq$  20 or  $\geq$  10 with clinically relevant facial, genital, or hand/ foot involvement) who were inadequately controlled with topical therapy and heliotherapy or phototherapy.

Patients received Adalimumab 0.8 mg/kg eow (up to 40 mg), 0.4 mg/kg eow (up to 20 mg), or methotrexate 0.1 – 0.4 mg/kg weekly (up to 25 mg). At Week 16, more patients randomized to Adalimumab 0.8 mg/kg had positive efficacy responses (e.g. PASI 75) than those randomized to 0.4 mg/kg eow or MTX.

**Table 27. Pediatric Plaque Psoriasis efficacy results at 16 weeks**

	MTX <sup>a</sup> N = 37	Adalimumab 0.8 mg/kg eow N = 38
PASI 75 <sup>b</sup>	12 (32.4%)	22 (57.9%)
PGA: Clear/minimal <sup>c</sup>	15 (40.5%)	23 (60.5%)

<sup>a</sup> MTX = methotrexate

<sup>b</sup> P=0.027, Adalimumab 0.8 mg/kg versus MTX

<sup>c</sup> P=0.083, Adalimumab 0.8 mg/kg versus MTX

Patients who achieved PASI 75 and PGA clear or minimal were withdrawn from treatment for up to 36 weeks and monitored for loss of disease control (i.e. a worsening of PGA by at least 2 grades). Patients were then re-treated with adalimumab 0.8 mg/kg eow for an additional 16 weeks and response rates observed during retreatment were similar to the previous double-blind period: PASI 75 response of 78.9% (15 of 19 subjects) and PGA clear or minimal of 52.6% (10 of 19 subjects).

In the open-label period of the study, PASI 75 and PGA clear or minimal responses were maintained for up to an additional 52 weeks with no new safety findings.

### *Adolescent hidradenitis suppurativa*

There are no clinical trials with Adalimumab in adolescent patients with HS. Efficacy of adalimumab for the treatment of adolescent patients with HS is predicted based on the demonstrated efficacy and exposure-response relationship in adult HS patients and the likelihood that the disease course, pathophysiology, and drug effects are substantially similar to that of adults at the same exposure levels. Safety of the recommended adalimumab dose in the adolescent HS population is based on cross-indication safety profile of adalimumab in both adults and pediatric patients at similar or more frequent doses (see section 5.2).

### *Pediatric Crohn's disease*

Adalimumab was assessed in a multicenter, randomized, double-blind clinical trial designed to evaluate the efficacy and safety of induction and maintenance treatment with doses dependent on body weight (< 40 kg or ≥ 40 kg) in 192 pediatric subjects between the ages of 6 and 17 (inclusive) years, with moderate to severe Crohn's disease (CD) defined as Pediatric Crohn's Disease Activity Index (PCDAI) score > 30. Subjects had to have failed conventional therapy (including a corticosteroid and/or an immunomodulator) for CD. Subjects may also have previously lost response or been intolerant to infliximab.

All subjects received open-label induction therapy at a dose based on their Baseline body weight: 160 mg at Week 0 and 80 mg at Week 2 for subjects ≥ 40 kg, and 80 mg and 40 mg, respectively, for subjects < 40 kg.

At Week 4, subjects were randomized 1:1 based on their body weight at the time to either the Low Dose or Standard Dose maintenance regimens as shown in table 28.

**Table 28. Maintenance regimen**

<b>Patient Weight</b>	<b>Low dose</b>	<b>Standard dose</b>
< 40 kg	10 mg eow	20 mg eow
≥ 40 kg	20 mg eow	40 mg eow

### *Efficacy results*

The primary endpoint of the study was clinical remission at Week 26, defined as PCDAI score ≤ 10. Clinical remission and clinical response (defined as reduction in PCDAI score of at least 15 points from Baseline) rates are presented in table 29. Rates of discontinuation of corticosteroids or immunomodulators are presented in table 30.

**Table 29. Pediatric CD study PCDAI clinical remission and response**

	<b>Standard Dose 40/20 mg eow N = 93</b>	<b>Low Dose 20/10 mg eow N = 95</b>	<b>p-value*</b>
<b>Week 26</b>			
Clinical remission	38.7%	28.4%	0.075
Clinical response	59.1%	48.4%	0.073
<b>Week 52</b>			
Clinical remission	33.3%	23.2%	0.100
Clinical response	41.9%	28.4%	0.038

\* p-value for Standard Dose *versus* Low Dose comparison.

**Table 30. Pediatric CD study discontinuation of corticosteroids or immunomodulators and fistula remission**

	<b>Standard Dose 40/20 mg eow</b>	<b>Low Dose 20/10 mg eow</b>	<b>p-value<sup>1</sup></b>
<b>Discontinued corticosteroids</b>	<b>N = 33</b>	<b>N = 38</b>	
Week 26	84.8%	65.8%	0.066
Week 52	69.7%	60.5%	0.420
<b>Discontinuation of Immunomodulators<sup>2</sup></b>	<b>N = 60</b>	<b>N = 57</b>	
Week 52	30.0%	29.8%	0.983
<b>Fistula remission<sup>3</sup></b>	<b>N = 15</b>	<b>N = 21</b>	
Week 26	46.7%	38.1%	0.608
Week 52	40.0%	23.8%	0.303

<sup>1</sup> p-value for Standard Dose *versus* Low Dose comparison.

<sup>2</sup> Immunosuppressant therapy could only be discontinued at or after Week 26 at the investigator's discretion if the subject met the clinical response criterion

<sup>3</sup> defined as a closure of all fistulas that were draining at Baseline for at least 2 consecutive post-Baseline visits

Statistically significant increases (improvement) from Baseline to Week 26 and 52 in Body Mass Index and height velocity were observed for both treatment groups.

Statistically and clinically significant improvements from Baseline were also observed in both treatment groups for quality of life parameters (including IMPACT III).

One hundred patients (n = 100) from the Pediatric CD study continued in an open-label long-term extension study. After 5 years of adalimumab therapy, 74.0% (37/50) of the 50 patients remaining in the study continued to be in clinical remission, and 92.0% (46/50) of patients continued to be in clinical response per PCDAI.

#### *Pediatric ulcerative colitis*

The safety and efficacy of Adalimumab was assessed in a multicenter, randomized, double-blind, trial in 93 pediatric patients from 5 to 17 years of age with moderate to severe ulcerative colitis (Mayo score 6 to 12 with endoscopy subscore of 2 to 3 points, confirmed by centrally read endoscopy) who had an inadequate response or intolerance to conventional therapy. Approximately 16% of patients in the study had failed prior anti-TNF treatment. Patients who received corticosteroids at enrollment were allowed to taper their corticosteroid therapy after Week 4.

In the induction period of the study, 77 patients were randomized 3:2 to receive double-blind treatment with Adalimumab at an induction dose of 2.4 mg/kg (maximum of 160 mg) at Week 0 and Week 1, and 1.2 mg/kg (maximum of 80 mg) at Week 2; or an induction dose of 2.4 mg/kg (maximum of 160 mg) at Week 0, placebo at Week 1, and 1.2 mg/kg (maximum of 80 mg) at Week 2. Both groups received 0.6 mg/kg (maximum of 40 mg) at Week 4 and Week 6. Following an amendment to the study design, the remaining 16 patients who enrolled in the induction period received open-label treatment with Adalimumab at the induction dose of 2.4 mg/kg (maximum of 160 mg) at Week 0 and Week 1, and 1.2 mg/kg (maximum of 80 mg) at Week 2.

At Week 8, 62 patients who demonstrated clinical response per Partial Mayo Score (PMS; defined as a decrease in PMS  $\geq 2$  points and  $\geq 30\%$  from Baseline) were randomized equally to receive double-blind maintenance treatment with Adalimumab at a dose of 0.6 mg/kg (maximum of 40 mg) every week (ew), or a maintenance dose of 0.6 mg/kg (maximum of 40 mg) every other week (eow). Prior to an amendment to the study design, 12 additional patients who demonstrated clinical response per PMS were randomized to receive placebo but were not included in the confirmatory analysis of efficacy.

Disease flare was defined as an increase in PMS of at least 3 points (for patients with PMS of 0 to 2 at Week 8), at least 2 points (for patients with PMS of 3 to 4 at Week 8), or at least 1 point (for patients with PMS of 5 to 6 at Week 8).

Patients who met criteria for disease flare at or after Week 12 were randomized to receive a re-induction dose of 2.4 mg/kg (maximum of 160 mg) or a dose of 0.6 mg/kg (maximum of 40 mg) and continued to receive their respective maintenance dose regimen afterwards.

### *Efficacy Results*

The co-primary endpoints of the study were clinical remission per PMS (defined as PMS  $\leq 2$  and no individual subscore  $> 1$ ) at Week 8, and clinical remission per FMS (Full Mayo Score) (defined as a Mayo Score  $\leq 2$  and no individual subscore  $> 1$ ) at Week 52 in patients who achieved clinical response per PMS at Week 8.

Clinical remission rates per PMS at Week 8 for patients in each of the Adalimumab double-blind induction groups are presented in table 31.

**Table 31. Clinical remission per PMS at 8 weeks**

	<b>Adalimumab<sup>a</sup> Maximum of 160 mg at Week 0 / Placebo at Week 1</b> N = 30	<b>Adalimumab<sup>b, c</sup> Maximum of 160 mg at Week 0 and Week 1</b> N = 47
Clinical remission	13/30 (43.3%)	28/47 (59.6%)

<sup>a</sup> Adalimumab 2.4 mg/kg (maximum of 160 mg) at Week 0, placebo at Week 1, and 1.2 mg/kg (maximum of 80 mg) at Week 2

<sup>b</sup> Adalimumab 2.4 mg/kg (maximum of 160 mg) at Week 0 and Week 1, and 1.2 mg/kg (maximum of 80 mg) at Week 2

<sup>c</sup> Not including open-label Induction dose of adalimumab 2.4 mg/kg (maximum of 160 mg) at Week 0 and Week 1, and 1.2 mg/kg (maximum of 80 mg) at Week 2

Note 1: Both induction groups received 0.6 mg/kg (maximum of 40 mg) at Week 4 and Week 6

Note 2: Patients with missing values at Week 8 were considered as not having met the endpoint

At Week 52, clinical remission per FMS in Week 8 responders, clinical response per FMS (defined as a decrease in Mayo Score  $\geq 3$  points and  $\geq 30\%$  from Baseline) in Week 8 responders, mucosal healing (defined as Mayo endoscopy subscore  $\leq 1$ ) in Week 8 responders, clinical remission per FMS in Week 8 remitters, and the proportion of subjects in corticosteroid-free remission per FMS in Week 8 responders were assessed in patients who received Adalimumab at the double-blind maximum 40 mg eow (0.6 mg/kg) and maximum 40 mg ew (0.6 mg/kg) maintenance doses (Table 32).

**Table 32. Efficacy results at 52 Weeks**

	<b>Adalimumab<sup>a</sup> Maximum of 40 mg eow</b> N = 31	<b>Adalimumab<sup>b</sup> Maximum of 40 mg ew</b> N = 31
Clinical remission in Week 8 PMS responders	9/31 (29.0%)	14/31 (45.2%)
Clinical response in Week 8 PMS responders	19/31 (61.3%)	21/31 (67.7%)
Mucosal healing in Week 8 PMS responders	12/31 (38.7%)	16/31 (51.6%)
Clinical remission in Week 8 PMS remitters	9/21 (42.9%)	10/22 (45.5%)
Corticosteroid-free remission in Week 8 PMS responders <sup>c</sup>	4/13 (30.8%)	5/16 (31.3%)

<sup>a</sup> Adalimumab 0.6 mg/kg (maximum of 40 mg) every other week

<sup>b</sup> Adalimumab 0.6 mg/kg (maximum of 40 mg) every week

<sup>c</sup> In patients receiving concomitant corticosteroids at baseline

Note: Patients with missing values at Week 52 or who were randomized to receive re-induction or maintenance treatment were considered non-responders for Week 52 endpoints

Additional exploratory efficacy endpoints included clinical response per the Pediatric Ulcerative Colitis Activity Index (PUCAI) (defined as a decrease in PUCAI  $\geq$  20 points from Baseline) and clinical remission per PUCAI (defined as PUCAI < 10) at Week 8 and Week 52 (Table 33).

**Table 33. Exploratory endpoints results per PUCAI**

	<b>Week 8</b>	
	<b>Adalimumab<sup>a</sup> Maximum of 160 mg at Week 0 / Placebo at Week 1</b> N = 30	<b>Adalimumab<sup>b,c</sup> Maximum of 160 mg at Week 0 and Week 1</b> N = 47
Clinical remission per PUCAI	10/30 (33.3%)	22/47 (46.8%)
Clinical response per PUCAI	15/30 (50.0%)	32/47 (68.1%)
	<b>Week 52</b>	
	<b>Adalimumab<sup>d</sup> Maximum of 40 mg eow</b> N = 31	<b>Adalimumab<sup>e</sup> Maximum of 40 mg ew</b> N = 31
Clinical remission per PUCAI in Week 8 PMS responders	14/31 (45.2%)	18/31 (58.1%)
Clinical response per PUCAI in Week 8 PMS responders	18/31 (58.1%)	16/31 (51.6%)

<sup>a</sup> Adalimumab 2.4 mg/kg (maximum of 160 mg) at Week 0, placebo at Week 1, and 1.2 mg/kg (maximum of 80 mg) at Week 2

<sup>b</sup> Adalimumab 2.4 mg/kg (maximum of 160 mg) at Week 0 and Week 1, and 1.2 mg/kg (maximum of 80 mg) at Week 2

<sup>c</sup> Not including open-label Induction dose of adalimumab 2.4 mg/kg (maximum of 160 mg) at Week 0 and Week 1, and 1.2 mg/kg (maximum of 80 mg) at Week 2

<sup>d</sup> Adalimumab 0.6 mg/kg (maximum of 40 mg) every other week

<sup>e</sup> Adalimumab 0.6 mg/kg (maximum of 40 mg) every week

Note 1: Both induction groups received 0.6 mg/kg (maximum of 40 mg) at Week 4 and Week 6

Note 2: Patients with missing values at Week 8 were considered as not having met the endpoints

Note 3: Patients with missing values at Week 52 or who were randomized to receive re-induction or

---

maintenance treatment were considered non-responders for Week 52 endpoints

Of the Adalimumab-treated patients who received re-induction treatment during the maintenance period, 2/6 (33%) achieved clinical response per FMS at Week 52.

### *Quality of life*

Clinically meaningful improvements from Baseline were observed in IMPACT III and the caregiver Work Productivity and Activity Impairment (WPAI) scores for the groups treated with Adalimumab.

Clinically meaningful increases (improvement) from Baseline in height velocity were observed for the groups treated with adalimumab, and clinically meaningful increases (improvement) from Baseline in Body Mass Index were observed for subjects on the high maintenance dose of maximum 40 mg (0.6 mg/kg) ew.

### *Pediatric Uveitis*


The safety and efficacy of Adalimumab was assessed in a randomized, double-masked, controlled study of 90 pediatric patients from 2 to < 18 years of age with active JIA-associated non-infectious anterior uveitis who were refractory to at least 12 weeks of methotrexate treatment. Patients received either placebo or 20 mg adalimumab (if < 30 kg) or 40 mg adalimumab (if  $\geq$  30 kg) every other week in combination with their baseline dose of methotrexate.

The primary endpoint was 'time to treatment failure'. The criteria determining treatment failure were worsening or sustained non-improvement in ocular inflammation, partial improvement with development of sustained ocular co-morbidities or worsening of ocular co-morbidities, non-permitted use of concomitant medications, and suspension of treatment for an extended period of time.

### *Clinical Response*

Adalimumab significantly delayed the time to treatment failure, as compared to placebo (see figure 3,  $P < 0.0001$  from log rank test). The median time to treatment failure was 24.1 weeks for subjects treated with placebo, whereas the median time to treatment failure was not estimable for subjects treated with adalimumab because less than one-half of these subjects experienced treatment failure. Adalimumab significantly decreased the risk of treatment failure by 75% relative to placebo, as shown by the hazard ratio (HR = 0.25 [95% CI: 0.12, 0.49]).

**Figure 3. Kaplan-Meier curves summarizing time to treatment failure in the Pediatric Uveitis Study**



Note: P = Placebo (Number at Risk); H = Adalimumab (Number at Risk).

## 5.2 Pharmacokinetic properties

### Absorption and distribution

After subcutaneous administration of a single 40 mg dose, absorption and distribution of adalimumab was slow, with peak serum concentrations being reached about 5 days after administration. The average absolute bioavailability of adalimumab estimated from three studies following a single 40 mg subcutaneous dose was 64%. After single intravenous doses ranging from 0.25 to 10 mg/kg, concentrations were dose proportional. After doses of 0.5 mg/kg (~40 mg), clearances ranged from 11 to 15 mL/hour, the distribution volume ( $V_{ss}$ ) ranged from 5 to 6 liters and the mean terminal phase half-life was approximately two weeks. Adalimumab concentrations in the synovial fluid from several rheumatoid arthritis patients ranged from 31-96% of those in serum.

Following subcutaneous administration of 40 mg of adalimumab every other week in adult rheumatoid arthritis (RA) patients the mean steady-state trough concentrations were approximately 5  $\mu\text{g/mL}$  (without concomitant methotrexate) and 8 to 9  $\mu\text{g/mL}$  (with concomitant methotrexate), respectively. The serum adalimumab trough levels at steady-state increased roughly proportionally with dose following 20, 40 and 80 mg subcutaneous dosing every other week and every week.

Following the administration of 24  $\text{mg/m}^2$  (maximum of 40 mg) subcutaneously every other week to patients with polyarticular juvenile idiopathic arthritis (JIA) who were 4 to 17 years the mean trough steady-state (values measured from Week 20 to 48) serum adalimumab concentration was  $5.6 \pm 5.6 \mu\text{g/mL}$  (102% CV) for adalimumab without concomitant methotrexate and  $10.9 \pm 5.2 \mu\text{g/mL}$  (47.7% CV) with concomitant methotrexate.

In patients with polyarticular JIA who were 2 to < 4 years old or aged 4 and above weighing < 15 kg dosed with adalimumab 24  $\text{mg/m}^2$ , the mean trough steady-state serum adalimumab concentrations

was  $6.0 \pm 6.1$   $\mu\text{g/mL}$  (101% CV) for adalimumab without concomitant methotrexate and  $7.9 \pm 5.6$   $\mu\text{g/mL}$  (71.2% CV) with concomitant methotrexate.

Following the administration of  $24 \text{ mg/m}^2$  (maximum of 40 mg) subcutaneously every other week to patients with enthesitis-related arthritis who were 6 to 17 years, the mean trough steady-state (values measured at Week 24) serum adalimumab concentrations were  $8.8 \pm 6.6$   $\mu\text{g/mL}$  for adalimumab without concomitant methotrexate and  $11.8 \pm 4.3$   $\mu\text{g/mL}$  with concomitant methotrexate.

Following subcutaneous administration of 40 mg of adalimumab every other week in adult non-radiographic axial spondyloarthritis patients, the mean ( $\pm$ SD) trough steady-state concentration at Week 68 was  $8.0 \pm 4.6$   $\mu\text{g/mL}$ .

In adult patients with psoriasis, the mean steady-state trough concentration was 5  $\mu\text{g/mL}$  during adalimumab 40 mg every other week monotherapy treatment.

Following the administration of 0.8 mg/kg (maximum of 40 mg) subcutaneously every other week to pediatric patients with chronic plaque psoriasis, the mean  $\pm$  SD steady-state adalimumab trough concentration was approximately  $7.4 \pm 5.8$   $\mu\text{g/mL}$  (79% CV).

In adult patients with hidradenitis suppurativa, a dose of 160 mg adalimumab on week 0 followed by 80 mg on week 2 achieved serum adalimumab trough concentrations of approximately 7 to 8  $\mu\text{g/mL}$  at week 2 and week 4. The mean steady-state trough concentration at week 12 through week 36 were approximately 8 to 10  $\mu\text{g/mL}$  during adalimumab 40 mg every week treatment.

Adalimumab exposure in adolescent HS patients was predicted using population pharmacokinetic modelling and simulation based on cross-indication pharmacokinetics in other pediatric patients (pediatric psoriasis, juvenile idiopathic arthritis, pediatric Crohn's disease, and enthesitis-related arthritis). The recommended adolescent HS dosing schedule is 40 mg every other week. Since exposure to adalimumab can be affected by body size, adolescents with higher body weight and inadequate response may benefit from receiving the recommended adult dose of 40 mg every week.

In patients with Crohn's disease, the loading dose of 80 mg adalimumab on week 0 followed by 40 mg adalimumab on week 2 achieves serum adalimumab trough concentrations of approximately 5.5  $\mu\text{g/mL}$  during the induction period. A loading dose of 160 mg adalimumab on week 0 followed by 80 mg adalimumab on week 2 achieves serum adalimumab trough concentrations of approximately 12  $\mu\text{g/mL}$  during the induction period. Mean steady-state trough levels of approximately 7  $\mu\text{g/mL}$  were observed in Crohn's disease patients who received a maintenance dose of 40 mg adalimumab every other week.

In pediatric patients with moderate to severe CD, the open-label adalimumab induction dose was 160/80 mg or 80/40 mg at Weeks 0 and 2, respectively, dependent on a body weight cut-off of 40 kg. At Week 4, patients were randomized 1:1 to either the Standard Dose (40/20 mg eow) or Low Dose (20/10 mg eow) maintenance treatment groups based on their body weight. The mean ( $\pm$ SD) serum adalimumab trough concentrations achieved at Week 4 were  $15.7 \pm 6.6$   $\mu\text{g/mL}$  for patients  $\geq 40$  kg (160/80 mg) and  $10.6 \pm 6.1$   $\mu\text{g/mL}$  for patients  $< 40$  kg (80/40 mg).

For patients who stayed on their randomized therapy, the mean ( $\pm$ SD) adalimumab trough concentrations at week 52 were  $9.5 \pm 5.6$   $\mu\text{g/mL}$  for the standard dose group and  $3.5 \pm 2.2$   $\mu\text{g/mL}$  for the low dose group. The mean trough concentrations were maintained in patients who continued to receive adalimumab treatment eow for 52 weeks. For patients who dose escalated from eow to weekly regimen, the mean ( $\pm$ SD) serum concentrations of adalimumab at week 52 were  $15.3 \pm 11.4$   $\mu\text{g/mL}$  (40/20 mg, weekly) and  $6.7 \pm 3.5$   $\mu\text{g/mL}$  (20/10 mg, weekly).

In patients with ulcerative colitis, a loading dose of 160 mg adalimumab on week 0 followed by 80 mg adalimumab on week 2 achieves serum adalimumab trough concentrations of approximately 12  $\mu\text{g/mL}$  during the induction period. Mean steady-state trough levels of approximately 8  $\mu\text{g/mL}$  were observed in ulcerative colitis patients who received a maintenance dose of 40 mg adalimumab every other week.

Following the subcutaneous administration of body weight-based dosing of 0.6 mg/kg (maximum of 40 mg) every other week to pediatric patients with ulcerative colitis, the mean trough steady-state serum adalimumab concentration was  $5.01 \pm 3.28$   $\mu\text{g/mL}$  at Week 52. For patients who received 0.6 mg/kg (maximum of 40 mg) every week, the mean ( $\pm$ SD) trough steady-state serum adalimumab concentration was  $15.7 \pm 5.60$   $\mu\text{g/mL}$  at Week 52.

In adult patients with uveitis, a loading dose of 80 mg adalimumab on week 0 followed by 40 mg adalimumab every other week starting at week 1, resulted in mean steady-state concentrations of approximately 8 to 10  $\mu\text{g/mL}$ .

Adalimumab exposure in pediatric uveitis patients was predicted using population pharmacokinetic modelling and simulation based on cross-indication pharmacokinetics in other pediatric patients (pediatric psoriasis, juvenile idiopathic arthritis, pediatric Crohn's disease, and enthesitis-related arthritis). No clinical exposure data are available on the use of a loading dose in children < 6 years. The predicted exposures indicate that in the absence of methotrexate, a loading dose may lead to an initial increase in systemic exposure.

Population pharmacokinetic and pharmacokinetic/pharmacodynamic modelling and simulation predicted comparable adalimumab exposure and efficacy in patients treated with 80 mg every other week when compared with 40 mg every week (including adult patients with RA, HS, UC, CD or Ps, patients with adolescent HS, and pediatric patients  $\geq 40$  kg with CD and UC).

#### Exposure-response relationship in pediatric population

On the basis of clinical trial data in patients with JIA (pJIA and ERA), an exposure-response relationship was established between plasma concentrations and PedACR 50 response. The apparent adalimumab plasma concentration that produces half the maximum probability of PedACR 50 response (EC<sub>50</sub>) was 3  $\mu\text{g/mL}$  (95% CI: 1-6  $\mu\text{g/mL}$ ).

Exposure-response relationships between adalimumab concentration and efficacy in pediatric patients with severe chronic plaque psoriasis were established for PASI 75 and PGA clear or minimal, respectively. PASI 75 and PGA clear or minimal increased with increasing adalimumab concentrations, both with a similar apparent EC<sub>50</sub> of approximately 4.5  $\mu\text{g/mL}$  (95% CI 0.4-47.6 and 1.9-10.5, respectively).

In subjects with Behcet's disease, the mean steady-state trough adalimumab serum concentration was approximately 9  $\mu\text{g/mL}$  during the treatment of 40 mg given every other week starting 4 week after an initial 160 mg dose on week 0 followed by 80 mg on week 2 as subcutaneous injections (Japanese Subjects).

#### Elimination

Population pharmacokinetic analyzes with data from over 1,300 RA patients revealed a trend toward higher apparent clearance of adalimumab with increasing body weight. After adjustment for weight differences, gender and age appeared to have a minimal effect on adalimumab clearance. The serum levels of free adalimumab (not bound to anti-adalimumab antibodies, AAA) were observed to be lower in patients with measurable AAA.

#### Hepatic or renal impairment

Adalimumab has not been studied in patients with hepatic or renal impairment.

### **5.3 Preclinical safety data**

Non-clinical data reveal no special hazard for humans based on studies of single dose toxicity, repeated dose toxicity, and genotoxicity.

An embryo-fetal developmental toxicity/perinatal developmental study has been performed in Cynomolgus monkeys at 0, 30 and 100 mg/kg (9-17 monkeys/group) and has revealed no evidence of

harm to the fetuses due to adalimumab. Neither carcinogenicity studies, nor a standard assessment of fertility and postnatal toxicity, were performed with adalimumab due to the lack of appropriate models for an antibody with limited cross-reactivity to rodent TNF and to the development of neutralizing antibodies in rodents.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

L-lactic acid

Sucrose

Polysorbate 80

Sodium hydroxide (for pH adjustment)

Water for injections

### **6.2 Incompatibilities**

In the absence of compatibility studies, this medicinal product must not be mixed with other medicinal products.

### **6.3 Shelf life**

The expiry date of the product is indicated on the packaging materials.

### **6.4 Special precautions for storage**

Store in a refrigerator (2°C – 8°C).

Do not freeze.

Keep the pre-filled syringe or pre-filled pen in the outer carton in order to protect from light.

The pre-filled syringe or pre-filled pen may be stored at temperatures up to a maximum of 25°C for a period of up to 14 days. The pre-filled syringe or pre-filled pen must be protected from light, and discarded if not used within the 14-day period.

### **6.5 Nature and contents of container**

#### AMGEVITA 100 mg/mL - 20 mg solution for injection in pre-filled syringe

0.2 mL solution in pre-filled syringe (type I glass) with a plunger stopper (bromobutyl rubber) and a stainless steel needle with a needle shield (thermoplastic elastomer).

Pack size of one pre-filled syringe.

#### AMGEVITA 100 mg/mL - 40 mg solution for injection in pre-filled syringe

0.4 mL solution in pre-filled syringe (type I glass) with a plunger stopper (bromobutyl rubber) and a stainless steel needle with a needle shield (thermoplastic elastomer).

Pack sizes of 1, 2 or multipack containing 6 (3 packs of 2) pre-filled syringes.

Not all pack sizes may be marketed.

## AMGEVITA 100 mg/mL - 40 mg solution for injection in pre-filled pen

0.4 mL solution for injection in pre-filled pen for patient use containing a pre-filled syringe (type I glass). The pen is a single use, disposable, handheld, mechanical injection device. The needle cover of the pre-filled pen is made from synthetic rubber.

Pack sizes of 1, 2 or multipack containing 6 (3 packs of 2) pre-filled pens.

Not all pack sizes may be marketed.

### **6.6 Special precautions for disposal and other handling**

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

## **7. MANUFACTURER**

Amgen Europe B.V.  
Minervum 7061  
NL-4817 ZK Breda  
The Netherlands

## **8. LICENSE HOLDER**

Amgen Europe B.V.  
P.O. BOX 53313  
Tel - Aviv  
Israel

## **9. LICENSE NUMBER**

179-87-38162-00

*Revised in January 2026.*