



פיזר פי אף אי פרמצבטיקה ישראל בע"מ
רח' שנקר 9, ת.ד. 12133
הרצליה פיתוח, ישראל 46725
טל: 972-9-9700500 פקס: 972-9-9700501

אוקטובר 2021

רופא/ה, רוקח/ת נכבד/ה,

חברת פיזר פי אף אי ישראל בע"מ, מבקשת להודיעכם על על עדכון בעלון לרופא של התכשיר Prostin E2 3 mg. הודעה זו מפרטת את העדכונים המהווים החמרה במידע הבטיחותי בלבד, למידע מלא יש לעיין בעלון. העלון לרופא מפורסם במאגר התרופות שבמשרד הבריאות, וניתן לקבלו מודפס על ידי פניה לבעל הרישום: חברת פיזר פי אף אי פרמצבטיקה ישראל בע"מ, שנקר 9, ת.ד. 12133, הרצליה פיתוח, 46725.

Prostin E2 3mg

שם התכשיר:

Each tablet contains 3 mg dinoproston

הרכב וחוזק:

Induction of labour

התוויה מאושרת:

להלן עדכוני הבטיחות בעלון (מסומנים בצהוב):

4.2 Posology and method of administration

Usage is restricted to qualified health care professionals and to hospitals and clinics with specialised obstetric units with facilities for continuous monitoring.

The recommended dose should not be exceeded, and the dosing interval should not be shortened as this increases the risk of uterine hyperstimulation, uterine rupture, uterine haemorrhage, foetal and neonatal death.

4.4 Special warnings and precautions for use

As with any oxytocic agent, the risk of uterine rupture should be considered. Concomitant medication, maternal and foetal status should be taken into consideration in order to minimise the risk of uterine hyperstimulation, uterine rupture, uterine haemorrhage, foetal and neonatal death. Continuous electronic monitoring of uterine activity and foetal heart rate should be conducted during use of dinoprostone. Patients who develop uterine hypertonus or hypercontractility, or in whom unusual foetal heart rate patterns develop, should be managed in a manner that addresses the welfare of the foetus and mother.

Caution should be exercised in the administration of Prostin E2 Vaginal Tablets for the induction of labour in patients with:

- asthma or a history of asthma
- epilepsy or a history of epilepsy
- glaucoma or raised intra-ocular pressure
- compromised cardiovascular, hepatic, or renal function
- hypertension.
- ruptured chorioamniotic membranes.

Dinoprostone should be used with caution in patients with multiple pregnancy.

4.8 Undesirable effects

Pregnancy, puerperium and perinatal conditions: Foetal death, stillbirth, neonatal death*
(Frequency not known- cannot be estimated from the available data)

Maternal-related conditions: Uterine hypertonus, uterine rupture, abruptio placenta, pulmonary amniotic fluid embolism, rapid cervical dilatation

Foetus-related conditions: Uterine hypercontractility with/without foetal bradycardia foetal distress/altered foetal heart rate (FHR)

Neonatal conditions: Neonatal distress, neonatal death, stillbirths, low Apgar score

*Foetal death, stillbirth, and neonatal death have been reported after application of dinoprostone, especially following the occurrence of serious events such as uterine rupture (see sections 4.2, 4.3 and 4.4).