

**Patient leaflet in accordance with the
Pharmacists' Regulations (Preparations) - 1986**

This medicine is dispensed without a doctor's prescription

Carbosylane Capsules

Active ingredients:

Each blue capsule (acts in the stomach) contains:

activated charcoal 140 mg
simethicone 45 mg

Each red capsule (acts in the intestine) contains:

activated charcoal 140 mg
simethicone 45 mg

Inactive ingredients and allergens: See section 6 'Additional information'.

Read the entire leaflet carefully before you start using this medicine. This leaflet contains concise information about this medicine. If you have any further questions, consult your doctor or pharmacist.

You must use this medicine according to the instructions in the dosage section in this leaflet. Consult your pharmacist if you need additional information. You should contact your doctor if you do not feel your condition has improved or if you feel worse after 10 days.

1. What is this medicine intended for?

Against accumulation of gas in the digestive system. For the symptomatic treatment of stomach ache resulting from accumulation of gas and relief of sensation of fullness and bloating and flatulence.

Therapeutic group: Antiflatulent/intestinal adsorbent.

2. Before using this medicine

Do not use this medicine if:

- You are sensitive (allergic) to the active ingredients or to any of the other ingredients in this medicine listed in section 6.

Special warnings about using this medicine

If you receive another medical treatment keep an interval between taking Carbosylane capsules and other medicines (more than 2 hours, if possible).

Talk to your doctor or pharmacist before taking Carbosylane capsules.

Children and adolescents

This medicine is not indicated for children under the age of 6 due to the risk of choking.

Drug interactions

If you are taking or have recently taken other medicines, including nonprescription medications and dietary supplements, tell your doctor or pharmacist.

This medicine can reduce the uptake of other medicines, and these should therefore be taken at an interval from it (more than 2 hours if possible).

Using this medicine and food

Take this medicine before meals.

Pregnancy and breastfeeding

You must consult your doctor before using this medicine during pregnancy.

If you discover that you are pregnant during treatment, consult your doctor, as only a doctor can decide whether it is necessary to continue the treatment.

This medicine may be given during breastfeeding.

If you are pregnant or breastfeeding, think you may be pregnant or are planning to have a baby, consult your doctor or pharmacist for advice before taking this medicine.

3. How should you use the medicine?

Check with your doctor or pharmacist if you are not sure about your dose or about how to take this medicine.

The recommended dosage lacking other orders from the doctor:

Take a blue capsule with a red capsule three times a day before meals.

Do not chew. Swallow the capsules whole with water.

Do not exceed the recommended dose.

If you took an overdose

The use of high dose levels can result in dark colouration of the stools. Contact your doctor or pharmacist immediately.

If you forget to take the medicine

Do not take a double dose to make up for the forgotten dose.

Do not take medicines in the dark! Check the label and dose every time you take medicine. Wear glasses if you need them. If you have any further questions about using this medicine, consult your doctor or pharmacist.

4. Side effects

Like all medicines, using Carbosylane capsules may cause side effects in some users. Do not be alarmed by this list of side effects. You may not experience any of them.

- Dark discolouration of the stools may occur.
- Cases of allergic reactions (urticaria, generalised allergic reaction) have been reported after the administration of Carbosylane capsules.

- Cases of digestive disorders (gastro-intestinal) (pain, vomiting, discomfort, constipation or diarrhoea) have been reported following administration of Carbosylane capsules.

If you experience any side effect, if any side effect gets worse, or if you experience a side effect not mentioned in this leaflet, consult your doctor.

You can report side effects to the Ministry of Health by following the link 'Reporting Side Effects of Drug Treatment' on the Ministry of Health home page (www.health.gov.il) which links to an online form for reporting side effects. You can also use this link:

<https://sideeffects.health.gov.il>

You may also report to e-mail:

safety@trima.co.il

5. How to store the medicine?

- Prevent poisoning! To prevent poisoning, keep this, and all other medicines, in a closed place, out of the reach and sight of children and/or infants. Do not induce vomiting unless explicitly instructed to do so by a doctor.
- Do not use the medicine after the expiry date (exp. date) which is stated on the package/blister. The expiry date refers to the last day of that month.
- **Storage conditions:** Store below 25°C.

6. Additional information

In addition to the active ingredients, this medicine also contains:

blue capsule (acts in the stomach):

gelatin, sorbitan oleate, polysorbate 80, titanium dioxide, indogotin.

red capsule (acts in the intestine):

gelatin, sorbitan oleate, cellulose acetate phthalate, polysorbate 80, diethyl acetate, erythrosine, titanium dioxide, indogotin.

What the medicine looks like and contents of the pack:

Box of 48 (24 doses) capsules.

24 red capsules that contain a black powder.

24 blue capsules that contain a black powder.

Manufacturer's name and address:

Laboratories Grimmer, 44 avenue Georges Pompidou, 92300 Levallois-Perret, France.

Registration holder's name and address:

Trima Israel Pharmaceutical Products Maabarot Ltd., Maabarot 4023000, Israel.

Revised in November 2021 according to MOH guidelines.

Registration number of the medicine in the Ministry of Health's National Drug Registry: 134-82-21600-07.

1221F

P00001465

PIL-1221-02