





Increlex® Patient Information Card

Increlex Instructions for Use

- S Increlex should be administered twice per day
- 😚 Increlex should be administered **shortly before** or **immediately after** a meal
- If a dose of Increlex cannot be administered shortly before or just after a meal, that dose of Increlex should not be given.
- So Look out for signs and symptoms, provided to you by the attending physician or the nurse, that indicate low blood sugar levels. Keep an edible sugar source with you at all times in case they develop.
- 😚 Use a different injection site in the skin for each dose



- If you miss a dose, never increase the subsequent doses of Increlex to make up for the missed dose. The next dose should be taken at the regular time.
- Increlex should be kept in the refrigerator and must not be frozen. Store at a temperature of 2°C 8°C.

Before using the product, read the package leaflet, which includes detailed injection instructions.

Fill in the patient's name and the required dose before giving the form to the patient - it contains useful and important information about the manner of use of Increlex.

Patient name:

Dose, in units:

Contact details

Physician name:

Nurse name:

Clinic / contact person telephone number:

information. You can help by reporting any side effects that your child may experience. If your child develops side effects, including any possible side effects not listed in the package leaflet, consult with your doctor, pharmacist or nurse.

By reporting side effects, you can help provide more information about the safety of this medication.

You can report side effects to the Ministry of Health, at https://sideeffects.health.gov.il You can also report side effects to the license holder, Medison Pharma, at pv@medison.co.il

For simplicity and ease of reading, this leaflet has been phrased in the masculine form. However, the product is designed for both sexes.

This guide was reviewed and approved by the Ministry of Health in November 2021.



