

DERMOVATE OINTMENT

1. NAME OF THE MEDICINAL PRODUCT

Dermovate Ointment

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Clobetasol 17-propionate 0.05% w/w

Excipients with known effect:

Propylene Glycol

For the full list of excipients, see section 6.1

3. PHARMACEUTICAL FORM

Ointment

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

Clobetasol is a very potent topical corticosteroid indicated for adults, elderly and children over 1 year for the short term treatment only of more resistant inflammatory and pruritic manifestations of steroid responsive dermatoses unresponsive to less potent corticosteroids.

These include the following:

- Psoriasis (excluding widespread plaque psoriasis)
- Recalcitrant dermatoses
- Lichen planus
- Discoid lupus erythematosus
- Other skin conditions which do not respond satisfactorily to less potent steroids.

4.2 Posology and method of administration

Clobetasol propionate belongs to the most potent class of topical corticosteroids (Group IV) and prolonged use may result in serious undesirable effects (see section 4.4). If treatment with a local corticosteroid is clinically justified beyond 4 weeks, a less potent corticosteroid preparation should be considered. Repeated but short courses of clobetasol propionate may be used to control exacerbations (see details below).

Route of administration: Cutaneous

Ointments are especially appropriate for dry, lichenified or scaly lesions.

Adults, Elderly and Children over 1 year

Apply thinly and gently rub in using only enough to cover the entire affected area once or twice a day until improvement occurs (in the more responsive conditions this may be within a few days), then reduce the frequency of application or change the treatment to a less potent preparation. Allow adequate time for absorption after each application before applying an emollient.

Repeated short courses of clobetasol propionate may be used to control exacerbations.

In more resistant lesions, especially where there is hyperkeratosis, the effect of clobetasol can be enhanced, if necessary, by occluding the treatment area with polythene film. Overnight occlusion only is usually adequate to bring about a satisfactory response. Thereafter improvement can usually be maintained by application without occlusion.

If the condition worsens or does not improve within 2-4 weeks, treatment and diagnosis should be re-evaluated.

Treatment should not be continued for more than 4 weeks. If continuous treatment is necessary, a less potent preparation should be used.

The maximum weekly dose should not exceed 50gms/week.

Therapy with clobetasol should be gradually discontinued once control is achieved and an emollient continued as maintenance therapy.

Rebound of pre-existing dermatoses can occur with abrupt discontinuation of clobetasol.

Recalcitrant dermatoses: Patients who frequently relapse

Once an acute episode has been treated effectively with a continuous course of topical corticosteroid, intermittent dosing (once daily, twice weekly, without occlusion) may be considered. This has been shown to be helpful in reducing the frequency of relapse.

Application should be continued to all previously affected sites or to known sites of potential relapse. This regimen should be combined with routine daily use of emollients. The condition and the benefits and risks of continued treatment must be re-evaluated on a regular basis.

Paediatric population

Clobetasol is contraindicated in children under one year of age.

Children are more likely to develop local and systemic side effects of topical corticosteroids and, in general, require shorter courses and less potent agents than adults.

Care should be taken when using clobetasol propionate to ensure the amount applied is the minimum that provides therapeutic benefit.

Duration of treatment for children and infants

Courses should be limited if possible to five days and reviewed weekly. Occlusion should not be used.

Application to the face

Courses should be limited to five days if possible and occlusion should not be used.

Elderly

Clinical studies have not identified differences in responses between the elderly and younger patients. The greater frequency of decreased hepatic or renal function in the elderly may delay elimination if systemic absorption occurs. Therefore the minimum quantity should be used for the shortest duration to achieve the desired clinical benefit.

Renal / Hepatic Impairment

In case of systemic absorption (when application is over a large surface area for a prolonged period) metabolism and elimination may be delayed therefore increasing the risk of systemic toxicity. Therefore the minimum quantity should be used for the shortest duration to achieve the desired clinical benefit.

4.3 Contraindications

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

The following conditions should not be treated with clobetasol:

- Untreated cutaneous infections (fungal, bacterial, viral)
- Rosacea
- Acne vulgaris
- Pruritus without inflammation
- Perianal and genital pruritus
- Perioral dermatitis.

Clobetasol is contraindicated in dermatoses in children under one year of age, including dermatitis and nappy eruptions.

4.4 Special warnings and precautions for use

Cases of osteonecrosis serious infections (including necrotizing fasciitis) and systemic immunosuppression (sometimes resulting in reversible Kaposi's sarcoma lesions) have been reported with long-term use of clobetasol propionate beyond the recommended doses (see section 4.2). In some cases patients used concomitantly other potent oral/topical corticosteroids or immunosuppressors (e.g. methotrexate, mycophenolate mofetil). If treatment

with local corticosteroids is clinically justified beyond 4 weeks, a less potent corticosteroid preparation should be considered.

Clobetasol should be used with caution in patients with a history of local hypersensitivity to other corticosteroids or to any of the excipients in the preparation. Local hypersensitivity reactions (see section 4.8) may resemble symptoms of the condition under treatment.

Manifestations of hypercortisolism (Cushing's syndrome) and reversible hypothalamic-pituitary-adrenal (HPA) axis suppression, leading to glucocorticosteroid insufficiency, can occur in some individuals as a result of increased systemic absorption of topical steroids. If either of the above are observed, withdraw the drug gradually by reducing the frequency of application, or by substituting a less potent corticosteroid. Abrupt withdrawal of treatment may result in glucocorticosteroid insufficiency (see section 4.8).

Risk factors for increased systemic effects are:

- Potency and formulation of topical steroid
- Duration of exposure
- Application to a large surface area
- Use on occluded areas of skin (e.g. on intertriginous areas or under occlusive dressings (in infants the nappy may act as an occlusive dressing))
- Increasing hydration of the stratum corneum
- Use on thin skin areas such as the face
- Use on broken skin or other conditions where the skin barrier may be impaired
- In comparison with adults, children and infants may absorb proportionally larger amounts of topical corticosteroids and thus be more susceptible to systemic adverse effects. This is because children have an immature skin barrier and a greater surface area to body weight ratio compared with adults.

Paediatric population

In infants and children under 12 years of age, long-term continuous topical corticosteroid therapy should be avoided where possible, as adrenal suppression can occur.

Children are more susceptible to develop atrophic changes with the use of topical corticosteroids.

Duration of treatment for children and infants

Courses should be limited if possible to five days and reviewed weekly. Occlusion should not be used.

Infection risk with occlusion

Bacterial infection is encouraged by the warm, moist conditions within skin folds or caused by occlusive dressings. When using occlusive dressings, the skin should be cleansed before a fresh dressing is applied.

Use in Psoriasis

Topical corticosteroids should be used with caution in psoriasis as rebound relapses, development of tolerances, risk of generalised pustular psoriasis and development of local or systemic toxicity due to impaired barrier function of the skin have been reported in some cases. If used in psoriasis careful patient supervision is important.

Concomitant infection

Appropriate antimicrobial therapy should be used whenever treating inflammatory lesions which have become infected. Any spread of infection requires withdrawal of topical corticosteroid therapy and administration of appropriate antimicrobial therapy.

Chronic leg ulcers

Topical corticosteroids are sometimes used to treat the dermatitis around chronic leg ulcers. However, this use may be associated with a higher occurrence of local hypersensitivity reactions and an increased risk of local infection.

Application to the face

Application to the face is undesirable as this area is more susceptible to atrophic changes.

If used on the face, treatment should be limited to only 5 days.

Application to the eyelids

If applied to the eyelids, care is needed to ensure that the preparation does not enter the eye, as cataract and glaucoma might result from repeated exposure. If clobetasol does enter the eye, the affected eye should be bathed in copious amounts of water.

Visual disturbance

Visual disturbance has been reported with systemic and topical corticosteroid use. If a patient presents with symptoms such as blurred vision or other visual disturbances, the patient should be considered for referral to an ophthalmologist for evaluation of possible causes which may include cataract, glaucoma or rare diseases such as central serous chorioretinopathy (CSCR) which have been reported after use of systemic and topical corticosteroids.

Healthcare professionals should be aware that if this product comes into contact with dressings, clothing and bedding, the fabric can be easily ignited with a naked flame. Patients should be warned of this risk and advised to keep away from fire when using this product.

Dermovate Ointment contains 50 mg propylene glycol per gram of product. Propylene glycol may cause skin irritation.

Clobetasol ointment contains paraffin. Instruct patients not to smoke or go near naked flames due to the risk of severe burns. Fabric (clothing, bedding,

dressings etc) that has been in contact with this product burns more easily and is a serious fire hazard. Washing clothing and bedding may reduce product build-up but not totally remove it.

Topical steroid withdrawal syndrome

Long term continuous or inappropriate use of topical steroids can result in the development of rebound flares after stopping treatment (topical steroid withdrawal syndrome). A severe form of rebound flare can develop which takes the form of a dermatitis with intense redness, stinging and burning that can spread beyond the initial treatment area. It is more likely to occur when delicate skin sites such as the face and flexures are treated. Should there be a reoccurrence of the condition within days to weeks after successful treatment a withdrawal reaction should be suspected. Reapplication should be with caution and specialist advice is recommended in these cases or other treatment options should be considered.

4.5 Interaction with other medicinal products and other forms of interaction

Co-administered drugs that can inhibit CYP3A4 (e.g. ritonavir and itraconazole) have been shown to inhibit the metabolism of corticosteroids leading to increased systemic exposure. The extent to which this interaction is clinically relevant depends on the dose and route of administration of the corticosteroids and the potency of the CYP3A4 inhibitor.

4.6. Fertility, pregnancy and lactation

Pregnancy

There are limited data from the use of clobetasol in pregnant women.

Topical administration of corticosteroids to pregnant animals can cause abnormalities of foetal development (see Nonclinical Information)

The relevance of this finding to humans has not been established. Administration of clobetasol during pregnancy should only be considered if the expected benefit to the mother outweighs the risk to the foetus. The minimum quantity should be used for the minimum duration.

Breast-feeding

The safe use of topical corticosteroids during lactation has not been established.

It is not known whether the topical administration of corticosteroids could result in sufficient systemic absorption to produce detectable amounts in breast milk. Administration of clobetasol during lactation should only be considered if the expected benefit to the mother outweighs the risk to the infant.

If used during lactation clobetasol should not be applied to the breasts to avoid accidental ingestion by the infant.

Fertility

There are no data in humans to evaluate the effect of topical corticosteroids on fertility

Clobetasol administered subcutaneously to rats had no effect upon mating performance; however, fertility was decreased at the highest dose (see section 5.2).

4.7 Effects on ability to drive and use machines

There have been no studies to investigate the effect of clobetasol on driving performance or the ability to operate machinery. A detrimental effect on such activities would not be anticipated from the adverse reaction profile of topical clobetasol.

4.8 Undesirable effects

Adverse drug reactions (ADRs) are listed below by MedDRA system organ class and by frequency. Frequencies are defined as: very common ($\geq 1/10$), common ($\geq 1/100$ and $< 1/10$), uncommon ($\geq 1/1,000$ and $< 1/100$), rare ($\geq 1/10,000$ and $< 1/1,000$) and very rare ($< 1/10,000$), including isolated reports.

Post-marketing data

Infections and Infestations

Very rare Opportunistic infection

Immune System Disorders

Very rare Hypersensitivity, generalised rash

Endocrine Disorders

Very rare Hypothalamic-pituitary adrenal (HPA) axis suppression:
Cushingoid features: (e.g. moon face, central obesity),
delayed weight gain/growth retardation in children,
osteoporosis, hyperglycaemia/glucosuria, hypertension,
increased weight/obesity, decreased endogenous cortisol
levels, alopecia, trichorrhexis

Skin and Subcutaneous Tissue Disorders

Common Pruritus, local skin burning /skin pain

Uncommon Skin atrophy*, striae*, telangiectasias*

Very rare Skin thinning*, skin wrinkling*, skin dryness*, pigmentation
changes*, hypertrichosis, exacerbation of underlying symptoms,
allergic contact dermatitis/dermatitis, pustular psoriasis,
erythema, rash, urticaria, acne

Not known Withdrawal reactions - redness of the skin which may extend
to areas beyond the initial affected area, burning or stinging
sensation, itch, skin peeling, oozing pustules. (see section 4.4)

**Skin features secondary to local and/or systemic effects of hypothalamic-pituitary adrenal (HPA) axis suppression.*

General Disorders and Administration Site Conditions

Very rare Application site irritation/pain

Eye disorders

Very rare Cataract, central serous chorioretinopathy, glaucoma

Not known (cannot be estimated from available data) Vision, blurred

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the risk/benefit balance of the medicinal product.

Any suspected adverse events should be reported to the Ministry of Health according to the National Regulation by using an online form

<https://sideeffects.health.gov.il>

Additionally, you should also report to GSK Israel (il.safety@gsk.com).

4.9 Overdose

Symptoms

Topically applied clobetasol may be absorbed in sufficient amounts to produce systemic effects. Acute overdosage is very unlikely to occur, however, in the case of chronic overdosage or misuse the features of hypercortisolism may occur (see section 4.8).

Management

In the event of overdose, clobetasol should be withdrawn gradually by reducing the frequency of application or by substituting a less potent corticosteroid because of the risk of glucocorticosteroid insufficiency.

Further management should be as clinically indicated or as recommended by the national poisons centre, where available.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Corticosteroids, very potent (group IV)

ATC code: D07AD

Mechanism of action

Topical corticosteroids act as anti-inflammatory agents via multiple mechanisms to inhibit late phase allergic reactions including decreasing the density of mast cells, decreasing chemotaxis and activation of eosinophils, decreasing cytokine production by lymphocytes, monocytes, mast cells and eosinophils, and inhibiting the metabolism of arachidonic acid.

Pharmacodynamic effects

Topical corticosteroids have anti-inflammatory, antipruritic, and vasoconstrictive properties.

5.2 Pharmacokinetic properties

Absorption

Topical corticosteroids can be systemically absorbed from intact healthy skin. The extent of percutaneous absorption of topical corticosteroids is determined by many factors, including the vehicle and the integrity of the epidermal barrier. Occlusion, inflammation and/or other disease processes in the skin may also increase percutaneous absorption.

Mean peak plasma clobetasol propionate concentrations of 0.63 nanograms/ml occurred in one study eight hours after the second application (13 h after an initial application) of 30 g clobetasol propionate 0.05 % ointment to normal individuals with healthy skin. Following the application of a second dose of 30 g clobetasol propionate cream 0.05 %, mean peak plasma concentrations were slightly higher than the ointment and occurred 10 h after application. In a separate study, mean peak plasma concentrations of approximately 2.3 nanograms/ml and 4.6 nanograms/ml occurred respectively in patients with psoriasis and eczema three hours after a single application of 25 g clobetasol propionate 0.05 % ointment.

Distribution

The use of pharmacodynamic endpoints for assessing the systemic exposure of topical corticosteroids is necessary due to the fact that circulating levels are well below the level of detection.

Metabolism

Once absorbed through the skin, topical corticosteroids are handled through pharmacokinetic pathways similar to systemically administered corticosteroids. They are metabolised, primarily in the liver.

Elimination

Topical corticosteroids are excreted by the kidneys. In addition, some corticosteroids and their metabolites are also excreted in the bile.

5.3 Preclinical safety data

Carcinogenesis / Mutagenesis

Carcinogenesis

Long-term animal studies have not been performed to evaluate the carcinogenic potential of clobetasol propionate.

Genotoxicity

Clobetasol propionate was not mutagenic in a range of *in vitro* bacterial cell assays.

Reproductive Toxicology

Fertility

In fertility studies, subcutaneous administration of clobetasol propionate to rats at doses of 6.25 to 50 micrograms/kg/day produced no effects on mating, and fertility was only decreased at 50 micrograms/kg/day.

Pregnancy

Subcutaneous administration of clobetasol propionate to mice (≥ 100 micrograms/kg/day), rats (400 micrograms/kg/day) or rabbits (1 to 10 micrograms/kg/day) during pregnancy produced foetal abnormalities including cleft palate and intrauterine growth retardation.

In the rat study, where some animals were allowed to litter, developmental delay was observed in the F1 generation at ≥ 100 micrograms/kg/day and survival was reduced at 400 micrograms/kg/day. No treatment-related effects were observed in F1 reproductive performance or in the F2 generation.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Propylene glycol
Sorbitan sesquioleate
White soft paraffin

6.2 Incompatibilities

None known.

6.3 Shelf life

The expiry date of the product is indicated on the packaging materials.
Use within 3 months after opening.

6.4 Special precautions for storage

Store below 25°C.

6.5 Nature and contents of container

Membrane-sealed, collapsible, aluminium tubes either coated internally with epoxy resin based lacquer or uncoated. The tubes are sealed with a latex band sealant and wadless polypropylene caps.

Pack sizes: 25 g, 30 g or 100 g.

Not all pack sizes may be marketed

6.6 Special precautions for disposal and other handling

Patients should be advised to wash their hands after applying Dermovate Ointment unless it is the hands that are being treated.

7. **MANUFACTURER**
GlaxoSmithKline Trading Services Limited, Dublin, Ireland
8. **LICENSE HOLDER AND IMPORTER**
GlaxoSmithKline (Israel) Ltd., 25 Basel St., Petach Tikva
9. **LICENSE NUMBER**
027-57-22063

Revised in June 2022 according to MOHs guidelines

Trade marks are owned by or licensed to the GSK group of companies.
©2022 GSK group of companies or its licensor

Der Oin DR v8