

## **SUMMARY OF PRODUCT CHARACTERISTICS**

### **1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Flugestone Acetate Intervet 20 mg Veterinary

### **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

Each polyester polyurethane sponge contains

#### **Active substance(s)**

Flugestone acetate, 20 mg.

#### **List of excipients**

Excipients qsp 1 sponge.

For a full list of excipients, see section 6.1

### **3. PHARMACEUTICAL FORM**

Vaginal sponge.

White cylindrical polyester polyurethane foam equipped with string.

### **4. CLINICAL PARTICULARS**

#### **4.1 Target species**

Sheep (ewes and ewe-lambs).

#### **4.2 Indications for use**

In ewes and ewe lambs, in combination with PMSG (Pregnant Mare Serum Gonadotrophin)

- Induction and synchronization of oestrus and ovulation (non cycling ewes during seasonal anoestrus and ewe lambs).
- Synchronization of oestrus and ovulation (cycling ewes and ewe-lambs).

#### **4.3 Contraindications**

Please refer to section 4.7 and section 4.8.

#### **4.4 Special warnings**

None.

#### **4.5 Special precautions for use**

- (i) Special precautions for use in animals

- The repeated treatment with the product combined with PMSG may trigger the appearance of PMSG antibodies in some ewes. This in turn may affect the time of ovulation and result in reduced fertility when combined with fixed time artificial insemination at 55h following sponge removal.
- The repeated use of sponges within one year has not been studied.
- The use of a vaginal applicator designed for ewes or ewe lambs is recommended to correctly insert sponges and to avoid vaginal injuries.

(ii) Special precautions to be taken by the person administering the medicinal product to animals

- Direct contact with the skin should be avoided. Personal protective clothing (single use gloves) should be worn when handling the product. If accidental contact with the skin occurs, wash the affected zone with soap and water. Wash hands after treatment and before meals.
- Human exposure to this product can affect fertility.
- Women who are pregnant, or suspected to be pregnant, must not use the product.

#### **4.6 Adverse reactions**

A muco-purulent discharge may occasionally be observed at sponge removal. It is not associated with clinical signs and does not alter fertility.

Reporting suspected adverse reactions:

Reporting suspected adverse reactions after authorization of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product.

Any suspected adverse events should be reported to the Ministry of Health according to the National Regulation by using an online form

<https://sideeffects.health.gov.il>

#### **4.7 Use during pregnancy and lactation**

Can be used during lactation.

The use is not recommended during gestation.

#### **4.8 Interaction with other medicinal products and other forms of interaction**

The sponges should not be used in conjunction with alcohols, cresols, phenols, sheep dips or similar disinfectants.

#### **4.9 Amounts to be administered and administration route**

The dose is one sponge per animal irrespective of body weight, breed, type (dairy or meat) and season.

The sponge is inserted intra-vaginally using an applicator.

Duration of sponge residence is 14 days. At the end of the administration period,

the sponge is gently removed by pulling on its string. To obtain an optimal synchronization of ovulation, an intra-muscular injection of PMSG (range 300-700 IU) is recommended (i.m.) at the time of sponge removal. In case fixed time artificial insemination is applied, it is recommended to perform it 55 h after sponge removal.

#### **4.10 Overdose (symptoms, emergency procedures, antidotes) (if necessary)**

A five time overdose of flugestone acetate (100 mg/sponge) did not result in observable side effects.

#### **4.11 Withdrawal periods**

Meat: 2 days after withdrawal of sponges.  
Milk: zero hours, including the treatment time.

### **5. PHARMACOLOGICAL PROPERTIES**

**Pharmacotherapeutic group:** progestagen  
**ATC vet code:** QG03D

#### **5.1 Pharmacodynamic properties**

Flugestone acetate is a synthetic analogue of progesterone. It is approximately 20 fold more potent than progesterone and displays progestational activity but no anti-progestational, anti-androgenic or androgenic properties together with a low glucocorticoid activity.

Owing to its binding to the progesterone receptors, flugestone acetate acts by negative feedback on the hypothalamo-pituitary axis, suppressing pituitary release of gonadotrophins and therefore terminal follicular growth and ovulation.

#### **5.2 Pharmacokinetic properties**

Flugestone acetate is readily absorbed during the 12-14 days period of intravaginal administration.  $T_{max}$  ranges between 8 and 24 h, whereas  $C_{max}$  varies between 1.4 and 3.7 ng/ml. Steady state is reached quickly following onset of the treatment. Plasma cronolone concentrations are relatively constant throughout treatment. One day after removal of the Flugestone Acetate Intervet 20 mg Veterinary, flugestone acetate concentrations have dropped below the limit of quantification (LOQ = 0.04 ng/mL).

Flugestone acetate is metabolised into hydroxylated metabolites, which are excreted in faeces and urine.

### **6. PHARMACEUTICAL PARTICULARS**

#### **6.1 List of excipients**

Hydroxypropylcellulose (e.g. Klucel E, or equivalent)  
Polyethylene glycol 4000

Purified water  
Polyester Polyurethane Sponge  
Multifilament polyamide string high resistance (1.17 g/Km)

## **6.2 Incompatibilities**

None known.

## **6.3 Shelf life**

The expiry date of the product is indicated on the packaging materials.

## **6.4 Special precautions for storage**

Store below 25°C in a dry place.

Store the product in its original immediate packaging.

Once packaging is opened, any unused product should be discarded.

## **6.5 Nature and composition of immediate packaging**

Bags made of polyester/ aluminium/ polyethylene containing 10 sponges, 25 sponges or 50 sponges.

## **6.6 Special precautions for the disposal of unused medicinal product or waste materials, if any**

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of as toxic waste, do not throw to sewer.

## **7. MANUFACTURER**

Intervet Productions SA  
Rue de Lyons  
27460 Igoville  
France

## **8. REGISTRATION HOLDER**

Intervet Israel Ltd.  
Industrial Zone Neve Ne'eman 2  
Hod Hasharon 45240  
Israel

## **9. LICENSE NUMBER**

157-89-34250-00

Revised in June 2022 according to MoHs guidelines.