

The medicine is dispensed with a veterinarian's prescription only  
For animal use only

### 1. NAME OF THE VETERINARY MEDICINE, ITS FORM AND STRENGTH

Phibrocol 10% Veterinary  
A solution for administration in drinking water

### 2. ACTIVE INGREDIENT AND QUANTITY IN A DOSAGE UNIT

Florfenicol 11.22% w/v  
1 gram of solution (equivalent to 0.89 ml) contains 100 mg florfenicol  
For the list of inactive ingredients in the preparation – see section 13 “Further Information”.

### 3. WHAT IS THE MEDICINE INTENDED FOR?

This medicine is intended for the treatment of infections susceptible to florfenicol in chickens.

**Therapeutic group:** An antibiotic of the Amphenicol group with a broad spectrum of activity against Gram-positive and Gram-negative bacteria.

### 4. CONTRAINDICATIONS

Do not use in hens laying eggs for human consumption.  
Do not administer to breeding poultry. The preparation may cause embryonic death and a sharp decrease in hatchability.

Do not use in case of hypersensitivity to the active ingredient florfenicol or to any of the ingredients of the medicine.

### 5. SIDE EFFECTS

A decrease in food and water consumption, as well as loose stool (diarrhea), were observed after use, and disappeared after stopping treatment.

Side effects can be reported to the Ministry of Health by clicking on the link “Report Side Effects of Drug Treatment” found on the Ministry of Health homepage ([www.health.gov.il](http://www.health.gov.il)) that directs you to the online form for reporting side effects, or by entering the link:  
<https://sideeffects.health.gov.il>

### 6. TARGET ANIMALS

Chickens.

### 7. METHOD OF ADMINISTRATION AND DOSAGE

30 mg of active ingredient per kilogram live weight in the drinking water (300 grams of solution per every ton of live weight). For administration in bulk in the drinking water for 6-8 hours over 3-5 days, or according to the recommendation of the attending veterinarian.

Below is a table of doses according to the weight of the birds:

Live weight	Grams of active ingredient per day	Solution weight per day (grams)	Solution volume per day (ml)
1 ton	30	300	270
2 ton	60	600	535
3 ton	90	900	800
4 ton	120	1200	1070

### 8. HOW TO USE THE PREPARATION

For administration in drinking water according to the instructions in section 7. A fresh solution should be prepared every day.

### 9. WITHDRAWAL PERIOD

Withdrawal time in chickens: 7 days.

### 10. WARNINGS

None

### 11. STORAGE INSTRUCTIONS

- Avoid poisoning! This medicine and any other medicine should be kept in a safe place out of the reach and sight of children and/or infants, in order to avoid poisoning.
- Do not use the medicine after the expiry date (exp. date) that appears on the package. The expiry date refers to the last day of that month.

- Stability in drinking water: 24 hours.
- Stability of the preparation after first opening: 48 hours.
- Storage conditions: Store in a shaded place, below 25°C.

### 12. INSTRUCTIONS REGARDING DISPOSAL OF THE PREPARATION/ REMNANTS OF THE PREPARATION AFTER USE

- Any veterinary medical preparation that was unused or any substance remaining after using the veterinary medical product must be disposed of as toxic waste; do not throw into a wastewater drain.

### 13. FURTHER INFORMATION

In addition to the active ingredient(s), the medicine also contains:

Polyethylene glycol, Propylene glycol, Soluphor p (pyrrolidone)

- What does the medicine look like and what is the content of the package: A bottle containing a yellowish liquid.
- Package sizes: 1 kg
- **License holder:** Abic Veterinary Products, P.O.B. 489 Beit Shemesh Industrial Zone

**This leaflet was revised in January 2022 according to MOH guidelines.**

Registration number of the medicine in the National Drug Registry of the Ministry of Health: 169-01-35774-00