

Consumer leaflet for a Veterinary Product

This medicine is marketed according to a veterinarian's prescription only.
For animal use only

1. Name of the veterinary medicine, form and strength

Galfenicol Solution Veterinary, solution for drinking water for chickens, containing Florfenicol 10%

2. Active ingredient and quantity in a single dose

Florfenicol 10% w/v

For full list of inactive ingredients – see in section 13 "additional information"

3. What is the medicine intended for

Treatment of bacterial infections susceptible to florfenicol in chickens

Therapeutic group: Antibacterial product of the Amphenicols group, with a wide range activity against gram-positive and gram-negative bacteria.

4. Contra-indications

Do not use in egg laying hens.

Do not use in birds for reproduction. The product might cause fetal mortality and a sharp decrease in hatching.

Do not use in case of hypersensitivity to the active ingredient (florfenicol).

5. Side effects

decrease in food and water consumption were observed following treatment, as well as loose stool (diarrhea), which disappeared following treatment stopping.

Side effects can be reported to the Ministry of Health by clicking on the link "Adverse Drug Reactions Report" that appears on the home page of the Ministry of Health web site (www.health.gov.il), which leads to an online form for reporting side effects. Alternatively you can use the following link: [/https://sideeffects.health.gov.il](https://sideeffects.health.gov.il)

6. Target animals: Chickens

7. Dosage and administration

30 mg/ kg body weight administered collectively in the drinking water during 6-8 hours for 3-5 days, or according to recommendation of the attending veterinarian.

8. How to use the product

For administration in drinking water according to the instructions in paragraph 7.

9. Withdrawal period - for slaughter for meat: 7 days.

10. Warnings

- Special warnings regarding the safety for the person administering the veterinary medicinal product to animals

The inactive ingredient N-methylpyrrolidone (NMP) can cause damage to

the fetus, therefore women of childbearing potential should take great care to avoid spilling the product on the skin during treatment with the product. If you are pregnant, think you may be pregnant or are trying to conceive, do not treat this medicine.

11. Storage instructions

- Avoid poisoning! This medicine, and any other medicine, must be kept in a closed place outside the reach of children and/or infants to prevent accidental poisoning.
- Do not use the medicine after its expiration date (exp. date) as it appears on the package. The expiration date refers to the last day of the stated month.
- Storage conditions: Store below 25°C.
- Shelf life after dilution in drinking water: 24 hours.

12. Instructions for disposing of the product / remaining product at the end of its use

Any veterinary medical product that was unused or any substance remaining after using the veterinary medical product must be disposed of as toxic waste; do not throw into the sewage system.

13. Additional information

- In addition to the active ingredient, this medicine also contains Propylene glycol, n-methylpyrrolidone
- What does the medicine look like and what is the content of the package – clear, colorless to yellowish solution
- Pack sizes 500 ml, 1 L. Not all pack sizes may be marketed.

Manufacturer and License holder: BIOVAC LTD., 30 Hailan St., OR-AKIVA 3060000

Revised on November 2022 in accordance with the Ministry of Health instructions

Registration number of this medicine in the Ministry of Health State Medicine Registry: 149-46-33423