

Patient leaflet in accordance with the Pharmacists' Regulations (Preparations) - 1986

This medicine is dispensed with a physician's prescription only

**LYUMJEV 100 units/ml
Solution for injection in a vial**

Active ingredient and its quantity:

Each vial contains:

insulin lispro 100 units/ml (zinc crystals)

Inactive ingredients and allergens in the preparation: see section 2 "Important information about some of the ingredients of this medicine" and section 6 "Additional Information".

Read this patient leaflet carefully in its entirety before using this medicine. This leaflet contains concise information about this medicine. If you have any further questions, contact your doctor or pharmacist.

This medicine has been prescribed to treat your illness. Do not pass it on to others. It may harm them, even if it seems to you that their illness is similar.

1. WHAT IS THIS MEDICINE INTENDED FOR?

Lyumjev 100 units/ml is used for the treatment of diabetes mellitus in adults.

Therapeutic group: drugs used in diabetes, insulins and analogues for injection, fast-acting.

Lyumjev 100 units/ml solution for injection in a vial contains the active ingredient insulin lispro. It is a mealtime insulin that works more quickly than other medicines containing insulin lispro. Lyumjev contains ingredients that speed up absorption of insulin lispro into the body.

Diabetes is a condition in which your body does not make enough insulin or does not use insulin effectively, which results in effects such as high levels of sugar in the blood. Lyumjev is an insulin medicine that is used in the treatment of diabetes and so controls blood sugar. Effective treatment of diabetes, with good control of blood sugar, prevents long-term complications from your diabetes.

Treatment with Lyumjev helps to control blood sugar in the long term and prevent complications from your diabetes. Lyumjev has its maximum effect 1 to 3 hours after injection and the effect lasts for up to 5 hours. You should use Lyumjev 0-2 minutes before the start of the meal. In exceptional cases, Lyumjev can be injected up to 20 minutes after starting the meal.

Your doctor may tell you to use Lyumjev as well as a longer- or intermediate-acting insulin. Do not change your insulin unless your doctor tells you to.

2. BEFORE USING THIS MEDICINE

Do not use this medicine if:

- you are sensitive (**allergic**) to the active ingredient (insulin lispro) or any of the other ingredients of this medicine (listed in section 6).
- you think your blood sugar is dropping (**hypoglycemia**). Further on, this leaflet tells you how to deal with low blood sugar (see section 3 under "If you accidentally take a higher dose").

Special warnings regarding the use of this medicine

Talk to your doctor, pharmacist or nurse before using Lyumjev.

If you cannot see very well, you will need help from someone who has been trained to give injections.

- **Low blood sugar** (hypoglycemia).

Low blood sugar can be serious and untreated hypoglycemia may even lead to death.

Lyumjev starts to lower blood sugar faster than some other mealtime insulins. If hypoglycemia occurs, you may experience it earlier after an injection of Lyumjev. If you often have hypoglycemia or have difficulty recognizing it, please discuss this with your doctor or nurse. If your blood sugar levels are well controlled by your current insulin therapy or after long duration of diabetes, you may not feel the warning symptoms when your blood sugar is falling too low. Warning signs are listed later in this leaflet. For symptoms see “Common problems of diabetes”.

You must think carefully about when to have your meals, how often to exercise and how much to do. You must also keep a close watch on your blood sugar levels by testing your blood sugar often. Making changes to the types of insulin you use may cause your blood sugar to rise or fall too much. It may be necessary to increase the frequency of blood sugar testing if you are at risk of low blood sugars. Your doctor may need to change the doses of your other diabetes medicines.

- **High blood sugar** (hyperglycemia).

Stopping or not taking enough insulin may lead to high blood sugar (hyperglycemia) and diabetic ketoacidosis, serious conditions that can even lead to death. For symptoms see “Common problems of diabetes”.

- If you are using an insulin pump and it stops working you will need to fix the problem immediately as this can lead to high blood sugar. You may need to take an injection of Lyumjev using an insulin pen or a syringe if your pump stops working.
- If your insulin treatment is being combined with one of a class of diabetes medicines called thiazolidinediones or glitazones, such as pioglitazone, tell your doctor as soon as possible if you get signs of heart failure such as unusual shortness of breath or rapid increase in weight or localized swelling caused by fluid retention (oedema).
- If you have a serious allergic reaction to insulin or any of the ingredients in Lyumjev, stop using this medicine and contact emergency medical services straight away.
- Always check the pack and the label for the name and type of the insulin when you get it from your pharmacy. Make sure you get the Lyumjev that your doctor has told you to use.
- Keep the carton, or keep a note of the batch number on the carton. If you have a side effect, you can then provide that number when you report the adverse side effect. See “Reporting of side effects”.
- Always use a new needle for each injection to prevent infections and blocked needles. If a needle is blocked replace it with a new needle.

- **Skin changes at the injection site.**

The injection site should be rotated to prevent skin changes such as lumps under the skin. The insulin may not work well if you inject into a lumpy area (See “How to use this medicine?”). Contact your doctor if you are currently injecting into a lumpy area before you start injecting into a different area. Your doctor may tell you to check your blood sugar more closely, and to adjust your insulin or your other antidiabetic medications dose.

Children and adolescents

This medicine is not intended for children and adolescents under 18 years of age.

There is no information on the safety and efficacy of this medicine in children and adolescents under 18 years of age.

Drug Interactions

If you are taking or have recently taken other medicines, including nonprescription medications and nutritional supplements, inform your doctor or pharmacist. Some medicines affect your blood sugar level - this may mean your insulin dose has to change.

Your blood sugar levels may fall (hypoglycemia) if you take:

- other medicines for diabetes (oral and injectable)
- sulphonamide antibiotics (for infections)
- acetylsalicylic acid (for pain and mild fever and to prevent blood clotting)
- some antidepressants (monoamine oxidase inhibitors (MAOIs) or selective serotonin reuptake inhibitors (SSRIs))
- some angiotensin converting enzyme (ACE) inhibitors (for example captopril, enalapril) (for some heart problems or high blood pressure)
- angiotensin II receptor blockers (for high blood pressure or heart problems)
- somatostatin analogues (such as octreotide, which are used to treat a rare condition involving too much growth hormone)

Your blood sugar levels may rise (hyperglycemia) if you take:

- danazol (for endometriosis)
- the contraceptive pill (birth control pills)
- thyroid hormone replacement therapy (for thyroid problems)
- human growth hormone (for growth hormone deficiency)
- diuretics (for high blood pressure or if you have a build-up of water in your body)
- sympathomimetic agents (for serious allergic reactions or used in some cold remedies)
- corticosteroids (to treat asthma or autoimmune conditions)

Beta-blockers (used for high blood pressure, arrhythmia or angina) make it harder to recognize the warning signs of low blood sugar.

Using this medicine and food

You should use Lyumjev 0-2 minutes before the start of the meal. In exceptional cases, Lyumjev can be injected up to 20 minutes after starting the meal.

Using this medicine and alcohol consumption

Your blood sugar levels may either rise or fall if you drink alcohol. Therefore, the amount of insulin needed may change. You should therefore monitor your blood sugar level more often than usual.

Pregnancy and breastfeeding

There is no available data on the use of Lyumjev in pregnant women.

If you are planning to get pregnant, think you may be pregnant, are pregnant or breastfeeding, ask your doctor, nurse or pharmacist for advice before taking this medicine. The amount of insulin you need usually falls during the first 3 months of pregnancy and increases for the remaining 6 months. After you have had your baby your insulin requirements will likely return to how much you needed before your pregnancy .

There is no available data on the use of Lyumjev in breastfeeding women.

Driving and using machines

Your ability to concentrate and react may be reduced if you have hypoglycemia. Please keep this possible problem in mind in all situations where you might put yourself and others at risk (e.g. driving a car or using machines). You should contact your doctor about driving if you have:

- frequent episodes of hypoglycemia
- reduced or absent warning signs of hypoglycemia

Important information about some of the ingredients of this medicine

This medicine contains less than 1 mmol sodium (23 mg) per dose, that is to say essentially 'sodium-free'.

3. HOW SHOULD YOU USE THE MEDICINE?

Always use this medicine according to your doctor's instructions. You must check with the doctor or pharmacist if you are not sure about the dosage and manner of treatment with this preparation. They will have told you exactly how much Lyumjev to use, when to use it, and how often. They will also tell you how often to visit your diabetes clinic .

The dosage and manner of treatment will be determined only by the doctor.

Do not exceed the recommended dose.

You should always have spare insulin and another injection device in case you need them.

If you are blind or visually impaired, you will need help from someone to make your injections.

Your doctor may tell you to use Lyumjev as well as a longer- or intermediate-acting insulin. Inject them separately. Lyumjev should not be mixed with any other insulin.

When to inject Lyumjev

Lyumjev is a mealtime insulin for subcutaneous injection and should be administered 0-2 minutes before the start of the meal. In exceptional cases, Lyumjev can be injected up to 20 minutes after starting the meal.

How much insulin to use

Your doctor will work out your dose based on your blood sugar and body weight and explain

- How much Lyumjev you need at each meal.
- How and when to check your blood sugar level .
- How to change your insulin dose depending on your blood sugar levels.
- What to do if you change your diet, or change how much you exercise, if you are ill or if you are using other medicines.
- If you change the type of insulin you use, you may have to take more or less than before. This might just be for the first injection or it may be a gradual change over several weeks or months.

Do not use Lyumjev

- If it does not look like water. Lyumjev must be clear, have no color and no solid pieces in it. Check each time you inject yourself.
- If Lyumjev has not been stored correctly (see section 5 "How to store the medicine?").
- If the plastic cap of the vial is damaged, do not use.

Where to inject Lyumjev

- Inject Lyumjev under the skin (subcutaneous injection).
- Do not inject yourself directly into a vein. Only your physician can administer Lyumjev by the intravenous use. He will only do this under special circumstances such as surgery or if you are ill and your sugar levels are too high.
- Make sure you inject at least 1 cm from the last injection and that you 'rotate' the places you inject (upper arm, thigh, buttocks or abdomen), as you have been taught.
- If you need to inject another insulin at the same time as Lyumjev, use a different injection site.

How to inject Lyumjev from a vial

- First wash your hands.
- Before you make an injection, clean your skin as you have been instructed. Clean the rubber stopper on the vial, but do not remove the stopper.
- Use a new, sterile syringe and needle to pierce the rubber stopper and draw in the amount of Lyumjev you want (use a syringe with 100 unit markings). Your doctor, nurse or clinic will tell you how to do this. **Do not share your needles and syringes.**

- Inject under the skin, as you were taught. After your injection, leave the needle in the skin for 5 seconds to make sure you receive the full dose.

Using Lyumjev in an insulin pump

- Only certain insulin infusion pumps may be used to infuse Lyumjev.
- Carefully follow the instructions supplied with your infusion pump.
- Be sure to use the correct reservoir and catheter for your pump. It is important to use the correct needle length on the filling system to avoid damaging the pump.
- Change the infusion set (tubing and needle) according to the instructions supplied with the infusion set.
- If repeated or severe low blood sugar levels occur, tell your doctor or nurse.
- A pump malfunction or obstruction of the infusion set can result in a rapid rise in sugar levels. If you think Lyumjev is not flowing, follow the pump instructions and if appropriate, notify your doctor or nurse.
- You may need to take an injection of Lyumjev if your pump does not work properly.

After injecting

If you are unsure how much you have injected then check your blood sugar levels before deciding if you need another injection.

If you accidentally take a higher dose

If you inject too much Lyumjev, or are unsure how much you have injected, a low blood sugar may occur. Check your blood sugar.

If your blood sugar is low (hypoglycemia) and you can treat yourself, eat glucose tablets, sugar or drink a sugary drink. Then eat fruit, biscuits, or a sandwich, as your doctor or nurse has advised you and have some rest. This will often get you over a low blood sugar or a minor insulin overdose. Check your blood sugar again after 15-20 mins until blood sugar is stabilized.

If you are unable to treat yourself (severe hypoglycemia) because you feel too dizzy, weak, confused, have difficulty talking, lose consciousness or have a seizure you may need to be treated with glucagon. This can be given by someone who knows how to use it. Eat glucose or sugar after the glucagon. If glucagon does not work, you will have to go to the hospital or call emergency services. Ask your doctor to tell you about glucagon.

Tell everyone you spend time with that you have diabetes. Tell them what could happen if your blood sugar gets too low, including the risk of passing out.

Let them know that if you pass out, they must: turn you on your side to avoid choking, get medical help straight away and not give you any food or drink because you may choke.

If you have taken an overdose or if a child has accidentally swallowed the medicine, contact a doctor or hospital Emergency Room immediately and bring the medicine pack with you.

If you forget to take the medicine

If you forget to use your insulin or you use less than you should, or are unsure how much you have injected, your blood sugar may get too high (hyperglycemia). Check your blood sugar level to decide if an insulin dose is needed. Resume your usual dosing schedule at your next meal.

If you stop taking the medicine

Do not stop or change your insulin unless your doctor tells you to. If you use less Lyumjev than you should, a high blood sugar may occur.

If high blood sugar (hyperglycemia) is not treated it can be very serious and cause headaches, nausea, vomiting, abdominal pain, dehydration, unconsciousness, coma or even death (see section 4).

Three simple steps to reduce your risk of hypoglycemia or hyperglycemia are:

- Always keep spare syringes and a spare vial of Lyumjev.
- Always carry something to show you are diabetic.
- Always carry sugar with you.

Do not take medicines in the dark! Check the label and the dose each time you take a medicine.

Wear glasses if you need them.

If you have any further questions regarding the use of this medicine, consult the doctor or pharmacist.

4. SIDE EFFECTS

Like all medicines, this medicine can cause side effects in some users. Do not be alarmed while reading the list of side effects. You may not experience any of them.

Low blood sugar (hypoglycemia) is very common with insulin treatment (may affect more than 1 in 10 users). It can be very serious. If your blood sugar level falls too much you may become unconscious. Serious hypoglycemia may cause brain damage and may be life-threatening. If you have symptoms of low blood sugar, act **immediately** to increase your blood sugar level. See section 3 under 'If you accidentally take a higher dose'.

Allergic reactions are common (may affect up to 1 in 10 users). They may be severe and they may include the following symptoms:

- rash over the whole body
- difficulty in breathing
- wheezing
- blood pressure dropping
- heart beating fast
- sweating

If you have a serious allergic reaction (including an anaphylactic attack) to insulin or any of the ingredients in Lyumjev, stop using this medicine and contact emergency medical services straight away.

Other side effects include:

Very Common Side Effects

Infusion site reactions. Some people get rash, redness, inflammation, pain, swelling, bruising or itching around the area of the insulin infusion. If you have infusion site reactions, tell your doctor.

Common Side Effects

Injection site reactions. Some people get rash, redness, inflammation, pain, swelling, bruising or itching around the area of the insulin injection. This usually clears up in a few minutes to a few weeks without needing to stop Lyumjev. If you have injection site reactions, tell your doctor.

Uncommon Side Effects (may affect up to 1 in 100 users)

Skin changes at the injection site: If you inject insulin too often at the same place, the fatty tissue may either shrink (lipoatrophy) or thicken (lipohypertrophy). Lumps under the skin may also be caused by the build-up of a protein called amyloid (cutaneous amyloidosis). The insulin may not work very well if you inject into a lumpy area. Change the injection site with each injection to help prevent these skin changes.

Other potential side effects

Swelling in arms or ankles due to fluid retention (oedema) particularly at the start of insulin therapy or during a change in your diabetes medicines.

If a side effect appears, if any side effect gets worse, or if you suffer from a side effect not mentioned in the leaflet, you should consult the doctor.

Reporting of Side Effects

Side effects can be reported to the Ministry of Health by clicking on the link “Reporting Side Effects due to Drug Treatment” that can be found on the Home Page of the Ministry of Health’s website (www.health.gov.il), which refers to the online form for reporting side effects, or via the following link: <https://sideeffects.health.gov.il>

Common problems of diabetes

Low blood sugar

Low blood sugar (hypoglycemia) means there is not enough sugar in the blood. This can be caused if:

- you take too much Lyumjev or other insulin;
- you miss or delay meals or change your diet;
- you exercise or work too hard just before or after a meal;
- you have an infection or illness (especially diarrhoea or vomiting);
- there is a change in your need for insulin for example if you lose weight; or you have trouble with your kidneys or liver which gets worse.

See section “If you accidentally take a higher dose”.

The first symptoms of low blood sugar usually come on quickly and include the following:

- tiredness
- rapid heart beat
- nervousness or shakiness
- nausea (feeling sick)
- headache
- cold sweat

If you are not confident about recognizing your warning symptoms, avoid situations such as driving a car, in which you or others would be put at risk by hypoglycemia.

High blood sugar (hyperglycemia) and diabetic ketoacidosis

Hyperglycemia (too much sugar in the blood) means that the levels of glucose in your body are too high. Hyperglycemia can be brought about by:

- not taking your insulin;
- using less insulin than your body needs;
- an imbalance between the amount of carbohydrates you eat and the insulin you take; or
- fever, infection or emotional stress.

The early symptoms of hyperglycemia are:

- being very thirsty
- headache
- feeling sleepy
- urinating more often

Hyperglycemia can lead to diabetic ketoacidosis. The first symptoms come on slowly over many hours or days. Additional symptoms include the following:

- nausea and/or vomiting
- abdominal pain
- rapid pulse
- heavy breathing
- moderate or large amounts of urine ketones. Ketones are produced when your body burns fat for energy instead of glucose.

If you have any of these symptoms and high sugars **get medical help immediately**. See section 'If you forget to take the medicine'.

Illness

If you are ill, especially if you feel sick (nauseous) or are sick (vomiting), the amount of insulin you need may change. **Even when you are not eating normally, you still need insulin.** Test your urine or blood, follow your 'sick rules', and tell your doctor.

5. HOW TO STORE THE MEDICINE?

Avoid poisoning! This medicine and any other medicine should be kept in a closed place out of the reach and sight of children and/or infants in order to avoid poisoning. Do not induce vomiting without an explicit instruction from the doctor.

Do not use the medicine after the expiry date (exp. date) which is stated on the label and the carton. The expiry date refers to the last day of that month.

Storage conditions

Do not freeze.

Keep in the outer carton in order to protect from light.

Before first use

Store in a refrigerator (2°C to 8°C).

After first use

Do not store above 30°C.

Do not freeze.

Discard after 28 days **even if some of the solution remains**.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help to protect the environment.

6. ADDITIONAL INFORMATION

In addition to the active ingredient, the medicine also contains:

glycerol, sodium citrate dihydrate, metacresol, magnesium chloride hexahydrate, zinc oxide, treprostinil sodium, hydrochloric acid and sodium hydroxide (for PH adjustment), water for injections.

What does the medicine look like and what are the contents of the pack:

Lyumjev 100 units/ml, solution for injection is a clear, colorless, aqueous solution in a vial. Each vial contains 1,000 units (10 milliliters). Pack sizes of 1, 2 or a multipack of 5 × 1 vials.

A syringe is not included in the package of the vial.

Not all pack sizes may be marketed.

Registration holder name and address:

Eli Lilly Israel Ltd., 4 HaSheizaf st., POB 4246 Ra'anana 4366411

Manufacturer name and address:

Eli Lilly & Company Ltd., Indianapolis, Indiana, USA

Revised in March 2022 according to MOH guidelines.

Registration number of the medicine in the National Drug Registry of the Ministry of Health:

169-13-36423-00

I LYUMVL I 02

I LYUMVL I 02