

פברואר 2023

רופא/ה נכבד/ה,  
רוקח/ת נכבד/ה,

### Fluconazole MBI 2mg/ml

חברת אמ.בי.איי פארמה בע"מ מבקשת להודיע על עדכונים בעלון לרופא של התכשיר שבנדון.

#### התווית התכשיר:

Fluconazole is indicated in the following fungal infections.

Fluconazole is indicated in adults for the treatment of:

- Cryptococcal meningitis (see section 4.4).
- Coccidioidomycosis (see section 4.4).
- Invasive candidiasis.
- Mucosal candidiasis (including oropharyngeal candidiasis, oesophageal candidiasis, candiduria and chronic mucocutaneous candidiasis).
- Chronic oral atrophic candidiasis (denture sore mouth) if dental hygiene or topical treatment are insufficient.

Fluconazole is indicated in adults for the prophylaxis of:

- Relapse of cryptococcal meningitis in patients with high risk of recurrence.
- Relapse of oropharyngeal or oesophageal candidiasis in patients infected with HIV who are at high risk of experiencing relapse.
- Prophylaxis of candidal infections in patients with prolonged neutropenia (such as patients with haematological malignancies receiving chemotherapy or patients receiving Haematopoietic Stem Cell Transplantation (see section 5.1)).

Fluconazole is indicated in term newborn infants, infants, toddlers, children and adolescents aged from 0 to 17 years old:

Fluconazole is used for the treatment of mucosal candidiasis (oropharyngeal, oesophageal), invasive candidiasis, cryptococcal meningitis and the prophylaxis of candidal infections in immunocompromised patients. Fluconazole can be used as maintenance therapy to prevent relapse of cryptococcal meningitis in children with high risk of reoccurrence (see section 4.4).

Therapy may be instituted before the results of the cultures and other laboratory studies are known; however, once these results become available, anti-infective therapy should be adjusted accordingly.

Consideration should be given to official guidance on the appropriate use of antifungals.

fluconazole 2mg/ml : **מרכיב פעיל:**

Solution for infusion : **צורת המתן של התכשיר:**

**עדכונים בעלון לרופא נעשו בסעיפים הבאים, כאשר החמרה מסומנת בצהוב:**

#### **4.5 Interaction with other medicinal products and other forms of interaction**

[...]

**Abrocitinib:** Fluconazole (inhibitor of CYP2C19, 2C9, 3A4) increased exposure of abrocitinib active moiety by 155%. If co-administered with fluconazole, adjust the dose of abrocitinib as instructed in the abrocitinib prescribing information.

## 6.5 Special precautions for storage

[...]

From a microbiological point of view, the dilutions should be used immediately. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user and would normally not be longer than 24 hours at 2° C to 8° C, unless dilution has taken place in controlled and validated aseptic conditions.

העלון מפורסם במאגר התרופות שבאתר משרד הבריאות:

<https://data.health.gov.il/drugs/index.html#!/byDrug>

לחילופין, ניתן לקבלו מודפס באמצעות פניה לבעל הרישום, חברת אמ.בי.איי פארמה, ת.ד. 5061, קדימה  
טל. 09-7719003

בברכה,

גלי קיסר

רוקחת ממונה