

Remicade

infliximab

Patient Reminder Card

Patient's name: _____

Doctor's name: _____

Doctor's telephone number: _____

This patient reminder card contains important safety information that you need to be aware of before and during treatment with Remicade.

Show this card to any doctor involved in your treatment.

Please read the Remicade 'Package Leaflet' carefully before you start using this medicine.

Date of Remicade therapy initiation: _____

Current administrations: _____

It is important that you and your doctor record the brand name and batch number of your medicine.

Ask your doctor to record the type and date of last screening(s) for tuberculosis (TB) below:

Test _____ Test _____

Date _____ Date _____

Result: _____ Result: _____

Please make sure you also have a list of all other medicines that you are using with you at any visit to a healthcare professional.

List of allergies: _____

List of other medicines: _____

Infections

Before treatment with Remicade

- Tell your doctor if you have an infection even if it is a very minor one
- It is very important that you tell your doctor if you have ever had TB, or if you have been in close contact with someone who has had TB. Your doctor will test you to see if you have TB. Ask your doctor to record the type and date of your last screening(s) for TB on the card

- Tell your doctor if you have hepatitis B or if you know or suspect you are a carrier of the hepatitis B virus.

During treatment with Remicade

- Tell your doctor straight away if you have signs of an infection. Signs include a fever, feeling tired, (persistent) cough, shortness of breath, weight loss, night sweats, diarrhea, wounds, dental problems, burning when urinating or 'flu-like' signs.

Pregnancy and vaccinations

- In case you have received Remicade while you were pregnant or if you are breast-feeding, it is important that you inform your baby's doctor about it before your baby receives any vaccine. Your

baby should not receive BCG live vaccine (used to prevent tuberculosis) within 12 months after birth unless your baby's doctor recommends otherwise. For all other live vaccines, a six-month waiting is recommended.

Keep this card with you for 4 months after your last dose of Remicade, or in case of pregnancy for 12 months after the birth of your baby. Side effects may occur a long time after your last dose.

Reporting Side Effects

This format and its contents have been updated and approved by the Ministry of Health in January 2023. For more information, see the patient package insert. You can report side effects to the Ministry of Health using the online form for reporting side effects on the Ministry of Health homepage: www.health.gov.il, or by clicking on the link: <https://sideeffects.health.gov.il/> Furthermore, you can report side effects directly to Janssen Israel, by calling: 09-9591111.