

## 1. NAME OF THE MEDICINAL PRODUCT

Veklury® 100 mg powder for concentrate for solution for infusion

## 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each vial contains 100 mg of remdesivir. After reconstitution, each vial contains 5 mg/mL of remdesivir solution.

### Excipients with known effect

Each vial contains 3 g betadex sulfobutyl ether sodium

For the full list of excipients, see section 6.1.

## 3. PHARMACEUTICAL FORM

Powder for concentrate for solution for infusion (powder for concentrate).  
White to off-white to yellow powder.

## 4. CLINICAL PARTICULARS

### 4.1 Therapeutic indications

Veklury is indicated for the treatment of coronavirus disease 2019 (COVID-19) in:

- adults and in adolescents (aged 12 to less than 18 years and weighing at least 40 kg) with pneumonia requiring supplemental oxygen (low- or high-flow oxygen or other non-invasive ventilation at start of treatment) (see section 5.1).

### 4.2 Posology and method of administration

Use of remdesivir is confined to healthcare facilities in which patients can be monitored closely (see section 4.4).

#### Posology

The recommended dosage of remdesivir in adults and adolescents (12 to less than 18 years of age and weighing at least 40 kg) is:

- Day 1 – single loading dose of remdesivir 200 mg given by intravenous infusion
- Day 2 onwards – 100 mg given once daily by intravenous infusion.

The recommended **duration** of treatment is 5 days.

#### Special populations

##### *Elderly*

No dose adjustment of remdesivir is required in patients over the age of 65 years (see sections 5.1 and 5.2).

##### *Renal impairment*

The pharmacokinetics of remdesivir have not been evaluated in patients with renal impairment. Patients with eGFR  $\geq$  30 mL/min have received remdesivir for treatment of COVID-19 with no dose adjustment. Remdesivir should not be used in patients with eGFR  $<$  30 mL/min (see sections 4.4 and 5.2).

### *Hepatic impairment*

The pharmacokinetics of remdesivir have not been evaluated in patients with hepatic impairment. It is not known if dosage adjustment is appropriate in patients with hepatic impairment (see sections 4.4 and 5.2).

### *Paediatric population*

Remdesivir is not indicated for children under the age of 12 years and weighing < 40 kg. The safety and efficacy of remdesivir in children under the age of 12 years and weighing < 40 kg have not yet been established. No data are available.

### Method of administration

For intravenous use.

Remdesivir is for administration by intravenous infusion after reconstitution and further dilution.

It must not be given as an intramuscular (IM) injection.

For instructions on reconstitution and dilution of the medicinal product before administration, see section 6.6.

**Table 1: Recommended rate of infusion – for reconstituted and diluted remdesivir powder for concentrate for solution for infusion**

<b>Infusion Bag Volume</b>	<b>Infusion Time</b>	<b>Rate of Infusion</b>
250 mL	30 min	8.33 mL/min
	60 min	4.17 mL/min
	120 min	2.08 mL/min
100 mL	30 min	3.33 mL/min
	60 min	1.67 mL/min
	120 min	0.83 mL/min

### **4.3 Contraindications**

Hypersensitivity to the active substance(s) or to any of the excipients listed in section 6.1.

### **4.4 Special warnings and precautions for use**

#### Hypersensitivity including infusion-related and anaphylactic reactions

Hypersensitivity reactions including infusion-related and anaphylactic reactions have been observed during and following administration of remdesivir. Signs and symptoms may include hypotension, hypertension, tachycardia, bradycardia, hypoxia, fever, dyspnoea, wheezing, angioedema, rash, nausea, vomiting, diaphoresis, and shivering. Slower infusion rates, with a maximum infusion time of up to 120 minutes, can be considered to potentially prevent these signs and symptoms. Monitor patients for hypersensitivity reactions during and following administration of remdesivir. If signs and symptoms of a clinically significant hypersensitivity reaction occur, immediately discontinue administration of remdesivir and initiate appropriate treatment.

#### Transaminase elevations

Transaminase elevations have been observed in the remdesivir clinical trials, including in healthy volunteers and patients with COVID-19. Liver function should be determined in all patients prior to starting remdesivir and should be monitored while receiving it as clinically appropriate. No clinical studies with remdesivir have been conducted in patients with hepatic impairment. Remdesivir should only be used in patients with hepatic impairment if the potential benefit outweighs the potential risk.

- Remdesivir should not be initiated in patients with alanine aminotransferase (ALT)  $\geq 5$  times the upper limit of normal at baseline
- Remdesivir should be discontinued in patients who develop:
  - ALT  $\geq 5$  times the upper limit of normal during treatment with remdesivir. It may be restarted when ALT is  $< 5$  times the upper limit of normal.
  - OR
  - ALT elevation accompanied by signs or symptoms of liver inflammation or increasing conjugated bilirubin, alkaline phosphatase, or international normalised ratio (INR) (see sections 4.8 and 5.2).

### Renal impairment

In animal studies on rats and monkeys, severe renal toxicity was observed (see section 5.3). The mechanism of this renal toxicity is not fully understood. A relevance for humans cannot be excluded.

All patients should have eGFR determined prior to starting remdesivir and while receiving it as clinically appropriate. Remdesivir should not be used in patients with eGFR  $< 30$  mL/min.

### Risk of reduced antiviral activity when coadministered with chloroquine or hydroxychloroquine

Coadministration of remdesivir and chloroquine phosphate or hydroxychloroquine sulphate is not recommended based on *in vitro* data demonstrating an antagonistic effect of chloroquine on the intracellular metabolic activation and antiviral activity of remdesivir (see sections 4.5 and 5.1)

### Excipients

Veklury contains betadex sulfobutyl ether sodium, which is renally cleared and accumulates in patients with decreased renal function, which may potentially adversely affect renal function. Therefore Veklury should not be used in patients with eGFR  $< 30$  mL/min (see sections 4.2 and 5.2).

## **4.5 Interaction with other medicinal products and other forms of interaction**

### Pharmacodynamic interactions

Due to antagonism observed *in vitro*, concomitant use of remdesivir with chloroquine phosphate or hydroxychloroquine sulphate is not recommended.

### Pharmacokinetic interactions

#### *Effects of other medicinal products on remdesivir*

*In vitro*, remdesivir is a substrate for esterases in plasma and tissue, drug metabolizing enzyme CYP3A4 and is a substrate for Organic Anion Transporting Polypeptides 1B1 (OATP1B1) and P-glycoprotein (P-gp) transporters. GS-704277 (a metabolite of remdesivir) is a substrate for OATP1B1 and OATP1B3.

A drug-drug interaction study was conducted with remdesivir. Table 2 summarises the pharmacokinetic effects of studied drugs on remdesivir and metabolites GS-704277 and GS-441524.

**Table 2: Effect of other drugs on remdesivir and metabolites GS-704277 and GS-441524**

Co-administered Drug Dose (mg)	Interaction Geometric mean change (%)	Recommendation concerning co- administration
Cyclosporin 400 single dose	remdesivir: C <sub>max</sub> ↑49% AUC <sub>inf</sub> ↑89% GS-704277: C <sub>max</sub> ↑151% AUC <sub>inf</sub> ↑197% GS-441524: C <sub>max</sub> ↑17% AUC <sub>inf</sub> ↔ No interactions are expected when co-administering remdesivir with inhibitors of OATP1B1/1B3 and/or P-gp.	No dose adjustment of remdesivir is required when it is co-administered with inhibitors of OATP1B1 and OATP1B3.
Carbamazepine 300 twice daily	remdesivir: C <sub>max</sub> ↓13% AUC <sub>inf</sub> ↓8% GS-704277: C <sub>max</sub> ↔ AUC <sub>inf</sub> ↔ GS-441524: C <sub>max</sub> ↔ AUC <sub>inf</sub> ↓17% No interactions are expected when co-administering remdesivir with strong CYP3A4 inducers or CYP3A4 inhibitors.	No dose adjustment of remdesivir is required when it is co-administered with strong CYP3A4 and/or P-gp inducers.

NOTE: Interaction study conducted in healthy volunteers.

#### *Effects of remdesivir on other medicinal products*

*In vitro*, remdesivir is an inhibitor of CYP3A4, UGT1A1, MATE1, OAT3, OCT1, OATP1B1 and OATP1B3. Until respective clinical data become available, the coadministration of sensitive substrates of these enzymes and/or transporters should be considered with caution. Remdesivir induced CYP1A2 and potentially CYP3A *in vitro*. Co-administration of remdesivir with CYP1A2 or CYP3A4 substrates with narrow therapeutic index may lead to loss of their efficacy.

Dexamethasone is a substrate of CYP3A4 and although remdesivir inhibits CYP3A4, due to remdesivir's rapid clearance after IV administration, remdesivir is unlikely to have a significant effect on dexamethasone exposure.

## 4.6 Fertility, pregnancy and lactation

### Pregnancy

There are no or limited amount of data (less than 300 pregnancy outcomes) from the use of remdesivir in pregnant women. Animal studies do not indicate direct or indirect harmful effects with respect to reproductive toxicity at exposures of the major metabolite of remdesivir that were around human therapeutic exposures (see section 5.3). Remdesivir should not be used during pregnancy unless the clinical condition of the women requires treatment with it.

Women of child-bearing potential have to use effective contraception during treatment.

### Breast-feeding

It is unknown whether remdesivir is excreted in human milk or the effects on the breast-fed infant, or the effects on milk production.

In animal studies, the nucleoside analog metabolite GS-441524 has been detected in the blood of nursing rat pups of mothers given remdesivir. Therefore, excretion of remdesivir and/or metabolites into the milk of lactating animals can be assumed.

Because of the potential for viral transmission to SARS-CoV-2-negative infants and adverse reactions from the drug in breast-feeding infants, a decision must be made whether to discontinue breast-feeding or to discontinue/abstain from remdesivir therapy taking into account the benefit of breast-feeding for the child and the benefit of therapy for the woman.

### Fertility

No human data on the effect of remdesivir on fertility are available. In male rats, there was no effect on mating or fertility with remdesivir treatment. In female rats, however, an impairment of fertility was observed (see section 5.3). The relevance for humans is unknown.

### **4.7 Effects on ability to drive and use machines**

Remdesivir is predicted to have no or negligible influence on these abilities.

### **4.8 Undesirable effects**

#### Summary of the safety profile

The most common adverse reaction in healthy volunteers is increased transaminases (14%). The most common adverse reaction in patients with COVID-19 is nausea (4%).

#### Tabulated summary of adverse reactions

The adverse reactions in Table 3 are listed below by system organ class and frequency. Frequencies are defined as follows: Very common ( $\geq 1/10$ ); common ( $\geq 1/100$  to  $< 1/10$ ); uncommon ( $\geq 1/1,000$  to  $< 1/100$ ); rare ( $\geq 1/10,000$  to  $< 1/1,000$ ); not known (cannot be estimated from the available data).

**Table 3: Tabulated list of adverse reactions**

Frequency	Adverse reaction
<i>Immune system disorders</i>	
Rare	hypersensitivity
Not known	anaphylactic reaction, anaphylactic shock
<i>Nervous system disorders</i>	
Common	headache
<i>Cardiac disorders</i>	
Not known	sinus bradycardia*
<i>Gastrointestinal disorders</i>	
Common	nausea
<i>Hepatobiliary disorders</i>	
Very common	transaminases increased
<i>Skin and subcutaneous tissue disorders</i>	
Common	rash
<i>Investigations</i>	
Very common	prothrombin time prolonged
<i>Injury, poisoning and procedural complications</i>	
Rare	infusion-related reaction

\*Reported in post-marketing, usually normalised within 4 days following last remdesivir administration without additional intervention

#### Description of selected adverse reactions

##### *Transaminases Increased*

In healthy volunteer studies, increases in ALT, aspartate aminotransferase (AST) or both in subjects who received remdesivir were grade 1 (10%) or grade 2 (4%). In a randomised, double-blind, placebo-controlled clinical study of patients with COVID-19 (NIAID ACTT-1), any grade ( $\geq 1.25 \times$  upper limit of normal (ULN)) laboratory abnormalities of increased AST and increased ALT occurred in 33% and 32% of patients, respectively, receiving remdesivir compared with 44% and 43% of patients,

respectively, receiving placebo. Grade  $\geq 3$  ( $\geq 5.0 \times \text{ULN}$ ) laboratory abnormalities of increased AST and increased ALT occurred in 6% and 3% of patients, respectively, receiving remdesivir compared with 8% and 6% of patients, respectively, receiving placebo. In a randomised, open-label multi-centre clinical trial (Study GS-US-540-5773) in hospitalised patients with severe COVID-19 receiving remdesivir for 5 (n=200) or 10 days (n=197), any grade laboratory abnormalities of increased AST and increased ALT occurred in 40% and 42% of patients, respectively, receiving remdesivir. Grade  $\geq 3$  laboratory abnormalities of increased AST and increased ALT both occurred in 7% of patients receiving remdesivir. In a randomised, open-label multi-centre clinical trial (Study GS-US-540-5774) in hospitalised patients with moderate COVID-19 receiving remdesivir for 5 (n=191) or 10 days (n=193) compared to standard of care (n=200), any grade laboratory abnormalities of increased AST and increased ALT occurred in 32% and 33% of patients, respectively, receiving remdesivir, and 33% and 39% of patients, respectively, receiving standard of care. Grade  $\geq 3$  laboratory abnormalities of increased AST and increased ALT occurred in 2% and 3% of patients, respectively, receiving remdesivir and 6% and 8%, respectively, receiving standard of care.

#### *Prothrombin time prolonged*

In a clinical study (NIAID ACTT-1) of patients with COVID-19, the incidence of prolonged prothrombin time or INR (predominantly Grades 1-2) was higher in subjects who received remdesivir compared to placebo, with no difference observed in the incidence of bleeding events between the two groups. Prothrombin time should be monitored while receiving remdesivir as clinically appropriate.

#### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Any suspected adverse events should be reported to the Ministry of Health according to the National Regulation by using an online form <https://sideeffects.health.gov.il>. Additionally, adverse events can be reported to the registration holder by email (DrugSafety.Israel@gilead.com).

## **4.9 Overdose**

Treatment of overdose with remdesivir should consist of general supportive measures including monitoring of vital signs and observation of the clinical status of the patient. There is no specific antidote for overdose with remdesivir.

## **5. PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: Antivirals for systemic use, direct acting antivirals, ATC code: J05AB16

#### Mechanism of action

Remdesivir is an adenosine nucleotide prodrug that is metabolized within host cells to form the pharmacologically active nucleoside triphosphate metabolite. Remdesivir triphosphate acts as an analog of adenosine triphosphate (ATP) and competes with the natural ATP substrate for incorporation into nascent RNA chains by the SARS-CoV-2 RNA-dependent RNA polymerase, which results in delayed chain termination during replication of the viral RNA. As an additional mechanism, remdesivir triphosphate can also inhibit viral RNA synthesis following its incorporation into the template viral RNA as a result of read-through by the viral polymerase that may occur in the presence of higher nucleotide concentrations. When remdesivir nucleotide is present in the viral RNA template, the efficiency of incorporation of the complementary natural nucleotide is compromised, thereby inhibiting viral RNA synthesis.

## Antiviral activity

Remdesivir exhibited *in vitro* activity against a clinical isolate of SARS-CoV-2 in primary human airway epithelial cells with a 50% effective concentration (EC<sub>50</sub>) of 9.9 nM after 48 hours of treatment. Remdesivir inhibited the replication of SARS-CoV-2 in the continuous human lung epithelial cell lines Calu-3 and A549-hACE2 with EC<sub>50</sub> values of 280 nM after 72 hours of treatment and 115 nM after 48 hours of treatment, respectively. The EC<sub>50</sub> values of remdesivir against SARS-CoV-2 in Vero cells were 137 nM at 24 hours and 750 nM at 48 hours post-treatment.

The antiviral activity of remdesivir was antagonised by chloroquine phosphate in a dose-dependent manner when the two drugs were co-incubated at clinically relevant concentrations in HEP-2 cells infected with respiratory syncytial virus (RSV). Higher remdesivir EC<sub>50</sub> values were observed with increasing concentrations of chloroquine phosphate. Increasing concentrations of chloroquine phosphate reduced formation of remdesivir triphosphate in A549-hACE2, HEP-2 and normal human bronchial epithelial cells.

Based on *in vitro* testing, remdesivir retained similar antiviral activity (< 2.5-fold change) against clinical isolates of SARS-CoV-2 variants containing the P323L substitution in the viral polymerase including Alpha (B.1.1.7), Beta (B.1.351), Gamma (P.1), Epsilon (B.1.429), Kappa (B.1.617.1), Lambda (C.37), Iota (B.1.526) and Zeta (P.2) variants compared to earlier lineage SARS-CoV-2 (lineage A) isolates. For the clinical isolates of the Delta (B.1.617.2) and Omicron (B.1.1.529, BA.1, BA.2, BA.2.12.1, BA.4 and BA.5) variants, remdesivir also maintained antiviral activity (< 0.7-fold change) relative to the lineage A SARS-CoV-2 isolates. The antiviral activity of remdesivir against SARS-CoV-2 variants is presented in Table 4.

**Table 4: Remdesivir antiviral activity against clinical isolates of SARS-CoV-2 variants**

SARS-CoV-2 Lineage	Country First Identified	WHO Nomenclature	Key Substitutions	Remdesivir EC <sub>50</sub> (nM)	Fold Change in Susceptibility	Change in Susceptibility
A	USA	-	-	110	1.0	
B.1.1.7	UK	Alpha	P323L	192	1.58	No change <sup>a</sup>
B.1.351	South Africa	Beta	P323L	141	1.19	No change <sup>a</sup>
P.1	Brazil	Gamma	P323L	97	0.82	No change <sup>a</sup>
B.1.617.2	India	Delta	P323L, G671S	70	0.59	No change <sup>a</sup>
B.1.429	USA	Epsilon	P323L	210	1.94	No change <sup>a</sup>
P.2	Brazil	Zeta	P323L	151	1.17	No change <sup>a</sup>
B.1.526	USA	Iota	P323L	258	2.33	No change <sup>a</sup>
B.1.617.1	India	Kappa	P323L	77	0.63	No change <sup>a</sup>
C.37	Peru	Lambda	P323L	175	1.37	No change <sup>a</sup>
B.1.1.529	South Africa	Omicron				
BA.1			P323L	44	0.45	No change <sup>a</sup>
BA.2			P323L	25	0.23	No change <sup>a</sup>
BA.2.12.1			P323L	33	0.20	No change <sup>a</sup>
BA.4			P323L	25	0.15	No change <sup>a</sup>
BA.5			P323L	106	0.66	No change <sup>a</sup>

a Fold-change: < 2.5- is not significant. All variants show no reduction in susceptibility.

## Resistance

### *In Cell Culture*

SARS-CoV-2 isolates with reduced susceptibility to remdesivir have been selected in cell culture. In one selection with GS-441524, the parent nucleoside of remdesivir, virus pools emerged expressing combinations of amino acid substitutions at V166A, N198S, S759A, V792I, C799F, and C799R in the viral RNA-dependent RNA polymerase, conferring EC<sub>50</sub> fold-changes of 2.7 up to 10.4. When individually introduced into a wild-type recombinant virus by site-directed mutagenesis, 1.7- to 3.5-fold reduced susceptibility to remdesivir was observed. In a second selection with remdesivir using a SARS-CoV-2 isolate containing the P323L substitution in the viral polymerase, a single amino acid substitution at V166L emerged. Recombinant viruses with substitutions at P323L alone or

P323L+V166L in combination exhibited 1.3- and 1.5-fold changes in remdesivir susceptibility, respectively.

Cell culture resistance profiling of remdesivir using the rodent CoV murine hepatitis virus identified two substitutions (F476L and V553L) in the viral RNA-dependent RNA polymerase at residues conserved across CoVs that conferred 5.6-fold reduced susceptibility to remdesivir. Introduction of the corresponding substitutions (F480L and V557L) into SARS-CoV resulted in 6-fold reduced susceptibility to remdesivir in cell culture and attenuated SARS-CoV pathogenesis in a mouse model. When individually introduced into a SARS-CoV-2 recombinant virus, the corresponding substitutions at F480L and V557L each conferred 2-fold reduced susceptibility to remdesivir.

#### *In Clinical Trials*

In NIAID ACTT-1 Study (CO-US-540-5776), among 61 patients with baseline and post-baseline sequencing data available, the rate of emerging substitutions in the viral RNA-dependent RNA polymerase was similar in patients treated with VEKLURY compared to placebo. In 2 patients treated with VEKLURY, substitutions in the RNA-dependent RNA polymerase previously identified in resistance selection experiments (V792I or C799F) and associated with low fold change in remdesivir susceptibility ( $\leq 3.4$ -fold) were observed. No other RNA-dependent RNA polymerase substitutions observed in patients treated with VEKLURY were associated with resistance to remdesivir.

In Study GS-US-540-5773, among 19 patients treated with remdesivir who had baseline and post-baseline sequencing data available, substitutions in the viral RNA-dependent RNA polymerase (nsp12) were observed in 4 patients. The substitutions T76I, A526V, A554V and C697F were not associated with resistance to remdesivir ( $\leq 1.45$ -fold change in susceptibility). The effect of substitution E665K on susceptibility to remdesivir could not be determined due to lack of replication.

#### Clinical efficacy and safety

##### *Clinical trials in patients with COVID-19*

##### NIAID ACTT-1 Study (CO-US-540-5776)

A randomised, double-blind, placebo-controlled clinical trial evaluated remdesivir 200 mg once daily for 1 day followed by remdesivir 100 mg once daily for up to 9 days (for a total of up to 10 days of intravenously administered therapy) in hospitalised adult patients with COVID-19 with evidence of lower respiratory tract involvement. The trial enrolled 1,062 hospitalised patients: 159 (15%) patients with mild/moderate disease (15% in both treatment groups) and 903 (85%) patients with severe disease (85% in both treatment groups). Mild/moderate disease was defined as SpO<sub>2</sub> > 94% and respiratory rate < 24 breaths/minute without supplemental oxygen; severe disease was defined as SpO<sub>2</sub>  $\leq$  94% on room air, a respiratory rate  $\geq$  24 breaths/min, and an oxygen requirement, or a requirement for mechanical ventilation. A total of 285 patients (26.8%) (n=131 received remdesivir) were on mechanical ventilation/Extracorporeal Membrane Oxygenation (ECMO). Patients were randomised 1:1, stratified by disease severity at enrolment, to receive remdesivir (n=541) or placebo (n=521), plus standard of care.

The baseline mean age was 59 years and 36% of patients were aged 65 or older. Sixty-four percent were male, 53% were White, 21% were Black, 13% were Asian. The most common comorbidities were hypertension (51%), obesity (45%) and type 2 diabetes mellitus (31%); the distribution of comorbidities was similar between the two treatment groups.

Approximately 38.4% (208/541) of the patients received a 10-day treatment course with remdesivir.

The primary clinical endpoint was time to recovery within 29 days after randomisation, defined as either discharged from hospital (with or without limitations of activity and with or without home oxygen requirements) or hospitalised but not requiring supplemental oxygen and no longer requiring ongoing medical care. The median time to recovery was 10 days in the remdesivir group compared to 15 days in the placebo group (recovery rate ratio 1.29; [95% CI 1.12 to 1.49],  $p < 0.001$ ).



No difference in time to recovery was seen in the stratum of patients with mild-moderate disease at enrolment (n=159). The median time to recovery was 5 days in the remdesivir and 7 days in the placebo groups (recovery rate ratio 1.10; [95% CI 0.8 to 1.53]); the odds of improvement in the ordinal scale in the remdesivir group at Day 15 when compared to the placebo group were as follows: odds ratio, 1.2; [95% CI 0.7 to 2.2, p = 0.562].

Among patients with severe disease at enrolment (n=903), the median time to recovery was 12 days in the remdesivir group compared to 19 days in the placebo group (recovery rate ratio, 1.34; [95% CI 1.14 to 1.58]; p < 0.001); the odds of improvement in the ordinal scale in the remdesivir group at Day 15 when compared to the placebo group were as follows: odds ratio, 1.6; [95% CI 1.3 to 2.0].

Overall, the odds of improvement in the ordinal scale were higher in the remdesivir group at Day 15 when compared to the placebo group (odds ratio, 1.6; [95% CI 1.3 to 1.9], p < 0.001).

The 29-day mortality in the overall population was 11.6% for the remdesivir group vs 15.4% for the placebo group (hazard ratio, 0.73; [95% CI 0.52 to 1.03]; p=0.07). A post-hoc analysis of 29-day mortality by ordinal scale is reported in Table 5.

**Table 5: 29-Day mortality outcomes by ordinal scale<sup>a</sup> at baseline—NIAID ACTT-1 trial**

	Ordinal Score at Baseline			
	5		6	
	Requiring low-flow oxygen		Requiring high-flow oxygen or non-invasive mechanical ventilation	
	Remdesivir (N=232)	Placebo (N=203)	Remdesivir (N=95)	Placebo (N=98)
<b>29-day mortality</b>	4.1	12.8	21.8	20.6
<b>Hazard ratio<sup>b</sup> (95% CI)</b>	0.30 (0.14, 0.64)		1.02 (0.54, 1.91)	

a Not a pre-specified analysis.

b Hazard ratios for baseline ordinal score subgroups are from unstratified Cox proportional hazards models.

#### Study GS-US-540-5773 in Patients with Severe COVID-19

A randomised, open-label multi-centre clinical trial (Study 5773) of patients at least 12 years of age with confirmed SARS-CoV-2 infection, oxygen saturation of ≤ 94% on room air, and radiological evidence of pneumonia compared 200 patients who received remdesivir for 5 days with 197 patients who received remdesivir for 10 days. All patients received 200 mg of remdesivir on Day 1 and 100 mg once daily on subsequent days, plus standard of care. The primary endpoint was clinical status on Day 14 assessed on a 7-point ordinal scale ranging from hospital discharge to increasing levels of oxygen and ventilatory support to death.

The odds of improvement at Day 14 for patients randomized to a 10-day course of remdesivir compared with those randomized to a 5-day course was 0.67 (odds ratio); [95% CI 0.46 to 0.98]. Statistically significant imbalances in baseline clinical status were observed in this study. After adjusting for between-group differences at baseline, the odds of improvement at Day 14 was 0.75 (odds ratio); [95% CI 0.51 to 1.12]. In addition, there were no statistically significant differences in recovery rates or mortality rates in the 5-day and 10-day groups once adjusted for between group differences at baseline. All-cause 28-day mortality was 12% vs 14% in the 5- and 10-day treatment groups, respectively.

#### QT

Current non-clinical and clinical data do not suggest a risk of QT prolongation, but QT prolongation has not been fully evaluated in humans.

#### **5.2 Pharmacokinetic properties**

The pharmacokinetic properties of remdesivir have been investigated in healthy volunteers and patients with COVID-19.

## Absorption

The pharmacokinetic properties of remdesivir and the predominant circulating metabolite GS-441524 have been evaluated in healthy adult subjects. Following intravenous administration of remdesivir adult dosage regimen, peak plasma concentration was observed at end of infusion, regardless of dose level, and declined rapidly thereafter with a half-life of approximately 1 hour. Peak plasma concentrations of GS-441524 were observed at 1.5 to 2.0 hours post start of a 30 minutes infusion.

## Distribution

Remdesivir is approximately 93% bound to human plasma proteins (ex-vivo data) with free fraction ranging from 6.4% to 7.4%. The binding is independent of drug concentration over the range of 1 to 10  $\mu\text{M}$ , with no evidence for saturation of remdesivir binding. After a single 150 mg dose of [ $^{14}\text{C}$ ]-remdesivir in healthy subjects, the blood to plasma ratio of [ $^{14}\text{C}$ ]-radioactivity was approximately 0.68 at 15 minutes from start of infusion, increased over time reaching ratio of 1.0 at 5 hours, indicating differential distribution of remdesivir and its metabolites to plasma or cellular components of blood.

## Biotransformation

Remdesivir is extensively metabolized to the pharmacologically active nucleoside analog triphosphate GS-443902 (formed intracellularly). The metabolic activation pathway involves hydrolysis by esterases, which leads to the formation of the intermediate metabolite, GS-704277. In the liver, carboxylesterase 1 and cathepsin A are the esterases responsible for 80% and 10% of remdesivir metabolism, respectively. Phosphoramidate cleavage followed by phosphorylation forms the active triphosphate, GS-443902. Dephosphorylation of all phosphorylated metabolites can result in the formation of nucleoside metabolite GS-441524 that itself is not efficiently re-phosphorylated. Decyanation of remdesivir and/or its metabolites, followed by subsequent rhodanese mediated conversion generates thiocyanate anion. The levels of thiocyanate detected following administration of 100 mg and 200 mg remdesivir were observed to be significantly below endogenous levels in human plasma.

## Elimination

Following a single 150 mg IV dose of [ $^{14}\text{C}$ ]-remdesivir, mean total recovery of the dose was 92%, consisting of approximately 74% and 18% recovered in urine and feces, respectively. The majority of the remdesivir dose recovered in urine was GS-441524 (49%), while 10% was recovered as remdesivir. These data indicate that renal clearance is the major elimination pathway for GS-441524. The median terminal half-lives of remdesivir and GS-441524 were approximately 1 and 27 hours, respectively.

## Pharmacokinetics of remdesivir and metabolites in adults with COVID-19

Pharmacokinetic exposures for remdesivir and its metabolites in adults with COVID-19 are provided in Table 6.

**Table 6: Multiple dose PK parameters<sup>a</sup> of remdesivir and metabolites (GS-441524 and GS-704277) following IV administration of remdesivir 100 mg to adults with COVID-19**

Parameters Mean <sup>b</sup> (95% CI)	Remdesivir	GS-441524	GS-704277
C <sub>max</sub> (ng/mL)	2700 (2440, 2990)	143 (135, 152)	198 (180, 218)
AUC <sub>tau</sub> (ng•h/mL)	1710 (1480, 1980)	2410 (2250, 2580)	392 (348, 442)
C <sub>tau</sub> (ng/mL)	ND	61.5 (56.5, 66.8)	ND

CI=Confidence Interval; ND=Not detectable (at 24 hours post-dose)

a. Population PK estimates for 30-minute IV infusion of remdesivir for 3 days (Study GS-US-540-9012, n=147).

b. Geometric mean estimates

## Other special populations

### *Gender, race and age*

Based on gender, race and age, pharmacokinetic differences on the exposures of remdesivir were evaluated using population pharmacokinetic analysis. Gender and race did not affect the pharmacokinetics of remdesivir and its metabolites (GS-704277 and GS-441524). Pharmacokinetic exposures of the GS-441524 metabolite were modestly increased in hospitalised COVID-19 patients  $\geq 60$  years of age, however no dose adjustment is needed in these patients.

### *Paediatric patients*

The pharmacokinetics in paediatric patients have not been evaluated.

### *Renal impairment*

The pharmacokinetics of remdesivir and GS-441524 in renal impairment have not been evaluated. Remdesivir is not cleared unchanged in urine to any substantial extent, but its main metabolite GS-441524 is renally cleared and the metabolite levels in plasma may theoretically increase in patients with impaired renal function. The excipient betadex sulfolbutyl ether sodium is renally cleared and accumulates in patients with decreased renal function. Veklury should not be used in patients with  $eGFR < 30$  mL/min.

### *Hepatic impairment*

The pharmacokinetics of remdesivir and GS-441524 in hepatic impairment have not been evaluated. The role of the liver in the metabolism of remdesivir is unknown.

### *Hospitalisation*

Pharmacokinetic exposures for remdesivir in hospitalised patients with severe COVID-19 pneumonia were generally within the range of the exposures in non-hospitalised patients. The GS-704277 and GS-441524 metabolite levels were modestly increased.

### *Interactions*

Remdesivir inhibited CYP3A4 *in vitro* (see section 4.5). At physiologically relevant concentrations (steady-state), remdesivir or its metabolites GS-441524 and GS-704277 did not inhibit CYP1A2, 2B6, 2C8, 2C9, 2C19, and 2D6 *in vitro*. Remdesivir is not a time-dependent inhibitor of CYP450 enzymes *in vitro*.

Remdesivir induced CYP1A2 and potentially CYP3A4, but not CYP2B6 *in vitro* (see section 4.5).

*In vitro* data indicates no clinically relevant inhibition of UGT1A3, 1A4, 1A6, 1A9 or 2B7 by remdesivir or its metabolites GS-441524 and GS-704277. Remdesivir, but not its metabolites, inhibited UGT1A1 *in vitro*.

For GS-441524 and GS-704277, the only enzyme for which metabolism could be detected was UGT1A3.

Remdesivir inhibited OAT3, MATE1, OCT1, OATP1B1 and OATP1B3 *in vitro* (see section 4.5). At physiologically relevant concentrations, remdesivir and its metabolites did not inhibit P-gp and BCRP *in vitro*.

## **5.3 Preclinical safety data**

### Toxicology

Following intravenous administration (slow bolus) of remdesivir to rhesus monkeys and rats, severe renal toxicity occurred after short treatment durations. In male rhesus monkeys at dosage levels of 5, 10, and 20 mg/kg/day for 7 days resulted, at all dose levels, in increased mean urea nitrogen and

increased mean creatinine, renal tubular atrophy, and basophilia and casts, and an unscheduled death of one animal at the 20 mg/kg/day dose level. In rats, dosage levels of >3 mg/kg/day for up to 4 weeks resulted in findings indicative of kidney injury and/or dysfunction. Systemic exposures (AUC) of the predominant circulating metabolite of remdesivir (GS-441524) were 0.1 times (monkeys at 5 mg/kg/day) and 0.3 times (rats at 3 mg/kg/day) the exposure in humans following intravenous administration at the recommended human dose (RHD).

### Carcinogenesis

Long-term animal studies to evaluate the carcinogenic potential of remdesivir have not been performed.

### Mutagenesis

Remdesivir was not genotoxic in a battery of assays, including bacterial mutagenicity, chromosome aberration using human peripheral blood lymphocytes, and *in vivo* rat micronucleus assays.

### Reproductive toxicity

In female rats, decreases in corpora lutea, numbers of implantation sites, and viable embryos, were seen when remdesivir was administered intravenously daily at a systemically toxic dose (10 mg/kg/day) 14 days prior to mating and during conception; exposures of the predominant circulating metabolite (GS-441524) were 1.3 times the exposure in humans at the RHD. There were no effects on female reproductive performance (mating, fertility, and conception) at this dose level.

In rats and rabbits, remdesivir demonstrated no adverse effect on embryofetal development when administered to pregnant animals at systemic exposures (AUC) of the predominant circulating metabolite of remdesivir (GS-441524) that were up to 4 times the exposure in humans at the RHD.

In rats, there were no adverse effects on pre- and post-natal development at systemic exposures (AUC) of the predominant circulating metabolite of remdesivir (GS-441524) that were similar to the exposure in humans at the RHD.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Betadex sulfobutyl ether sodium  
Hydrochloric acid (to adjust pH) (E507)  
Sodium hydroxide (to adjust pH) (E524)

### **6.2 Incompatibilities**

This medicinal product must not be mixed or administered simultaneously with other medicinal products in the same dedicated line except those mentioned in section 6.6.

### **6.3 Shelf life**

#### Unopened vials

The expiry date of the product is indicated on the packaging materials.

#### Reconstituted and diluted solution for infusion

After reconstitution dilute immediately.

Store diluted remdesivir solution for infusion up to 24 hours at room temperature (20°C to 25°C) or 48 hours in a refrigerator (2°C – 8°C). Dilute within the same day as administration.

#### 6.4 Special precautions for storage

No special storage conditions are required. It is recommended to store the medicinal product at room temperature.

For storage conditions after reconstitution and dilution of the medicinal product, see section 6.3.

#### 6.5 Nature and contents of container

Type I clear glass vial, an elastomeric closure, and an aluminium overseal with a flip-off cap.

Pack size: 1 vial

#### 6.6 Special precautions for disposal and other handling

Prepare solution for infusion under aseptic conditions and on the same day as administration. Remdesivir should be inspected visually for particulate matter and discolouration prior to administration, whenever solution and container permit. Should either be observed, the solution should be discarded and fresh solution prepared.

Remdesivir must be reconstituted with 19 mL sterile water for injections and diluted in sodium chloride 9 mg/mL (0.9%) solution for injection before being administered via intravenous infusion over 30 to 120 minutes.

##### Preparation of remdesivir solution for infusion

###### *Reconstitution*

Remove the required number of single-use vial(s) from storage. For each vial:

- Aseptically reconstitute remdesivir powder for concentrate for solution for infusion by addition of 19 mL of sterile water for injections using a suitably sized syringe and needle per vial.
  - Discard the vial if a vacuum does not pull the sterile water for injections into the vial.
- Only use **sterile water** for injection to reconstitute remdesivir powder.
- Immediately shake the vial for 30 seconds.
- Allow the contents of the vial to settle for 2 to 3 minutes. A clear solution should result.
- If the contents of the vial are not completely dissolved, shake the vial again for 30 seconds and allow the contents to settle for 2 to 3 minutes. Repeat this procedure as necessary until the contents of the vial are completely dissolved.
- Inspect the vial to ensure the container closure is free from defects and the solution is free of particulate matter.
- Dilute immediately after reconstitution.

###### *Dilution*

Care should be taken to prevent inadvertent microbial contamination. As there is no preservative or bacteriostatic agent present in this product, aseptic technique must be used in preparation of the final parenteral solution. It is recommended to administer immediately after preparation when possible.

- Using Table 7, determine the volume of sodium chloride 9 mg/mL (0.9%) solution for injection to withdraw from the infusion bag.

**Table 7: Recommended dilution instructions – Reconstituted remdesivir powder for concentrate for solution for infusion**

Remdesivir dose	Sodium chloride 9 mg/mL (0.9%) infusion bag volume to be used	Volume to be withdrawn and discarded from sodium chloride 9 mg/mL (0.9%) infusion bag	Required volume of reconstituted remdesivir
200 mg (2 vials)	250 mL	40 mL	2 × 20 mL
	100 mL	40 mL	2 × 20 mL
100 mg (1 vial)	250 mL	20 mL	20 mL
	100 mL	20 mL	20 mL

NOTE: 100 mL should be reserved for patients with severe fluid restriction, e.g. with ARDS or renal failure.

- Withdraw and discard the required volume of sodium chloride 9 mg/mL from the bag using an appropriately sized syringe and needle per Table 7.
- Withdraw the required volume of reconstituted remdesivir using an appropriately sized syringe per Table 7. Discard any unused portion remaining in the remdesivir vial.
- Transfer the required volume of reconstituted remdesivir to the selected infusion bag.
- Gently invert the bag 20 times to mix the solution in the bag. Do not shake.
- The prepared solution is stable for 24 hours at room temperature (20°C to 25°C) or 48 hours in the refrigerator (2°C to 8°C).

After infusion is complete, flush with at least 30 mL of sodium chloride 9 mg/mL.

### Disposal

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

## **7. MANUFACTURER**

Gilead Sciences Ireland UC  
Carrigtohill  
County Cork, T45 DP77  
Ireland

## **8. REGISTRATION HOLDER**

Gilead Sciences Israel Ltd.  
4 HaHarash Street  
Hod Hasharon,  
4524075  
Israel

## **9. REGISTRATION NUMBER**

165-09-36554-99

Revised in June 2023 in accordance with MoH guidelines

Reference: EU SmPC April 2023

IL-JUN23-EU-APR23