




Information about severe acne

Severe acne is a disease that disfigures the skin. Although acne is usually viewed as a condition of adolescence, acne can continue to a later age, up to the 30s and 40s. Men tend to suffer more from severe acne.

Acne develops in the oil (sebaceous) glands that produce skin oils. These glands can be of different sizes and structure. There is at least one gland in each hair follicle.



These glands secrete an oily substance called sebum, which normally passes through the hair follicle to the skin.

During adolescence these glands grow even further and secrete more than the normal amount of sebum, particularly in the face, chest, and back. Acne develops when the normal passage of sebum to the skin is blocked. In severe acne, the large amount of sebum that builds up in the glands tears the hair follicle wall and starts an inflammatory process under the skin (nodule). These nodules tend to leave scars. There are several factors that may make acne worse, such as stress, cosmetics, and some medicines (such as those containing iodide or bromide).

	Curatane 5 mg capsules	Curatane 10 mg capsules	Curatane 20 mg capsules	Curatane 40 mg capsules
Composition:	Each capsule contains: Isotretinoin 5 mg	Each capsule contains: Isotretinoin 10 mg	Each capsule contains: Isotretinoin 20 mg	Each capsule contains: Isotretinoin 40 mg

This medicine has been prescribed to treat your illness. Do not pass it on to others. It may harm them, even if it seems to you that their medical condition is similar to yours. Treatment with Curatane must be supervised by a dermatologist.

Special warnings for women:

- **Curatane is highly likely to harm an unborn baby (in medical language: the preparation is teratogenic). In addition, it increases the risk of miscarriage, and therefore:**
- **You must not take Curatane when you are pregnant.**
- **You must not take Curatane if you are breastfeeding.** There is a high probability that the medicine will pass into your breast milk and it may harm your baby.
- **You must not take Curatane if there is a chance that you could get pregnant during treatment, or during one month after stopping treatment.**

Women who could get pregnant are allowed to take Curatane under several strict conditions, because of the risk of birth defects (damage to the unborn baby). These are the conditions:

- **You can only use Curatane in cases of severe acne** that has not improved in response to any of the other anti-acne treatments, including antibiotics and skin treatments.
- **Your doctor has explained to you the risk of birth defects.** You understand why you must not get pregnant and what you need to do to prevent becoming pregnant.
- **You discussed contraception with your doctor.** The doctor will provide you with information about preventing pregnancy and may refer you to a specialist who will tailor a contraceptive method for you.
- **You must agree to use one, or preferably two, effective methods of contraception, including condoms, or a diaphragm plus spermicide,** for one month before starting treatment with Curatane, during treatment with Curatane, and for one month after stopping treatment. Before starting treatment with Curatane, your doctor will ask you to perform a pregnancy test, which must be negative in order to take the medicine.
- **You must use contraceptive methods even if you are not menstruating** or are not sexually active (unless the doctor decides this is not necessary).
- **You must be capable of complying with the necessary pregnancy prevention measures as listed in this brochure.**
- **You must agree to come to the doctor for monthly follow up visits** and to perform additional pregnancy tests, as decided by your doctor. You may be asked to perform a pregnancy test 5 weeks after stopping treatment with Curatane. You must not get pregnant during treatment with Curatane and for one month after completing the treatment.

If you nevertheless become pregnant during treatment with Curatane, or during the month after treatment has stopped, stop taking the medicine straight away, and refer to your doctor. Your doctor may refer you to another specialist for consultation.

Prescriptions for women who could get pregnant are limited to 30 days of treatment. A new prescription is necessary to continue treatment.

Each prescription will be valid for 7 days from date of issue.

Special warnings for men:

Curatane does not appear to damage sperm. Very low levels of isotretinoin (the medicine's active ingredient) can be found in the semen of men taking Curatane, but these levels are too low to harm your partner's unborn baby. You must remember not to share your medicine with anyone else, particularly not with women.

X Do not use the medicine if:

- you are sensitive (allergic) to the active ingredient isotretinoin, or to peanuts or soya or any of the other ingredients that this medicine contains. For a list of inactive ingredients, please see section 6: "Additional information" in the patient information leaflet enclosed in the medicine package.
- you are pregnant or think you may be pregnant.
- you are breastfeeding.
- you are able to become pregnant, but cannot or are not willing to follow the guidelines

- for preventing pregnancy that appear in the section “Special warnings for women”.
- you suffer from a liver disease.
 - you have very high levels of blood fats (e.g., high level of cholesterol or triglycerides).
 - you have very high levels of vitamin A in your body (hypervitaminosis A).
 - you are being treated with a tetracycline antibiotic during treatment with Curatane (see the section “Curatane and other medicines”).
 - do not use Curatane in children under the age of 12.

Use of Curatane in children over the age of 12 is only allowed after reaching puberty.
If any of the aforementioned situations apply to you, consult a doctor before taking Curatane.

Special warnings regarding the use of the medicine:

Consult a doctor or pharmacist before taking Curatane.

Warnings for all patients:

- You must tell your doctor if you suffer, or have suffered in the past, from any mental illness (including depression, suicidal behavior or psychosis), or if you are taking any medicines to treat any of these conditions.
- **Severe skin reactions** such as erythema multiforme, Stevens-Johnson syndrome, and toxic epidermal necrolysis have been reported with Curatane use. The rash may develop into widespread blisters or skin peeling. You should also pay attention to ulcers in the mouth, throat, nose, genitals, and conjunctivitis (red and swollen eyes).
- **In rare cases, Curatane may cause severe allergic reactions** some of which can affect the skin in the form of eczema, hives and bruises or red patches on the arms and legs. If you develop an allergic reaction, stop taking Curatane, refer to a doctor urgently and tell him or her that you are taking this medicine.
- **Cut down on intensive exercise and physical activity.** Curatane may cause muscle and joint pain, particularly in children and teenagers engaging in vigorous physical activity.
- **Curatane has been associated with inflammatory bowel disease.** The doctor will stop Curatane treatment if you have severe bloody diarrhea without any history of gastrointestinal disorders.
- **Curatane may cause dry eyes, intolerance to contact lenses and vision difficulties including decreased night vision.** Inform the doctor if you have any of these symptoms. The doctor may ask you to use lubricating eye ointment or artificial tears. If you use contact lenses and you have developed intolerance to contact lenses, you may need to wear glasses during treatment. The doctor may refer you to a specialist for advice if you have vision difficulties, and you may need to stop taking Curatane.
- **Benign intracranial hypertension has been reported with Curatane treatment** in some cases where the medicine Curatane was taken together with tetracyclines (a type of antibiotic). Stop taking Curatane and refer to a doctor urgently if you have symptoms such as: headache, nausea, vomiting, and vision disturbances. The doctor may refer you to a specialist to check for swelling of the optic disk in the eye (papilledema).
- **Curatane may cause an increase in liver enzyme levels.** Your doctor will perform blood tests before, during and after Curatane treatment to measure these levels. If your liver enzyme levels remain high, the doctor may lower the Curatane dosage or decide to stop Curatane treatment.
- **Frequently, Curatane increases blood fat levels,** such as cholesterol or triglycerides. Your doctor will perform blood tests to measure these levels in your blood before, during and after Curatane treatment. It is recommended not to drink alcoholic drinks, or, at least, to reduce the amount of alcoholic drinks you usually consume – for as long as you are being treated with Curatane. Inform your doctor if you are suffering from high blood fat levels, from diabetes (high blood sugar levels), are overweight, or if you suffer from alcohol dependence. You may need to perform blood tests more often. If your blood fat levels remain high, the doctor may lower your Curatane dosage, or decide to stop the treatment.

- **Inform the doctor if you have a kidney problem.** The doctor may start the treatment with a low dosage of Curatane and then increase it to the maximum dose you can tolerate.
- **Tell your doctor if you have a fructose intolerance.** The doctor will not prescribe you with Curatane if you have a fructose, or sorbitol intolerance.
- **Curatane may cause an increase in your blood sugar levels.** In rare cases, patients become diabetic. Your doctor may monitor your blood sugar levels during treatment, particularly if you already have diabetes, are overweight, or if you suffer from alcohol dependence.
- **Curatane can apparently cause dryness of the skin and lips.** It is recommended to use a skin moisturizing ointment or cream, and a lip balm during treatment. To prevent skin irritation, avoid using skin cleansers that peel the skin or anti-acne preparations.
- **Avoid extensive sun exposure and use of sun-lamps and sun-beds.** Your skin may be more sensitive to sunlight. Before you go out in the sun, use a sun protection product with a high protection factor (SPF 15 or higher).
- **Do not undergo cosmetic skin treatments.** Curatane may make your skin more fragile: Do not remove hair by waxing, do not mechanically peel the skin (dermabrasion) and do not perform laser treatments during the course of treatment with Curatane, and for at least 6 months after completing the treatment. These treatments could cause scarring, skin irritation, or in rare cases, changes in the color of the skin.
- **You must remember not to share this medicine with anyone else.** Return extra capsules at the end of the treatment. Consult the doctor or pharmacist regarding where to return the capsules.
- **Do not donate blood** during Curatane treatment and for 30 days after completing the treatment. If a woman who is pregnant receives your blood, the baby may be born with birth defects.

Curatane and other medicines

- If you are taking or have recently taken other medicines, including nonprescription medications and nutritional supplements, tell your doctor or pharmacist.
- Do not take vitamin A supplements or tetracyclines (a type of antibiotic), and do not use skin treatment for acne during treatment with Curatane. Moisturizers or skin emollients (creams or skin preparations that prevent water loss from the skin and have a skin softening effect) can be used.
- During treatment with Curatane, avoid using anti-acne preparations that cause skin peeling.

Pregnancy and breastfeeding

If you are pregnant or breastfeeding, think you may be pregnant or are planning to become pregnant, ask your doctor for advice before taking this medicine.

Pregnancy: Do not use Curatane during pregnancy. If you are able to get pregnant, you should use effective contraceptive methods for one month before starting treatment, during treatment, and for one month after completing treatment with Curatane.

If you do get pregnant during the course of treatment with Curatane, or during the month after completing the treatment, **stop taking the medicine straight away,** and refer to a doctor. The doctor may refer you to a specialist for consultation.

If you used Curatane during pregnancy, it may cause damage to the unborn baby (in medical language: the preparation is teratogenic).

It also increases the risk of miscarriage.

Curatane can cause serious abnormalities of brain, face, ear, eye, heart and certain glands (called the thymus gland and parathyroid gland) of the unborn baby.

Breastfeeding: Do not use Curatane if you are breastfeeding. There is a high probability that the medicine will pass into your breast milk and may harm your baby.

Driving and using machines

During treatment with this medicine, your night vision may be disturbed. This effect may happen suddenly, and in rare cases this effect also continues after stopping the treatment. Drowsiness and dizziness have been reported very rarely. If you experience these effects, do not drive or operate machinery.

How to use this medicine?

Always use according to the doctor's instructions. Check with your doctor or pharmacist if you are not sure.

Only your doctor will determine your dose and how you should take the medicine. Do not exceed the recommended dose.

Do not chew! Swallow the capsules whole, on a full stomach, with some fluids or a small amount of food. The capsules can be taken once or twice a day.

Duration of treatment

Treatment with Curatane usually lasts 16 to 24 weeks. Most patients need only one course of treatment. Your acne may continue to improve for up to 8 weeks after completing the treatment with Curatane, and therefore, an additional course of treatment is not usually started until 8 weeks have passed since completion of treatment.

Sometimes the acne gets worse during the first weeks of treatment. The condition usually improves as treatment progresses. Persist with the treatment as recommended by the doctor. Even if your health improves, do not stop taking this medicine without consulting your doctor.

Side effects

As with any medicine, using Curatane may cause side effects in some people. Do not be alarmed by this list of side effects. You may not experience any of them. The side effects usually wear off gradually or stop after completing the treatment. Other side effects can be serious and you will have to refer to the doctor immediately.

Side effects requiring immediate medical attention:

Skin problems

Side effects occurring at unknown frequency (frequency cannot be determined from available data): Serious skin rashes (erythema multiforme, Stevens-Johnson syndrome, and toxic epidermal necrolysis) that may be life-threatening and therefore require immediate medical treatment. They appear initially as circular patches, often with blisters in their center, usually on the arms and hands, or legs and feet; more severe rashes may also include blistering of the chest and back. Additional symptoms that may appear include eye infection (conjunctivitis) or ulcers in the mouth, throat or nose. Severe forms of rashes may progress to widespread peeling of the skin which may be life-threatening. These serious rashes are often preceded by headache, fever and body aches (flu-like symptoms).

If you develop a severe rash or the skin symptoms described above, stop taking Curatane and consult your doctor immediately.

Mental problems

Rare side effects (appear in 1-10 users out of 10,000):

- Depression or related disorders. Signs of this include a feeling of emptiness or sadness, mood changes, anxiety, crying spells, irritability, loss of ability to enjoy or show interest in social or sport activities, sleeping too much or too little, changes in weight or appetite, reduced performance at school or at work, trouble concentrating.
- Existing depression getting worse.
- Becoming violent or aggressive.

Very rare side effects (appear in less than one user out of 10,000):

- Some patients developed thoughts about hurting themselves or ending their own lives (suicidal thoughts), have tried to end their own lives (attempted suicide), or have ended their lives (committed suicide). These patients may not appear to be depressed.
- Unusual behavior.
- Signs of psychosis: loss of touch with reality, such as hearing voices or seeing things that do not exist.

If you experience signs of any of the above-mentioned mental effects, refer to your doctor immediately.

Your doctor may instruct you to stop taking Curatane. It is possible that stopping treatment with Curatane will not lead to the disappearance of the effects. You may need additional help and your doctor can help you with this.

Allergic reactions

Rare side effects (appear in 1-10 users out of 10,000):

Serious reactions (anaphylactic): difficulty breathing or swallowing, caused by sudden swelling of the throat, face, lips and mouth. Also, sudden swelling of the hands, legs and ankles.

Very rare side effects (appear in less than one user out of 10,000):

- Sudden tightness in the chest, shortness of breath and wheezing, particularly if you have asthma.

If you develop a serious reaction, seek emergency medical help immediately.

If you develop any allergic reaction, **stop taking Curatane and refer to your doctor.**

Bones and muscles

Side effects occurring at unknown frequency (frequency cannot be determined from available data):

Muscle weakness which can be life-threatening, may be associated with difficulty moving arms or legs, painful, swollen, bruised areas of the body, dark-colored urine, reduced or no urine output, confusion or dehydration. These are signs of rhabdomyolysis, a breakdown of muscle tissue which can lead to kidney failure. This may occur if you are doing intensive physical activity during the course of treatment with Curatane.

Liver and kidney problems

Very rare side effects (appear in less than one user out of 10,000):

- Yellow skin or eyes and a feeling of tiredness - these can be signs of jaundice (hepatitis, inflammation of the liver). **Stop taking Curatane straight away and refer to a doctor.**
- Difficulty passing urine, swelling of the eyelids, feeling excessively tired - these can be signs of kidney inflammation. **Stop taking Curatane straight away and refer to a doctor.**

Nervous system problems

Very rare side effects (appear in less than one user out of 10,000):

- Lasting headache, accompanied by nausea, vomiting and changes in vision, including blurred vision - these can be signs of benign intracranial hypertension, especially if Curatane is taken with antibiotics from the tetracycline family. **Stop taking Curatane straight away and refer to a doctor.**

Gut and stomach problems

Very rare side effects (appear in less than one user out of 10,000):

- Severe abdominal pain, with or without severe bloody diarrhea, nausea and vomiting - these can be signs of serious problems in your gut. **Stop taking Curatane straight away and refer to a doctor.**

Eye problems

Very rare side effects (appear in less than one user out of 10,000):

- Blurred vision. **If you develop blurred vision, stop taking Curatane straight away and refer to a doctor.**

If your sight is affected in any way, refer to the doctor as soon as possible.

Additional side effects

Very common side effects (appear in more than one user out of ten):

- Dryness of the skin, especially in the area of the lips and face; skin inflammations; chapped and inflamed lips; rash; itching and slight skin peeling. Use a moisturizing preparation when you start treatment.
- Skin more fragile and redder than usual, especially in the face area.
- Back pain, muscle pain, joint pain, particularly in children and teenagers.

To avoid worsening of bone and muscle problems, cut down on intensive physical activity during the course of treatment with Curatane.

- Inflammation of the eyes (conjunctivitis) and eyelid area; dry and irritated eyes. Consult the pharmacist regarding suitable eye drops. If you suffer from dry eyes and wear contact lenses, you may need to wear glasses instead of contact lenses.
- High liver enzyme levels in blood tests.
- Changed levels of fats in the blood (including HDL or triglycerides).
- Bruising, bleeding or blood clots occurring more readily (in cases in which there is an effect on the blood clotting cells).
- Anemia – weakness, dizziness, pale skin (in cases in which there is an effect on the red blood cells).

There are additional side effects which may occur with Curatane, and that are not listed above. A full description of side effects and their frequencies is available in the patient information leaflet for this medicine.

If you experience any side-effect, if any side effect gets worse, or if you experience a side effect not mentioned in the patient information leaflet, consult your doctor.

Reporting side effects

You can report side effects to the Ministry of Health by clicking on the link: “Reporting side effects due to medical treatment” that is found on the homepage of the Ministry of Health website (www.health.gov.il) which refers to the online form for reporting side effects. You can also use this link:

<https://forms.gov.il/globaldata/getsequence/getsequence.aspx?formType=AdversEffectMedic@moh.gov.il>

Side effects can also be reported directly to Taro by email at drug.safety@taro.co.il or by phone: **1800464664**.

Storage conditions

To prevent poisoning, keep in a closed place out of reach and sight of children and/or infants.

Do not use the medicine after the expiry date (exp. date) that appears on the package.

The expiry date refers to the last day of that month.

Store in the original package, in a dry place below 25°C. Store capsules in the blister tray, in the outer carton, to protect from light and moisture.

This information brochure was reviewed and approved by the Ministry of Health in: November 2016.

