



מרץ 2019

פיזר פי אף אי פרמצבטיקה ישראל בע"מ  
רח' שנקר 9, ת.ד. 12133  
הרצליה פיתוח, ישראל 46725  
טל: 972-9-9700500 פקס: 972-9-9700501  
פיזר פי אף אי פרמצבטיקה ישראל בע"מ

רופא/ה, רוקח/ת נכבד/ה,  
ברצוננו להודיעך על עדכונים בעלון לרופא של התכשיר **Triflucan IV** :  
קו תחתי משמעו תוספת טקסט, קו חוצה משמעו מחיקת טקסט, הדגשה משמעה החמרה.

Solution For Injection

**צורת מינון:**

Each ml contains 2 mg of fluconazole.

**הרכב וחוזק:**

**התוויה:**

Fluconazole is indicated in the following fungal infections.

Fluconazole is indicated in adults for the treatment of:

Cryptococcal meningitis. Coccidioidomycosis . Invasive candidiasis. Mucosal candidiasis (including oropharyngeal candidiasis, oesophageal candidiasis, candiduria and chronic mucocutaneous candidiasis). Chronic oral atrophic candidiasis (denture sore mouth) if dental hygiene or topical treatment are insufficient.

Fluconazole is indicated in adults for the prophylaxis of:

- Relapse of cryptococcal meningitis in patients with high risk of recurrence.
- Relapse of oropharyngeal or oesophageal candidiasis in patients infected with HIV who are at high risk of experiencing relapse.
- Prophylaxis of candidal infections in patients with prolonged neutropenia (such as patients with haematological malignancies receiving chemotherapy or patients receiving Haematopoietic Stem Cell Transplantation).

Fluconazole is indicated in term newborn infants, infants, toddlers, children and adolescents aged from 0 to 17 years old:

Fluconazole is used for the treatment of mucosal candidiasis (oropharyngeal, oesophageal), invasive candidiasis, cryptococcal meningitis and the prophylaxis of candidal infections in immunocompromised patients. Fluconazole can be used as maintenance therapy to prevent relapse of cryptococcal meningitis in children with high risk of reoccurrence.

Therapy may be instituted before the results of the cultures and other laboratory studies are known; however, once these results become available, anti-infective therapy should be adjusted accordingly.

Consideration should be given to official guidance on the appropriate use of antifungals.

### להלן ההחמרות בעלון לרופא:

#### **Interaction with other medicinal products and other forms of interaction**

**Tofacitinib:** Exposure of tofacitinib is increased when tofacitinib is co-administered with medications that result in both moderate inhibition of CYP3A4 and strong inhibition of CYP2C19 (e.g., fluconazole). Therefore, it is recommended to reduce tofacitinib dose to 5 mg once daily when it is combined with these drugs.

העלון לרופא המעודכן נשלח למשרד הבריאות לצורך פרסומו במאגר התרופות שבאתר משרד הבריאות:

<https://www.old.health.gov.il/units/pharmacy/trufot/index.asp?safa=h>

לחילופין, לקבלת עלון מלא מודפס ניתן לפנות לחברת פיזר פי אף אי פרמצבטיקה ישראל בע"מ

שנקר 9, ת.ד. 12133

הרצליה פיתוח, 46725.

בברכה,

גילי קבשה

רוקחת ממונה