

Pharmasin, Veterinary

Granules for solution per – os in drinking water

As per a veterinarian's prescription only

COMPOSITION:

Tylosin (as tartrate) 100%

INDICATIONS:

For prevention and treatment of respiratory diseases in broilers and laying hens. Pharmasin is active against mycoplasma species, which are the primary cause of respiratory diseases.

DOSAGE:

0.5 gram Pharmasin in 1 liter drinking water, every 24 hours, for 3-5 days. Add the entire contents of the package into a container containing 50 liters of water at room temperature and mix until fully dissolved. Afterwards, add to the chicken's drinking water container, up to a volume of 2000 liters (concentration of 0.5 gram/liter).

During treatment, administer this to the flock as the sole source of drinking water. On very hot days, reduce the dosage, due to increased drinking in poultry, as per the veterinarian's instructions.

WITHDRAWAL PERIOD:

In chickens for meat - 2 days, from completion of treatment.
For laying hens, eggs for consumption - 0 days.

WARNINGS:

- Do not use in cases of sensitivity to tylosin.
- Prepare a fresh solution every 24 hours.

For the operator:

- Use gloves.
- The substance may cause sensitivity upon contact with the skin.
- Avoid inhaling and contact with the eyes and skin.
- In case of contact with the skin, rinse the area thoroughly with water.
- Wash your hands after use.
- Keep out of reach of children.

STORAGE:

Store the product in its original package to protect from light, below 25°C.

SHELF LIFE:

After first opening - 3 months. After diluting - 24 hours.

CONTAINS: 1.1 kg.

MANUFACTURER:

Biovet, Petar Rakov 39, Peshtera 4550, Bulgaria.

REGISTRATION HOLDER:

M.P.Vet Ltd., P.O.B. 7004, Petach Tikva 49170.

The format of this leaflet was determined by the Ministry of Health and its content thereof was checked and approved on 19.12.13



FOR VETERINARY USE