Patient package insert in accordance with the Pharmacists' Regulations (Preparations) - 1986

Non-prescription medicine

Septadine Tarhitz

Liquid

Composition:

Active substances and their concentration: Chlorhexidine Gluconate 4%

For the list of the inactive substances: See Section 6 "Additional information". Read the insert carefully to the end before using the medication.

This insert contains abridged information on the medicine. If you have further questions, consult vour doctor or pharmacist.

Use the preparation correctly. Consult your pharmacist if you require additional information.

1. What the medicine is used for ?

This medicine is indicated as hand rub and for the disinfection of skin prior to invasive procedures.

Therapeutic Group - Antiseptic and antibacterial preparations

2. Before using the medicine

X Do not use the medicine:

- If you are sensitive (allergic) to the active substance or to any of the additional ingredients contained in the medicine. For the list of the inactive substances - see Section 6 "Additional information".
- · On the brain tissues, meninges and middle ear.

!! Special warnings concerning use of the medicine:

- Avoid contact with the eves. In case of contact, rinse thoroughly with water.
- If you are sensitive to any food or medicine, inform your physician before using this medicine.

If you are taking, or recently took. other medicines, including over the counter medicines and dietary supplements, inform your doctor or pharmacist.

■ Important information about some of the ingredients of the medicine: Colored liquid raw material in the preparation might stain clothing

3. How to use the medicine?

Ask your doctor or pharmacist if you are not sure.

The generally recommended manner of use is:

As a lotion in surgery: Wet your hands and arms, then add 5 ml of Septadine Tarhitz and rub for 1 minute. Use a nail brush. Repeat this action for another 2 minutes. Wash and dry properly.

For hands and skin washing and disinfection: Wet your hands or skin, then add 5 ml of Septadine Tarhitz and rub for 1 minute. Wash and dry properly.

Do not exceed the recommended dosage. Do not swallow. For external use only.

If you have taken an overdose or if a child has accidentally swallowed the medicine. go immediately to a doctor or

a hospital emergency room and bring the medicine package with you.

Do not take medicines in the dark! Check the label and the dose each time you take vour medicine. Wear glasses if you need them

If you have further questions concerning use of the medicine refer to your doctor or pharmacist.

4. Side Effects:

As for any medicine, the use of Septadine Tarhitz may cause side effects in some users.

Do not be alarmed by the list of side effects. You will probably not suffer from any of them.

Stop treatment and refer to your doctor:

In case of skin sensitivity (rare)

Other Side Effects:

Redness and local irritation

If one of the side effects becomes more serious, or you suffer from a side effect not mentioned in the insert, consult your doctor

Side Effects in Children:

Parents must inform the attending physician of any side effects and of any other medicine being taken by the child. See above for details of special side effects

5. How to store the medicine?

Avoid poisoning!

This medicine, and all other medicine, must be stored in a safe place out of the reach of children and/or infants, in order to avoid poisoning.

Do not induce vomiting unless explicitly instructed to do so by the doctor.

 Do not use the medicine after the expiry date appearing on the package.

The expiry date refers to the last day of that month

- Store in a cool place, below 25°C.
- Close tightly to prevent entry of air and
- Store in the original package.
- You can use the product after opening until expiry of its validity.

6. Additional information

In addition to the active substances the medicine also contains:

Purified Water, Poloxamer, Lauramine Oxide, Isopropyl Alcohol, Ponceau 4R (E124), Fragrance.

Appearance of the medicine and contents of the package:

Clear red liquid.

Contents: 100 ml 500 ml

Manufacturer and Reg. Owner:

Ben-Shimon Floris Ltd., Misgay Industrial Park, D.N. Misgav 2017400.

This leaflet was revised on 12.08.2020

Registration No. of the medicine in the National Drug Register of the Ministry of

Health: 112 44 29484 00